

# CONVERGENCE RATE OF A STABLE, MONOTONE AND CONSISTENT SCHEME FOR THE MONGE-AMPÈRE EQUATION

GERARD AWANOU

ABSTRACT. We prove a rate of convergence for smooth solutions of the Monge-Ampère equation of a stable, monotone and consistent discretization. We consider the Monge-Ampère equation with a small low order perturbation. No rate of convergence was known for the discretization we consider.

## 1. INTRODUCTION

We obtain a rate of convergence, in the case of smooth convex solutions, of the finite difference schemes introduced in [10, 8] for the elliptic Monge-Ampère equation

$$\det D^2u = f \text{ in } \Omega, u = g \text{ on } \partial\Omega. \quad (1.1)$$

Here, for a smooth function  $u$ ,  $D^2u = \left( (\partial^2u)/(\partial x_i \partial x_j) \right)_{i,j=1,\dots,d}$  is the Hessian of  $u$ , a symmetric matrix field. We assume that  $\Omega$  is a bounded convex domain of  $\mathbb{R}^d$ ,  $d \geq 2$ ,  $g \in C(\partial\Omega)$  can be extended to a function  $\tilde{g} \in C(\overline{\Omega})$  which is convex in  $\Omega$  and  $f > 0 \in C(\overline{\Omega})$ . We consider in this paper a finite difference scheme  $F_h(u^h) = 0$  which is stable, monotone and consistent for the perturbed Monge-Ampère equation

$$\det D^2u + \delta u = f \text{ in } \Omega, u = g \text{ on } \partial\Omega. \quad (1.2)$$

Here  $\delta > 0$  is a small parameter and, for the discretizations we consider, both the term  $\delta u$  and its discretization are often omitted by an abuse of notation. The scheme we consider was introduced in [8] but the approach we take also applies to the one introduced in [10]. The local consistency error is  $O(h^2 + d\theta)$  where  $h$  is the spatial resolution and  $d\theta$  the directional resolution. Our global error estimates are in terms of  $O(h + d\theta)$ . The stability of the schemes for smooth solutions is a direct consequence of our error estimates.

Rate of convergence for smooth solutions were previously established in the context of finite elements [4, 7, 5] or the standard finite difference method [1, 2]. The rate of convergence proven in this paper for a stable, monotone and consistent scheme is a key component of the theory developed in [2] for the convergence of finite difference discretizations to the Aleksandrov solution of the Monge-Ampère equation. It follows from the approach taken therein and the results of this paper, that the discretizations proposed in [10, 8] have approximations which converge to the weak solution, as defined in [11], of an approximate problem to (1.2), even in the general case where (1.2) does not have a smooth solution. For convergence results in the classical sense,

---

Gerard Awanou was partially supported by NSF grant DMS-1319640. This version takes into account both the published version and its erratum.

the rate of convergence given here is expected to help establish a rate of convergence for the scheme without a dependence on the smoothness of the solution. Such a result would eliminate the need for convergence to an approximate problem, instead of (1.2), for the theory developed in [2]. See also [3] for a different approach.

## 2. NOTATIONS AND PRELIMINARIES

We make the usual convention of using the letter  $C$  for various constants independent of the discretization. We make the assumption that  $\Omega = (0, 1)^d \subset \mathbb{R}^d$ . Let  $h > 0$  denote the mesh size. We assume without loss of generality that  $1/h \in \mathbb{Z}$ . Put

$$\begin{aligned}\mathbb{Z}_h &= \{x = (x_1, \dots, x_n)^T \in \mathbb{R}^n : x_i/h \in \mathbb{Z}\} \\ \Omega_0^h &= \Omega \cap \mathbb{Z}_h, \Omega^h = \bar{\Omega} \cap \mathbb{Z}_h, \partial\Omega^h = \partial\Omega \cap \mathbb{Z}_h = \Omega^h \setminus \Omega_0^h.\end{aligned}$$

For  $x \in \mathbb{R}^d$ , we denote the maximum norm of  $x$  by  $|x|_\infty = \max_{i=1, \dots, d} |x_i|$ . We will use the notation  $|\cdot|$  for the Euclidean norm.

Let  $\mathcal{M}(\Omega^h)$  denote the set of real valued functions defined on  $\Omega^h$ , i.e. the set of mesh functions. For a subset  $T_h$  of  $\Omega^h$ , and  $v^h \in \mathcal{M}(\Omega^h)$  we define

$$|v^h|_{\infty, T_h} = \max_{x \in T_h} |v^h(x)|_\infty.$$

Let  $v$  be a continuous function on  $\Omega$  and let  $r_h(v)$  denote the unique element of  $\mathcal{M}(\Omega^h)$  defined by

$$r_h(v)(x) = v(x), x \in \Omega^h.$$

We extend the operator  $r_h$  canonically to vector fields and matrix fields. For a function  $g$  defined on  $\partial\Omega$ ,  $r_h(g)$  defines the analogous restriction on  $\partial\Omega^h$ .

We are first interested in discrete versions of (1.1)

$$F_h(u^h)(x) = 0, x \in \Omega_0^h, u^h(x) = r_h(g)(x), x \in \partial\Omega^h, \quad (2.1)$$

where  $F_h$  denotes a finite difference discretization of (1.1).

**2.1. Consistency.** We recall that the consistency error of the scheme is defined as  $|F_h(r_h(u))|_{\infty, \Omega_0^h}$  and that the scheme is consistent if for  $u \in C^2(\Omega)$   $|F_h(r_h(u))|_{\infty, \Omega_0^h} \rightarrow 0$  as  $h \rightarrow 0$ . We will assume that the discretization  $F_h(u^h) = 0$  is consistent. For the scheme we consider, the consistency error is given in terms of the usual spatial resolution  $h$  and the directional resolution.

To introduce the directional resolution, we first note that the determinant of the Hessian of a smooth function is essentially a second order directional derivative. More precisely, if we let  $W$  denote the set of orthogonal bases of  $\mathbb{R}^d$ , we have [8]

$$\det D^2u(x) = \min_{(\nu_1, \dots, \nu_d) \in W} \prod_{i=1}^d \frac{\nu_i^T (D^2u(x)) \nu_i}{|\nu_i|^2},$$

where  $\nu_i^T$  denotes the transpose of  $\nu_i$ .

The local directional resolution at  $x$  is defined as

$$d\theta(x) = \max_{|v|=1} \min_{\alpha^h/x \pm \alpha^h \in \Omega^h} \left| v - \frac{\alpha^h}{|\alpha^h|} \right|.$$

We note that as  $h \rightarrow 0$ ,  $d\theta(x) \rightarrow 0$ . The directional resolution  $d\theta$  is then defined as

$$d\theta = \min_{x \in \Omega_0^h} d\theta(x).$$

The directional resolution can also be defined in terms of the angles between vectors. Here we have followed the approach used in [6].

Let  $x_0 \in \Omega_0^h$  such that

$$d\theta(x_0) = \min_{x \in \Omega_0^h} d\theta(x).$$

We denote by  $W_h$  the set of orthogonal bases of  $\mathbb{R}^d$  such that  $(\alpha_1, \dots, \alpha_d) \in W_h$  if and only if  $x_0 \pm \alpha_i \in \Omega^h, \forall i$ . Let  $v^h$  be a given mesh function and let  $x \in \Omega_0^h$ , such that  $x + \alpha_i \in \Omega^h$  but  $x - \alpha_i \notin \Omega^h$  for some  $i$ . We denote by  $x_{\alpha_i}$  the intersection with  $\partial\Omega$  of the line through  $x$  and  $x + \alpha_i$  closest to  $x$ . We then define  $v^h(x - \alpha_i)$  to be the value obtained by quadratic interpolation of  $v^h(x_{\alpha_i})$ ,  $v^h(x)$  and  $v^h(x + \alpha_i)$ . Similarly, if  $x - \alpha_i \in \Omega^h$  but  $x + \alpha_i \notin \Omega^h$  for some  $i$ , we define  $v^h(x + \alpha_i)$  by quadratic interpolation of  $v^h(x_{\alpha_i})$ ,  $v^h(x)$  and  $v^h(x - \alpha_i)$ , where now  $x_{\alpha_i}$  is the intersection with  $\partial\Omega$  of the line through  $x$  and  $x - \alpha_i$  closest to  $x$ .

We consider the discrete Monge-Ampère operator defined by

$$M[u^h](x) = \inf_{(\alpha_1, \dots, \alpha_d) \in W_h} \prod_{i=1}^d \frac{u^h(x + \alpha_i) - 2u^h(x) + u^h(x - \alpha_i)}{|\alpha_i|^2}. \quad (2.2)$$

The operator  $M[r_h(u)]$  is shown in [8] to be consistent with  $\det D^2u(x)$ . We give here a detailed proof.

By a Taylor series expansion

$$\begin{aligned} r_h(u)(x + \alpha) &= r_h(u)(x) + Dr_h(u)(x) \cdot \alpha + \frac{1}{2} \alpha^T D^2 r_h(u)(x) \alpha \\ &\quad + \frac{1}{6} D(\alpha^T D^2 r_h(u)(x) \alpha) \cdot \alpha + O(|\alpha|^4) \\ r_h(u)(x - \alpha) &= r_h(u)(x) - Dr_h(u)(x) \cdot \alpha + \frac{1}{2} \alpha^T D^2 r_h(u)(x) \alpha \\ &\quad - \frac{1}{6} D(\alpha^T D^2 r_h(u)(x) \alpha) \cdot \alpha + O(|\alpha|^4). \end{aligned}$$

Thus using  $|\alpha_i| = O(h)$  for  $\alpha_i \in \mathbb{R}^d$  such that  $x_0 \pm \alpha_i \in \Omega^h$ , we have

$$\frac{\alpha_i^T D^2 u(x) \alpha_i}{|\alpha_i|^2} = \frac{r_h(u)(x + \alpha_i) - 2r_h(u)(x) + r_h(u)(x - \alpha_i)}{|\alpha_i|^2} + O(h^2). \quad (2.3)$$

Moreover, for  $v \in \mathbb{R}^d$ ,  $|v| = 1$ , and  $\alpha \in \mathbb{R}^d$  with  $x_0 \pm \alpha \in \Omega^h$ , a direction vector closest to  $v$ , i.e.  $\alpha = \arg \min_{\zeta \in \mathbb{R}^d, x_0 \pm \zeta \in \Omega^h} |v - \zeta|/|\zeta|$ , we have

$$\left(v + \frac{\alpha}{|\alpha|}\right)^T D^2 u(x) \left(v - \frac{\alpha}{|\alpha|}\right) = v^T D^2 u(x) v - \frac{\alpha^T D^2 u(x) \alpha}{|\alpha|^2}.$$

By definition of  $\alpha$ ,  $|v - \alpha/|\alpha|| \leq d\theta$ . Let us assume that  $u \in C^4(\Omega)$  (so that second derivatives of  $u$  are locally bounded). Since  $|v + \alpha/|\alpha|| \leq 2$ , for a constant  $C > 0$  we

have  $|(v + \alpha/|\alpha|)^T D^2 u(x)(v - \alpha/|\alpha|)| \leq Cd\theta$ . Thus by (2.3)

$$v^T D^2 u(x)v = \frac{r_h(u)(x + \alpha) - 2r_h(u)(x) + r_h(u)(x - \alpha)}{|\alpha|^2} + O(h^2) + O(d\theta). \quad (2.4)$$

Let  $x \in \Omega_0^h$  and let  $\nu_h = (\nu_1, \dots, \nu_d) \in W_h$  such that

$$M[r_h(u)](x) = \prod_{i=1}^d \frac{r_h(u)(x + \nu_i) - 2r_h(u)(x) + r_h(u)(x - \nu_i)}{|\nu_i|^2}.$$

Thus

$$\begin{aligned} M[r_h(u)](x) &= \prod_{i=1}^d \frac{\nu_i^T D^2 u(x) \nu_i}{|\nu_i|^2} + O(h^2) + O(d\theta) \\ &\geq \det D^2 u(x) + Ch^2 + Cd\theta. \end{aligned}$$

It follows that

$$\lim_{h \rightarrow 0} M[r_h(u)](x) \geq \det D^2 u(x). \quad (2.5)$$

Next, if  $(\nu_1, \dots, \nu_d)$  is a basis of eigenvectors of  $D^2 u(x)$ , we know from [8] that  $\det D^2 u(x) = \prod_{i=1}^d \nu_i^T D^2 u(x) \nu_i / |\nu_i|^2$ . We claim that for each  $h > 0$ , we can find  $(\mu_1, \dots, \mu_d) \in W_h(x)$  such that  $|\nu_i/|\nu_i| - \mu_i/|\mu_i|| \leq d\theta$  for all  $i$ . This follows from the observation that for  $i \neq j$ ,  $\nu_i/|\nu_i|$  is obtained from  $\nu_j/|\nu_j|$  by an orthogonal transformation. We recall that orthogonal transformations preserve inner products and that for  $\mu \in \mathbb{R}^d$  such that  $x \pm \mu \in \Omega_h$ , the image of  $\mu$  by a rotation of angle  $\pi/2$  in a plane spanned by two axis vectors is a vector  $\mu'$  for which  $x \pm \mu' \in \Omega_h$ . Thus having found  $\mu_1$  such that  $|\nu_1/|\nu_1| - \mu_1/|\mu_1|| \leq d\theta$ , the other vectors  $\mu_i, i = 1, \dots, d$  are obtained by orthogonal transformations.

It then follows from (2.4) that

$$\begin{aligned} \det D^2 u(x) &= \prod_{i=1}^d \frac{r_h(u)(x + \mu_i) - 2r_h(u)(x) + r_h(u)(x - \mu_i)}{|\mu_i|^2} + O(h^2) + O(d\theta) \\ &\geq M[r_h(u)](x) + O(h^2) + O(d\theta). \end{aligned}$$

We conclude that

$$\det D^2 u(x) \geq \lim_{h \rightarrow 0} M[r_h(u)](x). \quad (2.6)$$

It follows from (2.5) and (2.6) that consistency holds for  $u \in C^4(\Omega)$ .

We note that because of the use of quadratic interpolation for values at points  $x \pm \alpha$  not on the grid, the consistency error will be globally  $O(h + d\theta)$ . To summarize

$$|M[r_h(u)] - \det D^2 u|_{\infty, \Omega_0^h} \leq C_2(h + d\theta). \quad (2.7)$$

### 3. RATE OF CONVERGENCE

The proof of the rate of convergence is an application of the combined fixed point iterative method used in [7].

To ensure convergence to a convex solution of (1.1), it is natural to use a suitable notion of discrete convexity. We require that at an interior grid point  $x$  and for  $\alpha \in \mathbb{R}^d$  such that  $x_0 \pm \alpha \in \Omega^h$

$$u^h(x + \alpha) - 2u^h(x) + u^h(x - \alpha) \geq 0, \quad (3.1)$$

with the usual assumption of the value of  $u^h(x + \alpha)$  (resp.  $u^h(x - \alpha)$ ) obtained by quadratic interpolation of  $u^h(x_\alpha)$ ,  $u^h(x)$  and  $u^h(x - \alpha)$  (resp.  $u^h(x_\alpha)$ ,  $u^h(x)$  and  $u^h(x + \alpha)$ ) when  $x + \alpha \notin \Omega^h$  (resp.  $x - \alpha \notin \Omega^h$ ). Here we have denoted by  $x_\alpha$  the intersection with  $\partial\Omega$  of the line through  $x$  and  $x - \alpha$  closest to  $x$  (resp. the line through  $x$  and  $x + \alpha$  closest to  $x$ ).

We denote by  $\mathcal{C}(\Omega^h)$  the set of mesh functions which satisfy (3.1).

As with [8], the discrete convexity conditions can be combined with a discretization of the differential operator in a single equation. Recall that  $x^+ = \max(x, 0)$  and define

$$M^+[u^h](x) = \inf_{(\alpha_1, \dots, \alpha_d) \in W_h} \prod_{i=1}^d \max \left( \frac{u^h(x + \alpha_i) - 2u^h(x) + u^h(x - \alpha_i)}{|\alpha_i|^2}, 0 \right).$$

Put

$$F_h(u^h)(x) = M^+[u^h](x) + \delta u_h - r_h(f)(x).$$

In practice the term  $\delta u_h$  is not used. It makes the discretization proper as defined below. And guarantees uniqueness of the discrete solution. For consistency, we are forced to consider the perturbed equation (1.2).

The discrete Monge-Ampère equation is given by

$$\begin{aligned} M^+[u^h](x) + \delta u_h - r_h(f)(x) &= 0, x \in \Omega_0^h \\ u^h(x) &= r_h(g)(x) \text{ on } \partial\Omega^h. \end{aligned} \quad (3.2)$$

Let  $N(x)$  denote the set of points  $x \pm \alpha$ ,  $\alpha \in W_h$  and let  $\#N(x)$  denote the cardinality of the set  $N(x)$ . We note that the discretization takes the form

$$F_h(u^h)(x) \equiv \hat{F}_h(u^h(x), u^h(y) - u^h(x)|_{y \neq x, y \in N(x)}),$$

where for  $x \in \Omega_0^h$ ,  $\hat{F}_h$  is a real valued map defined on  $\mathbb{R} \times \mathbb{R}^{\#N(x)}$ . For convenience we do not write explicitly the dependence of  $\hat{F}_h$  on  $x$ .

A scheme is proper if there is  $\delta > 0$  such that for  $x \in \Omega_0^h$  and for all  $a_0, a_1 \in \mathbb{R}$  and  $b \in \mathbb{R}^{\#N(x)}$ ,  $a_0 \leq a_1$  implies  $\hat{F}_h(a_0, b) - \hat{F}_h(a_1, b) \leq \delta(a_0 - a_1)$ .

Thus our scheme is proper and the constant  $\delta > 0$  can be chosen independently of  $h$ .

The scheme is degenerate elliptic since if it is nondecreasing in each of the variables  $u^h(x)$  and  $u^h(y) - u^h(x)$ ,  $y \in N(x)$ ,  $y \neq x$ .

A scheme  $F_h(u^h) = 0$  is Lipschitz continuous if there is  $K > 0$  such that for all  $x \in \Omega_0^h$  and  $(a_0, b_0), (a_1, b_1) \in \mathbb{R}^{\#N(x)+1}$

$$|\hat{F}_h((a_0, b_0)) - \hat{F}_h((a_1, b_1))| \leq K|(a_0, b_0) - (a_1, b_1)|_\infty.$$

We claim that our scheme is locally Lipschitz continuous with Lipschitz constant  $K = C/h^{2d}$ . For  $(a, b) \in \mathbb{R}^{\#N(x)+1}$ , we can find  $r \in \mathbb{R}^{2d+1}$  such that

$$\hat{F}_h((a, b)) = -f(x) + \delta r_0 + \prod_{i=1}^d \frac{r_i + r_{i+d}}{h^2}.$$

If  $M^+[v^h](x) = 0$  we take  $r_i = 0, i = 1, \dots, 2d$ . Otherwise we have

$$M^+[v^h](x) = \prod_{i=1}^d \frac{v^h(x + \alpha_i) - 2v^h(x) + v^h(x - \alpha_i)}{|\alpha_i|^2},$$

for an orthogonal basis  $(\alpha_1, \dots, \alpha_d)$ . We then take  $r_i = h^2/|\alpha_i|^2(v^h(x + \alpha_i) - v^h(x))$ , and  $r_{i+d} = h^2/|\alpha_i|^2(v^h(x - \alpha_i) - v^h(x))$ . In both cases,  $r_0 = v^h(x)$ .

Since the map  $r \rightarrow \prod_{i=1}^d (r_i + r_{i+d})/h^2$  is multilinear,  $\hat{F}_h$  is locally Lipschitz continuous with Lipschitz constant  $\max(\delta, C/h^{2d})$  which we can take as  $C/h^{2d}$  for  $h$  sufficiently small.

Next we define the mapping

$$S : \mathcal{M}(\Omega^h) \rightarrow \mathcal{M}(\Omega^h), \quad S(v^h)(x) = v^h(x) - \nu F_h(v^h)(x), \quad (3.3)$$

for  $\nu > 0$  and for  $\rho > 0$  the ball

$$B_h(\rho) = \{v^h \in \mathcal{C}(\Omega^h), |v_h - r_h(u)|_{\infty, \Omega_0^h} \leq \rho\}.$$

Note that on  $B_h(\rho)$ , the Lipschitz constant  $K$  depends on  $\rho$ . We will assume that  $\rho \leq 1$  so that  $K$  can be assumed to be independent of  $\rho$ . We have [9, Theorem 7]

**Lemma 3.1.** *There exists a positive constant  $a < 1$  such that for all  $v^h, w^h \in B_h(\rho), \rho \leq 1$ , we have*

$$|S(v^h) - S(w^h)|_{\infty, \Omega_0^h} \leq a|v^h - w^h|_{\infty, \Omega_0^h},$$

for  $C_0 \leq \nu \leq C_1$  where  $C_0$  and  $C_1$  are positive constants.

The proof of [9, Theorem 7] shows that under the assumption  $\nu K < 1$ , the constant  $a$  takes the form  $\max(1 - \nu\delta, \nu K)$ . If necessary, by taking  $\nu$  smaller we may assume that  $\nu K < 1/2$  and  $\nu\delta < 1/2$ . Thus the constant  $a$  takes the form

$$a = 1 - \nu\delta,$$

and since  $K = C/h^{2d}$ , we have  $\nu \leq Ch^{2d}$ .

We can now state the main result of this paper.

**Theorem 3.2.** *For  $u \in C^4(\overline{\Omega})$ , (3.2) has a solution  $u^h$  and we have*

$$|u^h - r_h(u)|_{\infty, \Omega_0^h} \leq \frac{C}{\delta}(h + d\theta),$$

for a constant  $C$  which is a scalar multiple of the maximum of the derivatives of  $u$  up to order 4 on  $\overline{\Omega}$  and  $d$ .

*Proof.* Let us first assume that (3.2) has a solution  $u^h$  which is a fixed point of the mapping  $S$ . Recall that the exact solution  $u$  solves the perturbed Monge-Ampère equation (1.2). We have using (2.7)

$$\begin{aligned}
|u^h - r_h(u)|_{\infty, \Omega_0^h} &= |S(u^h) - r_h(u)|_{\infty, \Omega_0^h} \\
&\leq |S(u^h) - S(r_h(u))|_{\infty, \Omega_0^h} + |S(r_h(u)) - r_h(u)|_{\infty, \Omega_0^h} \\
&\leq a|u^h - r_h(u)|_{\infty, \Omega_0^h} + |S(r_h(u)) - r_h(u)|_{\infty, \Omega_0^h} \\
&= a|u^h - r_h(u)|_{\infty, \Omega_0^h} + \nu|F_h(r_h u)|_{\infty, \Omega_0^h} \\
&= a|u^h - r_h(u)|_{\infty, \Omega_0^h} + \nu|M[r_h(u)] - r_h(\det D^2 u)|_{\infty, \Omega_0^h} \\
&\leq a|u^h - r_h(u)|_{\infty, \Omega_0^h} + C\nu(h + d\theta),
\end{aligned}$$

by the consistency of the scheme, the observation that  $r_h(u)$  is discrete convex and (1.2). The constant  $C$  is a consistency error constant and depends on the maximum of derivatives of  $u$  up to order 4 on  $\bar{\Omega}$  and  $d$ .

We therefore have

$$\begin{aligned}
|u^h - r_h(u)|_{\infty, \Omega_0^h} &\leq C \frac{\nu}{1-a} (h + d\theta) \\
&= \frac{C}{\delta} (h + d\theta).
\end{aligned}$$

To complete the proof we now show that (3.2) has a solution  $u^h$  in  $B_h(\rho)$  with  $\rho = C_2(h + d\theta)/\delta$ . This also gives the convergence rate. We note that for  $h$  sufficiently small, we have  $\rho \leq 1$ . Also,  $\rho = C_2\nu(h + d\theta)/(1-a)$ . As above  $|S(r_h(u)) - r_h(u)|_{\infty, \Omega_0^h} \leq C_2\nu(h + d\theta)$ . We have for  $v^h \in B_h(\rho)$

$$\begin{aligned}
|S(v^h) - r_h(u)|_{\infty, \Omega_0^h} &\leq |S(v^h) - S(r_h(u))|_{\infty, \Omega_0^h} + |S(r_h(u)) - r_h(u)|_{\infty, \Omega_0^h} \\
&\leq a|v^h - r_h(u)|_{\infty, \Omega_0^h} + C_2\nu(h + d\theta) \leq a\rho + C_2\nu(h + d\theta) \\
&= a\rho + (1-a)\rho \leq \rho.
\end{aligned}$$

This shows that  $S$  maps the ball  $B_h(\rho)$  into itself, and by Theorem 3.1 is a strict contraction in  $B_h(\rho)$ . The result then follows from the Banach fixed point theorem.  $\square$

#### ACKNOWLEDGEMENTS

The authors would like to thanks the referees for suggestions to improve the paper and Brittney Froese for many useful discussions. The author was partially supported by NSF grant DMS-1319640.

#### REFERENCES

- [1] Awanou, G.: Iterative methods for  $k$ -Hessian equations (2013). <http://arxiv.org/abs/1406.5366>
- [2] Awanou, G.: On standard finite difference discretizations of the elliptic Monge-Ampère equation (2014). <http://arxiv.org/abs/1311.2812>
- [3] Awanou, G., Awi, R.: Convergence of finite difference schemes to the Aleksandrov solution of the Monge-Ampère equation. *Acta Applicandae Mathematicae* **144**(1), 87–98 (2016). DOI 10.1007/s10440-016-0041-x. URL <http://dx.doi.org/10.1007/s10440-016-0041-x>

- [4] Böhmer, K.: On finite element methods for fully nonlinear elliptic equations of second order. *SIAM J. Numer. Anal.* **46**(3), 1212–1249 (2008)
- [5] Brenner, S.C., Gudi, T., Neilan, M., Sung, L.Y.:  $C^0$  penalty methods for the fully nonlinear Monge-Ampère equation. *Math. Comp.* **80**(276), 1979–1995 (2011)
- [6] Buades, A., Coll, B., Lisani, J.L., Sbert, C.: Conditional image diffusion. *Inverse Probl. Imaging* **1**(4), 593–608 (2007)
- [7] Feng, X., Neilan, M.: Analysis of Galerkin methods for the fully nonlinear Monge-Ampère equation. *J. Sci. Comput.* **47**(3), 303–327 (2011)
- [8] Froese, B., Oberman, A.: Convergent finite difference solvers for viscosity solutions of the elliptic Monge-Ampère equation in dimensions two and higher. *SIAM J. Numer. Anal.* **49**(4), 1692–1714 (2011)
- [9] Oberman, A.M.: Convergent difference schemes for degenerate elliptic and parabolic equations: Hamilton-Jacobi equations and free boundary problems. *SIAM J. Numer. Anal.* **44**(2), 879–895 (electronic) (2006)
- [10] Oberman, A.M.: Wide stencil finite difference schemes for the elliptic Monge-Ampère equation and functions of the eigenvalues of the Hessian. *Discrete Contin. Dyn. Syst. Ser. B* **10**(1), 221–238 (2008)
- [11] Trudinger, N.S., Wang, X.J.: Hessian measures. I. *Topol. Methods Nonlinear Anal.* **10**(2), 225–239 (1997). Dedicated to Olga Ladyzhenskaya

DEPARTMENT OF MATHEMATICS, STATISTICS, AND COMPUTER SCIENCE, M/C 249. UNIVERSITY OF ILLINOIS AT CHICAGO, CHICAGO, IL 60607-7045, USA

*E-mail address:* `awanou@uic.edu`

*URL:* `http://www.math.uic.edu/~awanou`