

Resolution of Singularities in Two Dimensions and the Stability of Integrals

Michael Greenblatt

June 9, 2009

1. Background and Statement of Results

Suppose $S(x, y)$ is a smooth real-valued function on a neighborhood of $(0, 0)$ with $S(0, 0) = 0$. For a small open set U containing the origin and a small $\epsilon > 0$, define $M_{S,U}(\epsilon)$ by

$$M_{S,U}(\epsilon) = |\{(x, y) \in U : |S(x, y)| < \epsilon\}| \quad (1.1)$$

By resolution of singularities (see [G2] for an elementary proof) if $S(x, y)$ is real-analytic then if U is sufficiently small then for some $C_U > 0$ one has

$$M_{S,U}(\epsilon) = C_U \epsilon^j |\ln \epsilon|^p + o(\epsilon^j |\ln \epsilon|^p) \quad (1.2)$$

Here j is a positive rational number and $p = 0$ or 1 . Also using resolution of singularities, it can be shown that (j, p) is independent of U for small enough U . We refer to j as the *growth index* of S and to p as the *multiplicity* of this growth index. If $S(x, y)$ is merely smooth, then it follows from [G1] that (1.2) still holds unless there is a smooth coordinate change fixing the origin after which the bisectrix intersects the Newton polygon of S in the interior of its horizontal ray (see below for the relevant definitions.) It is also true that except in those exceptional situations, C_U is independent of U for small enough U . In the exceptional situations, then we can at least say there is a j such that for small enough U , for some $C_U > 0$ we have $M_{S,U}(\epsilon) < C_U \epsilon^j$, while for $j' > j$ there is no C'_U for which the estimate $M_{S,U}(\epsilon) < C'_U \epsilon^{j'}$ holds. Thus for the smooth situation, we also have a natural definition of the growth index of $S(x, y)$ and its multiplicity.

A natural question to consider is the effect of perturbing $S(x, y)$ on the growth index and its multiplicity. Besides being of intrinsic interest, these questions and their oscillatory integral analogues (described later in this section) are important in the analysis of Fourier transforms of surface-supported measures such as in [IKeM] and [IoSa]. Complex and higher-dimensional analogues of these questions are also connected to various issues in complex geometry.

Since by the implicit function theorem one has $M_{S,U}(\epsilon) \sim |\nabla S(0, 0)|^{-1} \epsilon$ when $\nabla S(0, 0) \neq 0$ and U is sufficiently small, one typically assumes that $S(x, y)$ has a critical point at the origin. That is, one assumes that $S(0, 0) = 0$ and $\nabla S(0, 0) = 0$.

This research was supported in part by NSF grant DMS-0654073

For the analogous problem in one dimension, the growth index is just the reciprocal of the order of vanishing of $S(x)$ at $x = 0$, so a small perturbation of the phase can only result in the growth index staying the same or increasing. A famous example of Varchenko [V] shows that the analogous phenomenon does not hold in three or higher dimensions. Thus a general result can hold only in dimension two. The strongest result in the current literature is the following theorem due to Karpushkin [K3]. Let D_r denote the open disk in \mathbf{R}^2 of radius r centered at the origin, and E_r the open disk in \mathbf{C}^2 of radius r centered at the origin. For a function $f(x, y)$ real-analytic on D_r , let $\tilde{f}(z_1, z_2)$ denote the unique holomorphic extension of $f(x, y)$ to E_r . Then Karpushkin's theorem is:

Theorem: ([K3]) Suppose $S(x, y)$ is real-analytic on D_r satisfying $S(0, 0) = 0$ and $\nabla S(0, 0) = 0$ such that $S(x, y)$ has growth index j with multiplicity p at the origin. Then there is a $\delta > 0$, an $s < r$, and a positive constant C_S depending on $S(x, y)$ such that if $f(x, y)$ is real-analytic on D_s and $\tilde{f}(z_1, z_2)$ extends to a continuous function on \bar{E}_s with $|\tilde{f}(z_1, z_2)| < \delta$ for all $(z_1, z_2) \in \bar{E}_s$ then for $0 < \epsilon < \frac{1}{2}$

$$M_{S+f, D_s}(\epsilon) \leq C_S \epsilon^j |\ln \epsilon|^p \quad (1.3)$$

Thus Karpushkin's theorem shows that for real-analytic functions not only does the growth index and multiplicity either improve or stay the same under a small perturbation, but also one has uniformity in the radius s and in the constant C_S . Karpushkin's proofs involve ideas from singularity theory, in particular the theory of versal deformations which turns arbitrary perturbations of $S(x, y)$ into a number of canonical forms which then may be considered individually.

Another method for dealing with stability of $M_{S,U}(\epsilon)$ was introduced in [PSSt] where the above theorem is proven modulo the logarithmic factors. Their methods are often referred to in this subject as the method of algebraic estimates, which also give partial analogues in higher dimensions (the example of Varchenko shows the full analogues not feasible). Also, in the case of *linear* perturbations of smooth functions, results of the above nature are proven in [IKeM].

The purpose of this paper is to show how the use of resolution of singularities algorithms in two dimensions along with some one-dimensional Van der Corput-type lemmas can be used to give new estimates and theorems for the $M_{S,U}(\epsilon)$ as well as for oscillatory integral analogues. Since these algorithms will apply to all smooth functions, our theorems will hold for all smooth functions as opposed to the earlier real-analytic results of [K1]-[K3] and [PSSt]. We also will make use of the superadapted coordinate systems of [G1] that put functions in certain canonical forms suitable for these problems. They are a refinement of the adapted coordinate systems of [V]. Adapted coordinate systems are also used in [K1]-[K3] and [PSSt].

For our sharpest estimates (Theorems 1.1 and 1.5) our condition on the perturbation function $f(x, y)$ will be that finitely many derivatives of $f(x, y)$ at the origin will be less than some δ which depends on $S(x, y)$. We will get uniformity in the coefficient

C_S of (1.3), but there will not be uniformity in the radius s since such uniformity does not hold in the general smooth case. For a given $f(x, y)$, there will also be finitely many t with $|t| < 1$ such that the case when $tf(x, y)$ is the perturbation function is excluded from the theorems. This is due to certain error terms being affected by the zeroes of certain one-dimensional polynomials induced by $S(x, y) + tf(x, y)$. Since these issues only affect error terms, other than for these exceptional values we will get the uniform estimates.

Before we state our theorems we first define some relevant terminology.

Definition 1.1. Let $S(x, y) = \sum_{a,b} s_{ab}x^a y^b$ denote the Taylor expansion of $S(x, y)$ at the origin. Assume there is at least one (a, b) for which s_{ab} is nonzero. For any (a, b) for which $s_{ab} \neq 0$, let Q_{ab} be the quadrant $\{(x, y) \in \mathbf{R}^2 : x \geq a, y \geq b\}$. Then the *Newton polygon* $N(S)$ of $S(x, y)$ is defined to be the convex hull of the union of all Q_{ab} .

A Newton polygon consists of finitely many (possibly zero) bounded edges of negative slope as well as an unbounded vertical ray and an unbounded horizontal ray. More generally, one can define the Newton polygon of a power series in $x^{\frac{1}{N}}$ and $y^{\frac{1}{N}}$ for a positive integer N analogously to Definition 1.1.

Definition 1.2. The *Newton distance* $d(S)$ of $S(x, y)$ is defined to be $\inf\{t : (t, t) \in N(S)\}$.

Throughout this paper, we will use the (a, b) coordinates to write equations of lines relating to Newton polygons, so as to distinguish from the x - y variables of the domain of $S(x, y)$. The line in the a - b plane with equation $a = b$ comes up so frequently it has its own name:

Definition 1.3. The *bisectrix* is the line in the a - b plane with equation $a = b$.

In Theorems 1.1-1.3 below, $S(x, y)$ is a smooth function on a neighborhood of the origin with nonvanishing Taylor expansion at the origin and satisfying $S(0, 0) = 0$ and $\nabla S(0, 0) = 0$. Denote the growth index of $S(x, y)$ by j and its multiplicity by p . Our first and sharpest theorem is the following.

Theorem 1.1. There is a positive integer l and a $\delta > 0$ such that if $f(x, y)$ is a smooth function on a neighborhood of the origin with $\sup_{|\alpha| \leq l} |\partial^\alpha f(0, 0)| < \delta$, then for all but finitely many t with $|t| < 1$, if D is a sufficiently small disk centered at the origin (depending on $S + tf$), then for all $0 < \epsilon < \frac{1}{2}$ we have

$$M_{S+tf, D}(\epsilon) \leq C_S \epsilon^j |\ln \epsilon|^p \quad (1.4)$$

There is no uniformity in the radius of D in Theorem 1.1 as such a statement is false for general smooth functions. It should also be pointed out that since in most cases the leading coefficient of (1.2) is independent of U for small enough U , simply shrinking down D does not typically help in getting a uniform constant in the right-hand side of (1.4).

Our next result says that as long as $S(x, y)$ and $f(x, y)$ do not both have Morse (nondegenerate) critical points at the origin, then Theorem 1.1 still holds for $|t| \geq 1$ as well, although now the constant now depends on t and f as well as S . Another way of saying this is that the growth index and multiplicity of $S(x, y) + tf(x, y)$ is at least as good as that of $S(x, y)$ for all but finitely many t .

Theorem 1.2. Let $f(x, y)$ be a smooth function on a neighborhood of the origin with a critical point there and assume that $S(x, y)$ and $f(x, y)$ do not both have Morse critical points at the origin. Let j_t denote the growth index of $S(x, y) + tf(x, y)$ at the origin and p_t its multiplicity. Then for all but finitely many real values of t we have $(-j_t, p_t) \leq (-j, p)$ (under the lexicographic ordering).

Theorem 1.2 does not hold if $S(x, y)$ and $f(x, y)$ are both Morse as can be seen by taking $S(x, y) = x^2 + y^2$ and $f(x, y) = x^2 - y^2$. However, in such situations the growth index of all but finitely many $S + tf$ is still going to be 1. Also, note that the condition excluding finitely many t may be necessary; for example, when $f(x, y) = -S(x, y)$ plus a small error term.

Since the growth index and multiplicity of $\alpha S_1(x, y) + \beta S_2(x, y)$ is the same as that of $S_1(x, y) + \frac{\beta}{\alpha} S_2(x, y)$ for any $\alpha \neq 0$, Theorem 1.2 and symmetry imply the following:

Theorem 1.3. Suppose $S_1(x, y)$ and $S_2(x, y)$ are two smooth functions on a neighborhood of the origin with critical points at the origin, not both Morse. Let (j_1, p_1) and (j_2, p_2) be their growth indices and multiplicities. Let $j_{\alpha, \beta}$ and $p_{\alpha, \beta}$ be the growth index and multiplicity of $\alpha S_1(x, y) + \beta S_2(x, y)$. Then there is a finite set of real numbers such that unless $\frac{\beta}{\alpha}$ is in this set, $(-j_{\alpha, \beta}, p_{\alpha, \beta}) \leq \min((-j_1, p_1), (-j_2, p_2))$ under the lexicographic ordering.

One includes ∞ as a possible value of $\frac{\beta}{\alpha}$ in Theorem 1.3. Also, note that one can make appropriate generalizations of Theorem 1.3 for several functions.

We now give an idea of how Theorems 1.1 and 1.2 are proved. First consider the simple case where D is a small disk centered at the origin and $S(x, y)$ and $f(x, y)$ are monomials $a_1 x^{\alpha_1} y^{\beta_1}$ and $a_2 x^{\alpha_2} y^{\beta_2}$. Then using an elementary argument, one can evaluate $M_{S+tf, D}(\epsilon)$ directly to show Theorems 1.1 and 1.2. Already one might have to exclude one value of t ; this occurs if $\alpha_1 = \alpha_2$, $\beta_1 = \beta_2$, and $t = -\frac{\alpha_1}{\alpha_2}$. Next, suppose that instead of being monomials, $S(x, y)$ and $f(x, y)$ are comparable to monomials. That is, suppose there are smooth functions $a_1(x, y)$ and $a_2(x, y)$, both nonvanishing at the origin, such that $S(x, y) = a_1(x, y)x^{\alpha_1}y^{\beta_1}$ and $f(x, y) = a_2(x, y)x^{\alpha_2}y^{\beta_2}$. Then roughly speaking one has the same behavior as in the monomial case. There is an added difficulty if $\alpha_1 = \alpha_2$ and $\beta_1 = \beta_2$, for in this case one must also have to consider the zeroes of $a_1(x, y) + ta_2(x, y)$.

More generally, the strong form of resolution of singularities says that in the real-analytic case there is a coordinate change ϕ such that $S \circ \phi(x, y)$ and $f \circ \phi(x, y)$ are locally

comparable to monomials in the above sense. However, the best one can automatically say about the Jacobian of this coordinate change is that it too is comparable to a monomial. Hence when looking at integrals one cannot automatically reduce to the above situations in general. Fortunately, in two dimensions there are substitutes for such resolution of singularities algorithms that reduce to situations similar to where $S \circ \phi$ and $f \circ \phi$ are locally comparable to monomials, and which have determinant one. Even better, these algorithms hold for the general smooth case.

Specifically, in section 3, we will take a small disk D centered at the origin and write $D = \cup_{i=1}^n D_i$. On each D_i there will be a coordinate change ϕ_i such that on each $\phi_i^{-1}D_i$, the function $S \circ \phi_i$ is comparable to a monomial in a certain sense. This can be done in such a way that $f \circ \phi_i$ is also comparable to a monomial, although we won't explicitly use this fact for certain technical reasons. Each $\phi_i(x, y)$ is of the form $(\pm x, \pm y - g_i(x))$, and the domain $\phi_i^{-1}D_i$ is a "curved triangle" consisting of the points in $\phi_i^{-1}(D_i)$ between two curves $y = p_i(x)$ and $y = q_i(x)$ such that $p_i(0) = q_i(0) = 0$ and $p_i(x^N)$ and $q_i(x^N)$ are smooth for some N . Since each ϕ_i now has Jacobian determinant ± 1 , in examining $M_{S,D}(\epsilon)$ one can switch to considering $S \circ \phi_i$ and $f \circ \phi_i$ on the set $\phi_i^{-1}D_i$. Although these two functions aren't strictly speaking comparable to monomials, there are enough similarities with that situation such that after some effort one can prove Theorem 1.2. One has to exclude finitely many values of t for each D_i to avoid cancellations such as in the monomial case.

The idea of dividing into curved triangles related to the singularities of $S(x, y)$ to simplify the behavior of integrals related to $S(x, y)$ goes a while back. It was used in the various Phong-Stein papers on oscillatory integral operators such as [PS] and then in the author's earlier work such as [G3]-[G4]. The Phong-Stein papers use curved triangles deriving from Puiseux expansions of real-analytic functions, while [G3]-[G4] uses explicit resolution of singularities algorithms such as in this paper. Since the problems being considered here are rather different from the earlier problems, we will derive from first principles a resolution of singularities theorem amenable to the situations at hand.

Proving the stronger result of Theorem 1.1 requires additional ideas. In fact, if our goal was only to prove Theorem 1.2 and its consequences, then section 4 would be noticeably shorter. To get the sharper estimates of Theorem 1.1, we will draw on the results of [G1]. Specifically, we first put $S(x, y)$ into what in [G1] are called superadapted coordinates. These are a generalization of the notion of adapted coordinates of [V]. Then we apply the resolution of singularities algorithm of section 3 to $S(x, y)$, getting the resulting D_i . One next focuses on the D_i which give the dominant terms in $M_{S,D}(\epsilon)$. For these D_i , one subdivides further into sets D_{ij} . This will be a coarse subdivision related to the resolution of singularities of $S(x, y) + tf(x, y)$; however, one does not have to use the full resolution of singularities theorem here. For the D_{ij} that give the largest contribution to $M_{S,D}(\epsilon)$, one uses estimates from section 2 related to one-dimensional Van der Corput-type lemmas to prove the sharp estimates of Theorem 1.1.

On the remaining D_{ij} , as well as the D_i that do not give dominant terms of $M_{S,D}(\epsilon)$, one now applies the full resolution of singularities theorem to $f \circ \phi_i(x, y)$. The resulting functions, call them $f \circ \Phi_{ij}$ and $S \circ \Phi_{ij}$, are now comparable to monomials in the new coordinates, and the considerations used for Theorem 1.2 can now be used. Because the contributions here are error terms for $M_{S,D}(\epsilon)$ (that is, they give higher powers of ϵ than the estimates sought), we do not have to worry about constants here if ϵ is small enough, which we will see we can assume. However, we still have to exclude finitely many values of t as in Theorem 1.2; it is conceivable that for such t the power of epsilon appearing in such error terms becomes as small or smaller than the desired power for $M_{S,D}(\epsilon)$. Note that this phenomenon does not appear in the real-analytic results [K1]-[K3] or [PSSt]. The author does not know if this is a result of the weaker assumptions of Theorem 1.1, or if one can avoid excluding finitely many values of t in the context of Theorem 1.1 with an additional argument. As indicated above, this cannot be avoided in the context of Theorem 1.2.

Oscillatory Integrals.

Let $S(x, y)$ be a smooth function on a neighborhood D of the origin with $S(0, 0) = 0$. Suppose $\phi(x, y)$ is a real-valued smooth function supported in D . We consider the oscillatory integral defined by

$$J_{S,\phi}(\lambda) = \int_{\mathbf{R}^2} e^{i\lambda S(x,y)} \phi(x, y) dx dy \quad (1.5)$$

Here λ is a real parameter and we are interested in the behavior of $J_{S,\phi}(\lambda)$ as $|\lambda| \rightarrow \infty$. Since $J_{S,\phi}(-\lambda)$ is the complex conjugate of $J_{S,\phi}(\lambda)$, one only needs to consider the situation as $\lambda \rightarrow \infty$. As is well-known, the analysis of (1.5) is closely related to the analysis of the sublevel areas above. Specifically, in the real-analytic case, if $S(0, 0) = 0$ and $\nabla S(0, 0) = 0$ then in analogy to (1.2) we have nontrivial asymptotics of the form

$$J_{S,\phi}(\lambda) = D_\phi \lambda^{-j} (\ln \lambda)^p + o(\lambda^{-j} (\ln \lambda)^j) \quad (1.6)$$

Here we write (1.6) such that $(-j, p)$ is maximal (under the lexicographic ordering) such that D_ϕ is nonzero for at least one ϕ in any sufficiently small neighborhood of the origin. We refer to j as the oscillatory index of $J_{S,\phi}$ and p as its multiplicity. Using well-known arguments from [AGV], it can be shown that (in the real-analytic case) the oscillatory index is equal to the growth index, with the multiplicity the same in both cases, unless there is a coordinate system near the origin in which $S(x, y) = x^2 - y^2$. In this case, the growth and oscillatory indices are both 1, with the multiplicity of the growth index being 1 and the multiplicity of the oscillatory index being zero.

Furthermore, Karpushkin's methods work for the oscillatory integral case as well, and in [K1]-[K2] analogues for oscillatory integrals (1.6) to his above-mentioned theorem on sublevel areas are proven.

Using some results of [G1], Theorem 1.3 directly implies analogues for oscillatory integrals. To see why this is the case, we first give some background from [G1] which

will also be used in the proof of Theorem 1.1. Suppose $S(x, y)$ is a smooth function on a neighborhood of the origin such that $S(0, 0) = 0$ and $\nabla S(x, y) \neq 0$. Write the Taylor series of $S(x, y)$ as $\sum_{a,b} s_{ab} x^a y^b$ and denote the Newton distance of $S(x, y)$ by d . Then it is proven in [G1] that there is a smooth coordinate change taking the origin to itself, such that after the coordinate change $S(x, y)$ is in "superadapted coordinates", which means the following.

Definition. For a compact edge e of $N(S)$ with equation $a + mb = \alpha$, let $S_e(x, y)$ denote the sum of all terms of $\sum_{a,b} s_{ab} x^a y^b$ with $a + mb = \alpha$. $S(x, y)$ is said to be in *superadapted coordinates* if for any compact edge e of $N(S)$ intersecting the bisectrix, any zero of $S_e(1, y)$ or $S_e(-1, y)$ has order less than $d(S)$.

In [G1] it is proven that in superadapted coordinates, the growth index j is given by $\frac{1}{d}$. The multiplicity p is 1 if and only if the bisectrix intersects $N(S)$ at a vertex. It is also shown that $d = 1$ if $S(x, y)$ is Morse and $d > 1$ otherwise. It is further shown that in superadapted coordinates, on a small enough neighborhood U of the origin for these values of j and p one has $J_{S,\phi}(\lambda) < C\lambda^{-j}(\ln \lambda)^p$ for any ϕ supported in U , and that for any $(-j', p') < (-j, p)$ (with respect to the lexicographic ordering) there is some ϕ supported on U for which the estimate $J_{S,\phi}(\lambda) < C\lambda^{-j'}(\ln \lambda)^{p'}$ does not hold for *any* C .

Thus in the non-Morse case, for a general smooth function it makes sense to define the oscillatory index and multiplicity to be the same as the growth index and multiplicity. In the Morse case one defines them to be inherited from the Morse coordinates. Note that this definition agrees with the old definition for the real-analytic case. Note also that the analogue of Theorem 1.3 for oscillatory integrals follows immediately from Theorem 1.3. Since the two types of Morse critical points have the same oscillatory indices and multiplicities, the case where $S_1(x, y)$ and $S_2(x, y)$ both have Morse critical points at the origin may be included in the oscillatory integral result:

Theorem 1.4. Suppose $S_1(x, y)$ and $S_2(x, y)$ are smooth functions on a neighborhood of the origin with critical points at the origin. Let (j_1, p_1) and (j_2, p_2) be their oscillatory indices and multiplicities at the origin. Let $j_{\alpha,\beta}$ and $p_{\alpha,\beta}$ be the oscillatory index and multiplicity of $\alpha S_1(x, y) + \beta S_2(x, y)$. Then there is a finite set of real numbers such that unless $\frac{\beta}{\alpha}$ is in this set, then $(-j_{\alpha,\beta}, p_{\alpha,\beta}) \leq \min((-j_1, p_1), (-j_2, p_2))$ under the lexicographic ordering.

Again, here we include ∞ as a possible value of $\frac{\beta}{\alpha}$. Note that Theorem 1.4 pertains to integrals of the form $\int_{\mathbb{R}^2} e^{i\alpha S_1(x,y) + i\beta S_2(x,y)} \phi(x, y) dx dy$, which are tied to Fourier transforms of surface-supported measures.

For oscillatory integrals, getting constants depending on $S(x, y)$ and not $f(x, y)$ in analogy with Theorem 1.1 is difficult for a few reasons. For one, since the integrands of oscillatory integrals have both positive and negative values, even if one had precise one-dimensional Van der Corput lemmas for the oscillatory integral case, averaging the

resulting estimates over a second dimension might give extra cancellation that needs to be taken into account. Secondly, and perhaps more importantly, applying one-dimensional Van der Corput lemmas on the integrands of $J_{S,\phi}$ will result in bounds depending on the supremum of $|\nabla\phi|$, and such upper bounds do not necessarily behave well under coordinate changes. However, if the perturbed function $S + tf$ is real-analytic there are explicit formulas from [AGV] for transforming the estimates of Theorem 1.1 into explicit estimates for large enough λ , and we get the following.

Theorem 1.5. Suppose $S(x, y)$, $f(x, y)$, l , δ , j , p , and D are as in Theorem 1.1. There is a $B_S > 0$ such that for all but finitely many $|t| < 1$, if ϕ is supported in D and $S + tf$ is real-analytic, then for sufficiently large λ we have

$$|J_{S+tf,\phi}(\lambda)| < B_S \|\phi\|_\infty \lambda^{-j} (\ln \lambda)^p \quad (1.7)$$

Proof. We may assume $S(x, y)$ is not Morse since the Morse case follows from the explicit asymptotic expansions known in the Morse situation. By Theorem 1.2 of [G1], the initial coefficient D_ϕ of the asymptotics (1.6) for $S + tf$ satisfies

$$|D_\phi| \leq j\Gamma(j) \lim_{\epsilon \rightarrow 0} \left| \frac{I_{S+tf,\phi}(\epsilon)}{\epsilon^j (\ln \epsilon)^p} \right| \quad (1.8)$$

Here $I_{S+tf,\phi}(\epsilon) = \int_{|S+tf| < \epsilon} \phi(x, y) dx dy$. Note that we have

$$\left| \frac{I_{S+tf,\phi}(\epsilon)}{\epsilon^j (\ln \epsilon)^p} \right| \leq \|\phi\|_\infty \left| \frac{M_{S+tf,D}(\epsilon)}{\epsilon^j (\ln \epsilon)^p} \right|$$

Since Theorem 1.1 holds for $S + tf$ on D , the above is at most

$$C_S \|\phi\|_\infty$$

Thus since $j\Gamma(j) \leq 2$, the limit in (1.8) is at most $2C_S \|\phi\|_\infty$. By taking λ sufficiently large that the other terms of the asymptotics (1.6) are small in comparison, we obtain (1.7) with $B_S = 3C_S$ and the theorem follows.

2. Lemmas about curved triangles and one-dimensional Van der Corput Lemmas

We make extensive use of the classical Van der Corput lemma throughout this paper. Although we don't need very sharp constants, to simplify our notation we use the following version that follows from [R]. We refer to [CaCW] for more information on this general subject.

Lemma 2.1. ([R]) Suppose for a positive integer k , $f(t)$ is a C^k function on an interval I such that for some positive constant c , $|\frac{d^k f}{dt^k}| > ck!$ on I . Then for any $\epsilon > 0$ we have

$$|\{t \in I : |f(t)| < \epsilon\}| \leq \min(|I|, 4c^{-\frac{1}{k}} \epsilon^{\frac{1}{k}}) \quad (2.1)$$

An immediate consequence of Lemma 2.1 that will be useful in analyzing our functions on curved triangles is the following.

Lemma 2.2. Let $A_{m,N}$ denote the set $\{(x, y) : 0 < x < x_0, 0 < y < Nx^m\}$, where $m > 0$. Suppose $g(x, y)$ is a function on $A_{m,N}$ such that $|\partial_y^\beta g(x, y)| > a\beta!x^\alpha$ on $A_{m,N}$, where $a, \alpha > 0$ and β is a positive integer. Then for a fixed x we have

$$|\{y : (x, y) \in A_{m,N}, |g(x, y)| < \epsilon\}| \leq 4|\{y : (x, y) \in A_{m,N}, ax^\alpha y^\beta < \epsilon\}| \quad (2.2)$$

Lemma 2.2 will often be used in conjunction with the following lemma.

Lemma 2.3. Let $h(x, y) = ax^\alpha y^\beta$ for some $a, \alpha, \beta \geq 0$. Let $A_{m,N}$ be as in Lemma 2.2.

a) If $\beta > \alpha$, then there are constants $c, C > 0$ depending on a, m, α , and β such that for sufficiently small ϵ we have

$$cx_0^{\frac{\beta-\alpha}{\beta}} \epsilon^{\frac{1}{\beta}} < |\{(x, y) \in A_{m,1} : |h(x, y)| < \epsilon\}| < Cx_0^{\frac{\beta-\alpha}{\beta}} \epsilon^{\frac{1}{\beta}} \quad (2.3)$$

b) If $\beta = \alpha$, then there are constants $c, C > 0$ depending on a, α, m , and β such that for sufficiently small ϵ we have the estimate

$$c\epsilon^{\frac{1}{\beta}} |\ln \epsilon| < |\{(x, y) \in A_{m,1} : |h(x, y)| < \epsilon\}| < C\epsilon^{\frac{1}{\beta}} |\ln \epsilon| \quad (2.4)$$

c) If $\beta < \alpha$, then there are constants $c, C > 0$ depending on a, m, α , and β such that for sufficiently small ϵ we have

$$cN^{\frac{\alpha-\beta}{\alpha+m\beta}} \epsilon^{\frac{m+1}{\alpha+m\beta}} < |\{(x, y) \in A_{m,N} : |h(x, y)| < \epsilon\}| < CN^{\frac{\alpha-\beta}{\alpha+m\beta}} \epsilon^{\frac{m+1}{\alpha+m\beta}} \quad (2.5)$$

Proof: Viewing $|\{(x, y) \in A_{m,N} : |h(x, y)| < \epsilon\}|$ as the integral of the characteristic function of $\{|h(x, y)| < \epsilon\}$ over $A_{m,N}$, we change variables twice, first by replacing y by $x^m y'$ and then by replacing x by $(x')^{\frac{1}{m+1}}$. We obtain that $|\{(x, y) \in A_{m,N} : |h(x, y)| < \epsilon\}|$ is given by

$$(m+1)^{-1} |\{(x', y') \in [0, x_0^{m+1}] \times [0, N] : a(x')^{\frac{\alpha+m\beta}{m+1}} (y')^\beta < \epsilon\}| \quad (2.6)$$

The x' -measure in (2.6) for fixed y' is given by $\min(x_0^{m+1}, c\epsilon^{\frac{m+1}{\alpha+m\beta}} (y')^{-\frac{m\beta+\beta}{\alpha+m\beta}})$. Note that the two terms in the minimum are equal at $y = y'_0 = c'x_0^{-\frac{\alpha+m\beta}{\beta}} \epsilon^{\frac{1}{\beta}}$. Also note that the power $\frac{m\beta+\beta}{\alpha+m\beta}$ of y' is greater than 1 if $\beta > \alpha$, and less than 1 if $\beta < \alpha$. In the former case, if ϵ is sufficiently small the measure of the set (2.6) is comparable to the portion where $y' < y'_0$, given by $y'_0 x_0^{m+1} = c'x_0^{\frac{\beta-\alpha}{\beta}} \epsilon^{\frac{1}{\beta}}$ and thus the formula (2.3) holds. If $\beta = \alpha$, then the exponent is exactly 1, and one obtains the additional logarithmic factor of (2.4). Lastly, if $\beta < \alpha$, the measure of the set (2.6) is comparable to the measure of the part where $N > y' > \frac{N}{2}$, giving (2.5) for small enough ϵ . This completes the proof of the lemma.

We will make frequent use of the next lemma in conjunction with the above lemmas.

Lemma 2.4. Suppose $R(x, y)$ is a smooth function on a neighborhood of $(0, 0)$ such that $N(R)$ has a vertex at (c, d) . Let $r_{cd}x^c y^d$ be the associated term of the Taylor expansion of $R(x, y)$ at $(0, 0)$. Then the following hold.

a) If (c, d) is the intersection of two compact edges of $N(R)$, with equations $a + mb = \alpha$ and $a + m'b = \alpha'$ respectively with $m' > m$, then for any $\delta > 0$ there is an $r > 0$ and a $\xi > 0$ such that for $0 < x < r$ and $\xi^{-1}x^{m'} < y < \xi x^m$ one has

$$|R(x, y) - r_{cd}x^c y^d| < \delta |r_{cd}| x^c y^d \quad (2.7)$$

b) If (c, d) is the intersection of the horizontal ray of $N(R)$ with a compact edge with equation $a + mb = \alpha$, then for sufficiently small $\eta > 0$, for any $\delta > 0$ there is an $r > 0$ and a $\xi > 0$ such that for $0 < x < r$ and $x^{\frac{1}{\eta}} < y < \xi x^m$ equation (2.7) holds.

c) In the setting of case b), if we also have that $d = 0$ then for any $\delta > 0$ there is an $r > 0$ and a $\xi > 0$ such that for $0 < x < r$ and $0 < y < \xi x^m$ equation (2.7) holds.

Proof. We start with a). Let the Taylor expansion of $R(x, y)$ at the origin be written as $\sum_{ab} r_{ab} x^a y^b$. For a large M we may write

$$\begin{aligned} R(x, y) - r_{cd}x^c y^d &= \sum_{c \leq a < M, d \leq b < M, (a,b) \neq (c,d)} r_{cd}x^a y^b \\ &+ \sum_{a < c, d < b < M, a+mb \geq \alpha} r_{ab}x^a y^b + \sum_{c < a < M, b < d, a+m'b \geq \alpha'} r_{ab}x^a y^b + E_M(x, y) \end{aligned} \quad (2.8)$$

Here $E_M(x, y)$ satisfies

$$|E_M(x, y)| < C(|x|^M + |y|^M) \quad (2.9)$$

The first sum in (2.7) can be made less than $\frac{\delta}{4}|r_{cd}|x^c y^d$ in absolute value by shrinking the radius of D appropriately. As for the second sum, if one changes coordinates from (x, y) to (x, y') , where $y' = x^m y$, then $(x, y') \in [0, 1] \times [0, \xi]$. Observe that a given term $r_{ab}x^a y^b$ of the second sum becomes $r_{ab}x^{a+mb}(y')^b$. Since $a + mb \geq \alpha$ and $b > d$ in each term in the second sum, the entire sum can be written as $x^\alpha (y')^d (y' f(x, y'))$ for some $f(x, y')$ which is a polynomial in y' and a fractional power of x . Thus by shrinking ξ appropriately, since $y' < \xi$ the sum can be made of absolute value less than $\frac{\delta}{4}|r_{cd}|x^\alpha (y')^d$. Note that $r_{cd}x^c y^d = r_{cd}x^{c+dm}(y')^d$, and this is equal to $r_{cd}x^\alpha (y')^d$ since (c, d) is on the edge with equation $a + mb = \alpha$. Thus by choosing ξ sufficiently small, we can have that the second sum is of absolute value at most $\frac{\delta}{4}|r_{cd}|x^c y^d$ (in the original coordinates.) These are the bounds we need.

The third sum is dealt with in exactly the same way, reversing the roles of the x and y axes. Lastly, since $x^m > y > x^{m'}$ the error term $E_M(x, y)$ can be made less than

$\frac{\delta}{4}|r_{cd}|x^c y^d$ in absolute value for small x if M is chosen sufficiently large. Putting these all together, we obtain $|R(x, y) - r_{cd}x^c y^d| < \delta|r_{cd}|x^c y^d$ as needed. This completes the proof of part a). of the lemma.

Moving on now to part b), we once again look at the expression (2.8). This time there is no third sum, and all the other terms can be bounded exactly as in part a); the condition that $y > x^{\frac{1}{\eta}}$ for some small η ensures that for large enough M the error term will still be smaller than $\frac{\delta}{4}|r_{cd}|x^c y^d$ using the inequality (2.9). Moving now to c), we again examine (2.8). Once again there is no third sum, and the first two terms are bounded as they were in part a). This time the error term is bounded by $C(|x|^M + |y|^M) \leq C'(|x|^M + |x|^{Mm})$. Since $d = 0$, by taking M large enough the error term can be made less than $\frac{\delta}{4}|r_{cd}|x^c y^d = \frac{\delta}{4}|r_{cd}|x^c$ for small enough x . This gives us part c) and completes the proof of Lemma 2.4.

3. The resolution of singularities algorithm.

In this section we prove the version of two-dimensional resolution of singularities we need for the analysis in section 4. In keeping with the philosophy of [G3] as well as its antecedents such as [PS] or [V], it involves dividing a neighborhood of the origin into "curved triangles" each of which has a coordinate system in which $S(x, y)$ behaves like a monomial in an appropriate sense. The theorem we use is the following.

Theorem 3.1. Suppose $S(x, y)$ is a smooth function defined on a neighborhood of the origin with $S(0, 0) = 0$ such that the Taylor expansion $\sum_{ab} s_{ab}x^a y^b$ of $S(x, y)$ at the origin has at least one nonvanishing term. Then for any sufficiently small $\eta > 0$, and any sufficiently small disk D centered at the origin, we may, up to a set of measure zero, write $D \cap \{(x, y) : |y| < |x|^\eta\}$ as a finite union $\cup_i D_i$, where the D_i have the following properties.

Let M denote the difference between the y coordinates of the uppermost and lowermost vertices of $N(S)$. If $N(S)$ has one vertex let $M = 1$. Then there is a positive integer M' depending on M such that for each i there is a function $g_i(x)$ with $g_i(x^{M'})$ smooth and a function $\phi_i(x, y)$ of the form $(\pm x, \pm y - g_i(x))$ such that $f \circ \phi_i(x, y)$ is approximately a nonzero monomial $b_i x^{\alpha_i} y^{\beta_i}$ ($\alpha_i, \beta_i \geq 0, \beta_i$ an integer) on the curved triangle $D'_i = \phi_i^{-1} D_i$ in the following sense.

a) The set D'_i is of the form $\{(x, y) \in \phi_i^{-1}(D) : x > 0, f_i(x) < y < F_i(x)\}$ where each $f_i(x^{M'})$ and $F_i(x^{M'})$ are smooth. The initial term of the Taylor expansion of $F_i(x)$ is of the form $A_i x^{N_i}$, where $A_i, N_i > 0$. The function $f_i(x)$ is either the zero function or has a Taylor series with initial term $a_i x^{n_i}$ where $a_i, n_i > 0$ and $n_i > N_i$.

b) If $\beta_i = 0$, then $f_i(x)$ is the zero function and there are positive constants c_i and C_i such that on D'_i one has the estimates

$$c_i x^{\alpha_i} < S \circ \phi_i(x, y) < C_i x^{\alpha_i}$$

c) If $\beta_i > 0$, then we can write $S = S_1 + S_2$ as follows. S_2 has a zero of infinite order at

$(0,0)$ and is the zero function if S is real-analytic. Also, $S_2(x^{M'}, y)$ is a smooth function. As for S_1 , for any preselected $\delta > 0$ we can arrange that the decomposition is such that for some nonzero constant b_i , for any $0 \leq k \leq \alpha_i$, $0 \leq l \leq \beta_i$ on D'_i we have

$$\begin{aligned} & |\partial_x^k \partial_y^l S \circ \phi_i(x, y) - b_i \alpha_i (\alpha_i - 1) \dots (\alpha_i - k + 1) \beta_i (\beta_i - 1) \dots (\beta_i - l + 1) x^{\alpha_i - k} y^{\beta_i - l}| \\ & \leq \delta |b_i| x^{\alpha_i - l} y^{\beta_i - k} \end{aligned} \quad (3.1)$$

d) The total number of sets D_i can be bounded in terms of M .

The following corollary will follow immediately from the proof of Theorem 3.1.

Corollary 3.2. If $S(x^p, y)$ is a smooth function for some positive integer p , then Theorem 3.1 still holds, except the exponent M' is replaced by pM' .

Proof of Theorem 3.1. We first dispense with the case where $N(S)$ has exactly one vertex (a, b) . Let $s_{ab}x^a y^b$ be the corresponding term of the Taylor expansion of $S(x, y)$. Thus for any N we have $S(x, y) = s_{ab}x^a y^b + O(|x|^N + |y|^N)$. We divide a small disk D centered at the origin into 8 curved triangles by slicing along the x and y axes as well as along the lines $y = \pm x$. We make these triangles the D_i 's with each $\phi_i(x) = (\pm x, \pm y)$ or $(\pm y, \pm x)$ and each $f_i(x) = 0$. Then condition c) above automatically holds and we are done.

Thus we now assume that $N(S)$ has multiple vertices. Hence $N(S)$ has at least one compact edge. We write the equations of these edges as $a + m_j b = \alpha_j$, where $m_1 > m_2 > \dots > m_n$. Clearly it suffices to divide the $x > 0, y > 0$ portion of a small disk D centered at the origin according to the lemma, so we restrict our attention to $D_0 = D \cap \{x > 0, y > 0\}$. For a small $\xi > 0$ to be determined, up to a set of measure zero we write $D_0 \cap \{y < x^\eta\}$ as the finite union of sets U_j and T_j , where the U_j are all possible sets of the form $\{(x, y) \in D_0 : \xi x^{m_j} < y < \xi^{-i} x^{m_j}\}$ and the T_j are all possible sets of the form $\{(x, y) \in D_0 : \xi^{-i} x^{m_{j+1}} < y < \xi x^{m_j}\}$ as well as the sets $T_n = \{(x, y) \in D_0 : 0 < y < \xi x^{m_n}\}$ and $T_0 = \{(x, y) \in D_0 : \xi^{-i} x^{m_1} < y < x^\eta\}$. (We assume $\eta < m_1$). Observe that each T_j corresponds to a unique vertex of $N(S)$. We will turn each T_j into one of the D_i 's with the associated ϕ_i just the identity map. As for the U_j , we will subdivide each U_j further into V_{jk} and U_{jl} , where the U_{jl} will also become D_i 's for which part b) of the theorem holds, and where the V_{jk} will undergo further subdivisions and coordinate changes.

We start with the T_j 's for $j \neq n$. Let (c, d) denote the vertex of $N(S)$ corresponding to T_j and let $s_{cd}x^c y^d$ denote the associated term of the Taylor expansion. Then by Lemma 2.4 applied to $S(x, y)$ and its various y partials, for any $\delta > 0$ we can choose ξ such that if D is sufficiently small then on T_j , we have the inequality $|S(x, y) - s_{cd}x^c y^d| < \delta |s_{cd}| x^c y^d$ as well as its analogues for any $\partial_x^k \partial_y^l S(x, y)$ for $k \leq c$, $l \leq d$. Thus a) and c) of the theorem hold with ϕ_i the identity map, which is what we need for these T_j .

Next, we look at $T_n = \{(x, y) \in D_0 : 0 < y < \xi x^{m_n}\}$. This time we cannot apply Lemma 2.4 automatically. By a famous theorem of Borel (see [H] for a proof), one can let $s_0(x, y)$ be a smooth function on a neighborhood of the origin whose Taylor expansion at the origin is given by $\sum_{ab} s_{ab} x^a y^{b-d}$. Then Lemma 2.4c) applies to $s_0(x, y)$ and we can assume that on T_n we have

$$|s_0(x, y) - s_{cd} x^c| < \delta |s_{cd} x^c| \quad (3.2)$$

Let $s_1(x, y) = y^d s_0(x, y)$. Then $s_1(x, y)$ has the same Taylor expansion at the origin as $S(x, y)$ and is equal to $S(x, y)$ when $S(x, y)$ is real-analytic. We also have the desired inequality

$$|s_1(x, y) - s_{cd} x^c y^d| < \delta |s_{cd} x^c y^d| \quad (3.3)$$

Note also that the analogues of (3.3) for the x and y partials also hold; for example, the Newton polygon of $s_0(x, y)$ is such that taking a y derivative of $s_0(x, y)$ incurs a factor of at most Cx^{-m_n} which is much smaller than $\frac{1}{y}$ on T_n if ξ is appropriately small. Thus we have a) and c) of Theorem 3.1 and we are done with the analysis of the T_j 's.

Next, we proceed to the analysis of the U_j 's. Let $S_{m_j}(x, y)$ denote the sum of the terms $s_{ab} x^a y^b$ of the Taylor expansion lying on the edge with equation $a + m_j b = \alpha_j$. Note that $S_{m_j}(x, y)$ is a polynomial and is the sum of $s_{ab} x^a y^b$ minimizing $a + m_j b$. Let $r_{j1} < \dots < r_{jN_j}$ denote the positive zeroes of $S_{m_j}(1, y)$ if there are any. Define $V_{jk} = \{(x, y) \in U_j : (r_{jk} - \xi)x^{m_j} < y < (r_{jk} + \xi)x^{m_j}\}$. As long as ξ is small enough, we may write $U_j - \cup_k V_{jk}$ as a union $\cup_l U_{jl}$ where each U_{jl} is of the form $\{(x, y) \in U_j : (r_{j, k+1} - \xi)x^{m_j} < y < (r_{jk} - \xi)x^{m_j}\}$, $\{(x, y) \in U_j : \xi x^{m_j} < y < (r_1 - \xi)x^{m_j}\}$, or $\{(x, y) \in U_j : (r_{N_j} + \xi)x^{m_j} < y < \xi^{-1} x^{m_j}\}$. In the case that $S_{m_j}(1, y)$ has no positive zeroes, there are no V_{kl} 's and we just set $U_{j1} = U_j$.

On each U_{jl} , $S(x, y)$ is already in the form required in Theorem 3.1. To see this, one does the coordinate change $(x, y) = (x, x^{m_j} y')$, turning U_{jl} into a set U'_{jl} contained in $[0, \eta] \times [a_{jl}, b_{jl}]$, where η denotes the radius of the disk D and where $a_{jl}, b_{jl} > 0$. Note that $S_{m_j}(1, y')$ has no zeroes on $[a_{jl}, b_{jl}]$. In the new coordinates, the finite Taylor expansion $S(x, y) = \sum_{a, b < M} s_{ab} x^a y^b + O(|x|^M + |y|^M)$ becomes of the form

$$S(x, x^{m_j} y') = x^{\alpha_j} S_{m_j}(1, y') + x^{\alpha_j + \zeta_j} f(x, y') + O(|x|^M + |x|^{mM} |y'|^M) \quad (3.4)$$

Here $f(x, y')$ is a polynomial in y' and a fractional power of x and $\zeta_j > 0$. Since $S_{m_j}(1, y')$ has no zeroes on $[a_{jl}, b_{jl}]$, there are $C_{jl}, \epsilon_{jl} > 0$ such that $C_{jl} x^{\alpha_j} > |x^{\alpha_j} S_{m_j}(1, y')| > \epsilon_{jl} x^{\alpha_j}$ on U'_{jl} . Furthermore, if η is sufficiently small and M is sufficiently large, we have that $|x^{\alpha_j + \zeta} f(x, y')|$ and the $O(|x|^M + |x|^{mM} |y'|^M)$ terms are both less than $\min(\frac{C_{jl}}{4} x^{\alpha_j}, \frac{\epsilon_{jl}}{4} x^{\alpha_j})$. As a result, shrinking the disk D if necessary we can assume that on U'_{jl} we have

$$\frac{C_{jl}}{2} x^{\alpha_j} > |S(x, x^{m_j} y')| > \frac{\epsilon_{jl}}{2} x^{\alpha_j} \quad (3.5)$$

Translating back into the coordinates of U_{jl} , this means that on U_{jl} we have

$$\frac{C_{jl}}{2} x^{\alpha_j} > |S(x, y)| > \frac{\epsilon_{jl}}{2} x^{\alpha_j} \quad (3.6)$$

Thus on U_{jl} , $S(x, y)$ satisfies the conditions of Theorem 3.1b) with $\beta_i = 0$, if we let the coordinate change ϕ_i associated with U_{jl} be given by $(x, y) \rightarrow (x, y + c_{jl}x^m)$, where $y = c_{jl}x^{m_j}$ denotes the equation of the lower boundary curve of U_{jl} . This completes the analysis of the U_{jl} .

We now move to the analysis of any V_{jk} that may exist. Let o_{jk} denote the order of the zero of $S_{m_j}(1, y)$ at $y = r_{jk}$. The idea is as follows. Using ideas from resolution of singularities, we will do a coordinate change of the form $(x, y) \rightarrow (x, y - r_{jk}x^{m_j} + \text{higher order terms})$ such that in the new coordinates $S(x, y)$ becomes a function whose analogues to the zeroes r_{jk} each has order $< o_{jk}$. Thus after at most $\max_{j,k} o_{jk}$ iterations, there will no longer be any sets V_{jk} and we will have divided D_0 into sets each of which is a U_{jl} or T_j in some iteration. Since each coordinate change will be of the form $(x, y) \rightarrow (x, \pm y - g(x))$, the composition of finitely many such coordinate changes is of the desired form. (We can get $-y$ as well as y since after one of these coordinate changes the resulting set lies in both the upper right and lower right quadrants). By the above analysis of $S(x, y)$ on the T_j and U_{jl} , the resulting $S \circ \phi_i(x, y)$ will satisfy Theorem 3.1 as needed.

The coordinate change on V_{jk} is chosen in the following way. We once again switch to the (x, y') coordinates and make use of (3.4). Let V'_{jk} denote V_{jk} in the new coordinates and define $s_j(x, y') = \frac{S(x, x^{m_j}y')}{x^{\alpha_j}}$, where recall $\alpha_j = a + m_j b$ for (a, b) on e_j . We claim that $s_j(x^N, y')$ is a smooth function of x and y' on V'_{jk} for some N . To see this, observe that by (3.4), for any (p, q) we have

$$\begin{aligned} \partial_x^p \partial_{y'}^q (s_j(x^N, y')) &= \partial_x^p \partial_{y'}^q (S_{m_j}(1, y')) + \partial_x^p \partial_{y'}^q (x^{N\zeta_j} f(x, y')) \\ &+ O(|x|^{N(M-p)} + |x|^{N(mM-p)} |y'|^{M-q}) \end{aligned} \quad (3.7)$$

The magnitude of the error term here follows from corresponding bounds on the derivatives of the error term in the partial Taylor expansion $S(x, y) = \sum_{a,b \leq M} s_{ab} x^a y^b + O(|x|^M + |y|^M)$. Thus as long as $N\zeta$ is an integer, by taking M large enough we have that $\partial_x^p \partial_{y'}^q s_j(x^N, y')$ exists and is continuous on V'_{jk} . Hence $s_j(x^N, y')$ is a smooth function of x and y as needed.

Furthermore, $\partial_{y'}^{o_{jk}} s_j(x, r_{jk}) \neq 0$, while $\partial_{y'}^{o_{jk}-1} s_j(x, r_{jk}) = 0$. Hence the implicit function theorem (applied to $s_j(x^N, y')$) says that there is a function $t_{jk}(x)$ for small x with $t_{jk}(x^N)$ smooth such that $t_{jk}(0) = r_{jk}$ and

$$\partial_{y'}^{o_{jk}-1} s_j(x, t_{jk}(x)) = 0 \quad (3.8)$$

More generally, we also have

$$c|y' - t_{jk}(x)| \leq |\partial_{y'}^{o_{jk}-1} s_j(x, y')| \leq C|y' - t_{jk}(x)| \quad (3.9)$$

Translating in terms of $S(x, y)$, on V_{jk} we have

$$\partial_y^{o_{jk}-1} S(x, x^{m_j} t_{jk}(x)) = 0 \quad (3.10a)$$

$$cx^{\alpha_j - m_j o_{jk}} |y - x^{m_j} t_{jk}(x)| \leq |\partial_y^{o_{jk} - 1} S(x, y)| \leq Cx^{\alpha_j - m_j o_{jk}} |y - x^{m_j} t_{jk}(x)| \quad (3.10b)$$

Since the terms of $S_{m_j}(x, y)$ are on the line $a + m_j b = \alpha_j$, the maximum possible order of a zero of $S_{m_j}(1, y)$ is $\frac{\alpha_j}{m_j}$. Hence $o_{jk} \leq \frac{\alpha_j}{m_j}$ and the exponent in (3.10b) is a nonnegative number which we denote by p . Thus if we make the coordinate change $\phi_{jk}(x, y) = (x, y + x^{m_j} t_{jk}(x))$ and let $R_{jk}(x, y) = S \circ \phi_{jk}(x, y)$, then we have

$$\partial_y^{o_{jk} - 1} R_{jk}(x, 0) = 0 \quad (3.11a)$$

$$cx^p |y| \leq |\partial_y^{o_{jk} - 1} R_{jk}(x, y)| \leq Cx^p |y| \quad (3.11b)$$

We now iterate the above algorithm to $R_{jk}(x, y)$ on $\phi_{jk}^{-1}(V_{jk})$. We first slice into two pieces along the x -axis. These two pieces are done the same way, so we focus our attention on the $y > 0$ piece which we denote by W . We divide W into pieces $T'_{j'}$, $U'_{j'l'}$, and $V'_{j'k'}$ exactly as done above. To simplify notation, write $R(x, y) = R_{jk}(x, y)$, with the understanding that any subscripts on R really refer to subscripts on a fixed R_{jk} .

We will show that any positive zero $r'_{j'k'}$ of any $R_{m'_j}(1, y)$ has order at most $o_{jk} - 1$. Thus after at most o_{jk} iterations there will be no more $V'_{j'k'}$ and we will be done. The fact that $R(x, y)$ is a smooth function of $x^{\frac{1}{N}}$ and y rather than x and y does not cause any problems in future stages; it just means after the next stage N may be replaced by a large multiple of N . Also, there are no problems caused by the fact that the upper boundary of $\phi_{jk}^{-1}(V_{jk})$ is some curve $y = \xi x^{m_j} +$ higher order terms instead of $y = x^n$ as before; by shrinking ξ if necessary we can ensure that this curve lies harmlessly inside one of the new $U'_{j'l'}$ whereupon the only effect is to shrink this $U'_{j'l'}$ somewhat.

So we turn our attention to showing that the order of any new positive zero $r'_{j'k'}$ of $R_{m'_j}(1, y)$ is at most $o_{jk} - 1$. For this, we will use (3.11b). Note that such a zero occurs for $V'_{j'k'}$ of the form $\{(x, y) : (r'_{j'k'} - \xi)x^{m'_j} < y < (r'_{j'k'} + \xi)x^{m'_j}\}$. The analogue of (3.4) for $R(x, y)$ implies that on $V'_{j'k'}$ we have an expression

$$\begin{aligned} \partial_y^{o_{jk} - 1} R(x, y) &= x^{\alpha'_j} \partial_y^{o_{jk} - 1} R_{m'_j}(1, y) + x^{\alpha'_j + \zeta'_j} \partial_y^{o_{jk} - 1} f(x, y) \\ &\quad + O(|x|^M + |y|^{M - o_{jk} + 1}) \end{aligned} \quad (3.12)$$

But by (3.11b), along any curve $y = cx^{m'_j}$, the function $\partial_y^{o_{jk} - 1} R(x, y)$ ($= \partial_y^{o_{jk} - 1} R_{jk}(x, y)$) will always have a zero of the same order $x^{p + m'_j}$ as $x \rightarrow 0$. Thus if the zero $r'_{j'k'}$ of $R'_{j'}(1, y)$ were of order o_{jk} or greater (3.12) gives a contradiction: on the curve $y = r'_{j'k'} x^{m'_j}$ the function $\partial_y^{o_{jk} - 1} R(x, y)$ vanishes to order greater than α'_j , while on nearby curves $y = cx^{m'_j}$, $c \neq r'_{j'k'}$, it vanishes to order α'_j . Thus we conclude that the order of the zero $r'_{j'k'}$ is at most $o_{jk} - 1$ and therefore that the induction ends after finitely many steps.

Lastly, part d) and the fact that each $g_i(x^{M'})$ is smooth for some positive integer M' depending on M follows by induction. Namely, if V_{jk} comes from an edge e with

vertices (a, b) and (a', b') , the coordinate change coming from that stage of the induction is a smooth function of $x^{\frac{1}{b'-b}}$. In addition, each o_{jk} is at most $b' - b$, and the corresponding difference in y coordinates will be at most o_{jk} in future iterations. Also, the number of different U_{jl} and V_{jk} coming from that edge is bounded by twice the number of possible zeroes of $S_e(1, y)$, or $2(b' - b)$. Since there are at most o_{jk} iterations of the algorithm, the result follows. This completes the proof of Theorem 3.1.

4. The beginning of the proofs of Theorems 1.1 and 1.2; the first decomposition and preliminary lemmas.

We will prove Theorems 1.1 and 1.2 simultaneously; the proofs have much in common. Furthermore, in proving Theorem 1.1 we may assume that $S(x, y)$ has a degenerate critical point at $(0, 0)$, as the nondegenerate (Morse) result can be read off from the explicit formulas (given for example in [G1]) for the leading term of the asymptotics for $M_{S,D}(\epsilon)$. Given this and that Theorem 1.2 excludes the nondegenerate case, the arguments of this section will always be under the assumption that $S(x, y)$ has a degenerate critical point at $(0, 0)$. Note also that we may assume that $S(x, y)$ is in superadapted coordinates; a fixed coordinate change does not affect the statements of Theorem 1.1 or 1.2.

Note next that for a given vertex v of $N(S)$ there is at most one value of t for which the Taylor expansion of $S + tf$ at the origin does not have a $c_v x^v$ term. Thus there are at most finitely many values of t for which the Taylor expansion of $S + tf$ at the origin does not contain a $c_v x^v$ term for each vertex v of $N(S)$. In other words, other than for these values one has $N(S) \subset N(S + tf)$. Thus excluding these values of t , in proving Theorems 1.1 and 1.2 we may assume that $N(S) \subset N(S + tf)$.

Next, notice that in proving Theorem 1.1 it actually suffices to show (1.4) holds for all sufficiently small $\epsilon > 0$ and not for all ϵ . The reason is as follows. Suppose for any sufficiently small disk D centered at the origin one has (1.3) for sufficiently small ϵ . Fix one such neighborhood D_0 . Then for all $\epsilon < \epsilon_0$ we have

$$M_{S,D_0}(\epsilon) \leq C_S \epsilon^j |\ln \epsilon|^p$$

Since $M_{S,D}(\epsilon)$ is monotone in D , for all $D \subset D_0$, for $\epsilon < \epsilon_0$ one also has

$$M_{S,D}(\epsilon) \leq C_S \epsilon^j |\ln \epsilon|^p$$

But by shrinking D enough the inequality will also hold for $\frac{1}{2} > \epsilon \geq \epsilon_0$. Thus if we fix one such shrunken D , call it D_1 , then (1.4) will now hold for all $\frac{1}{2} > \epsilon > 0$ for any $D \subset D_1$.

We now begin the proofs of Theorem 1.1 and 1.2. As indicated above, we can assume that $S(x, y)$ is in superadapted coordinates and that $N(S) \subset N(S + tf)$. Our goal will be to show that under the hypotheses of Theorem 1.1, for all but finitely many t (1.4) holds for sufficiently small ϵ , and that under the hypotheses of Theorem 1.2, for all but finitely many t one has the estimate $M_{S+tf,D}(\epsilon) \leq C \epsilon^j |\ln \epsilon|^p$. Here C may depend on S, t, f , and D .

We now fix a small disk D centered at the origin. We divide D into eight pieces through the x and y axes and two curves l_1 and l_2 chosen as follows. If the bisectrix intersects the interior of a compact edge e of $N(S)$ with equation $a + mb = \alpha$, then we choose l_1 and l_2 to be any two curves of the form $y = c|x|^m$ and $y = -c|x|^m$ so long as c is not a zero of $S_e(1, y)$ or $S_e(-1, y)$. (Recall that the polynomial $S_e(x, y)$ is the sum of the terms $s_{ab}x^a y^b$ of $S(x, y)$'s Taylor expansion at the origin with $(a, b) \in e$). If the bisectrix intersects $N(S)$ at a vertex (d, d) , we choose l_1 to be $y = |x|^m$ and l_2 to be $y = -|x|^m$, where m is such that a line with equation $a + mb = \alpha$ intersects $N(S)$ at the single point (d, d) . If the bisectrix intersects $N(S)$ in the interior of the horizontal or vertical rays, switching the roles of the x and y axes if necessary we can assume it's the horizontal ray and $N(S)$'s lowest vertex is of the form (c, d) for $c < d$. In this case we choose l_1 to be $y = |x|^m$ and l_2 to be $y = -|x|^m$ where a line with equation $a + mb = \alpha$ intersects $N(S)$ at the single point (c, d) .

In the above fashion D is divided into eight pieces E_1, \dots, E_8 . Clearly it suffices to show the desired bounds for each $M_{S+tf, E_i}(\epsilon)$. The argument for each E_i is the same, so we will focus our attention on the piece from the upper right quadrant between the x axis and the curve $y = cx^m$ or $y = x^m$. Denote this piece by E . We now apply the resolution of singularities algorithm of section 3 to $S(x, y)$, obtaining the corresponding sets D_i . Define $D'_i = D_i \cap E$. Clearly it suffices to show the desired upper bounds for each $M_{S+tf, D'_i}(\epsilon)$, which is what we will do.

Next, let ϕ_i be the maps of Theorem 3.1. Parts b) and c) of this theorem say that $S \circ \phi_i(x)$ is approximately a monomial in the precise sense given there. Denote $\phi_i^{-1}D'_i$ by F_i , $S \circ \phi_i(x)$ by $S_i(x)$, and $f \circ \phi_i(x)$ by $f_i(x)$. Since ϕ_i has determinant ± 1 , $M_{S+tf, D'_i}(\epsilon) = M_{S_i+tf_i, F_i}(\epsilon)$. Thus our task is to prove good upper bounds for $M_{S_i+tf_i, F_i}(\epsilon)$. Also, in Theorem 1.1 the smallness assumptions on the suprema of derivatives of f are implied by corresponding smallness assumptions on the derivatives of f_i (possibly with a different δ), so in our future arguments we may always assume the conditions on f_i instead of f without loss of generality.

Analogous to above, excluding at most finitely many t we can assume that $N(S_i) \subset N(R_i)$. Also note that by Theorem 3.1, the set F_i is a "curved triangle" in the sense that there are $h_i(x)$ and $H_i(x)$ such that for some $\eta_2 > \eta_1 > 0$ we have

$$\{(x, y) : 0 < x < \eta_1, h_i(x) < y < H_i(x)\} \subset F_i \subset \{(x, y) : 0 < x < \eta_2, h_i(x) < y < H_i(x)\}$$

Here $h_i(x^{M'})$ and $H_i(x^{M'})$ are smooth for the M given by the theorem. The function $h_i(x)$ may or may not be the zero function, and the Taylor expansion of $H_i(x)$ in fractional powers of x has some nonvanishing initial term $A_i x_i^N$. Theorem 3.1 also says that we can let (a_i, b_i) be such that $S_i(x, y)$ is comparable to a multiple of $x^{a_i} y^{b_i}$ on F_i in the sense of part b) or c) of the theorem. The following lemma gives us various conditions satisfied by (a_i, b_i) , N_i , and $N(S_i)$ what we will make use of.

Lemma 4.1.

a) If the bisectrix intersects $N(S)$ in the interior of a compact edge, then if $S_i(x)$ is viewed as a function on the $x > 0$ part of a neighborhood of the origin, $S_i(x)$ is in superadapted coordinates with the same Newton distance d that $N(S)$ has, and the bisectrix intersects $N(S_i)$ in the interior of a compact edge e_i . If the equation of this edge is denoted by $a + m_i b = \alpha_i$, then $N_i \geq m_i$. Furthermore $b_i < d$ and $a_i > d$. The ordinate of the intersection of the bisectrix with the line of slope $-\frac{1}{N_i}$ containing (a_i, b_i) , given by $\frac{a_i + N_i b_i}{1 + N_i}$, is at most the Newton distance d of $N(S)$.

b) If the bisectrix intersects $N(S)$ at the vertex (d, d) , the same is true for $N(S_i)$, and once again S_i is in superadapted coordinates with Newton distance d . Either $(a_i, b_i) = (d, d)$, which happens for at least one i , or $b_i < d$ and $a_i > d$ like above. In the latter case we again have $\frac{a_i + N_i b_i}{1 + N_i} \leq d$.

c) If the bisectrix intersects $N(S)$ in the interior of one of the rays, then the same is true for $N(S_i)$ and again S_i is in superadapted coordinates with Newton distance d . One of two things occurs. The first possibility is that $(a_i, b_i) = (c, d)$ for some $c < d$, the lower boundary of F_i is the x -axis, and part c) of Theorem 3.1 holds. The other possibility is that $b_i < d$ and $a_i \geq d$. In either case, we have the strict inequality $\frac{a_i + N_i b_i}{1 + N_i} < d$.

Proof. Recall that we divided a disk D centered at the origin into 8 pieces, each of which after a coordinate change of the form $(x, y) \rightarrow (\pm x, \pm y)$ or $(\pm y, \pm x)$ becomes of the form $E = \{(x, y) \in D : 0 < y < cx^m\}$, and we are focusing our attention on E . In the new coordinates, $S(x, y)$ becomes a function which we denote by $S_0(x, y)$. Note that $S_0(x, y)$ is automatically still in superadapted coordinates. In the setting of part a) of this lemma, the bisectrix intersects $N(S_0)$ at the point (d, d) which is in the interior of a compact edge e with equation $a + mb = \alpha$ for some α, m as above. In the setting of part b), the intersection is still (d, d) which is now a vertex of $N(S_0)$, and in the setting of part c), the intersection is (d, d) which is in the interior of the horizontal ray.

We now give some facts that are immediate consequences of the proof of the resolution of singularities algorithm of section 3, applied to $S_0(x, y)$ on E . First, each $\phi_i(x, y)$ is of the form $(x, \pm y - g_i(x))$, where $g_i(x^K)$ is a smooth function for some K . Next, if $g_i(x)$ is not the zero function then the Taylor expansion of $g_i(x)$ in fractional powers of x has initial term $p_i x^{l_i}$ where $l_i \geq m$ is such that $N(S_0)$ has an edge with equation $a + l_i b = \alpha_i$ for some i . The definition of E is such that (d, d) is either the upper vertex of this edge, or the edge lies entirely below (d, d) . The number N_i , defined such that the upper boundary of F_i has equation $y = q_i x^{N_i} +$ higher order terms, satisfies $N_i \geq m$ and the algorithm ensures that (a_i, b_i) is a vertex of $N(S_i)$. The definition of E is such that in the settings of part a) or b) of this lemma, either $(a_i, b_i) = (d, d)$ and $N_i = m$, or (a_i, b_i) is the lower vertex of a compact edge of $N(S_i)$ of slope $-\frac{1}{N_i}$. In the latter case either (d, d) is the upper vertex of the edge or it lies entirely below (d, d) .

We now proceed with the proof, starting with part a). If $l_i > m$, then the coordinate change $\phi_i(x, y)$ will not affect $(S_0)_e(x, y)$, the sum of the terms of $S_0(x, y)$'s

Taylor series at $(0, 0)$ corresponding to the edge $a + mb = \alpha$. Hence the resulting function $S_i(x, y)$ will be in superadapted coordinates like before. On the other hand, if $l_i = m$, then $(S_0)_e(x, y)$ becomes $(S_0)_e(x, \pm y - p_i x^m)$ after the coordinate change. Hence $(S_0)_e(1, y)$ becomes $(S_0)_e(1, \pm y - p_i)$ and $(S_0)_e(-1, y)$ becomes $(S_0)_e(-1, \pm y - p_i)$. Shifting the y variable does not change the fact that the definition of superadapted coordinates holds; the condition is that these polynomials have zeroes of order less than d . Hence in the $l_i = m$ case we are in superadapted coordinates as well. In either case, the bisectrix still intersects $N(S_i)$ at (d, d) , which is in the interior of a compact edge e_i with equation $a + mb = \alpha$ for some α . As a result (a_i, b_i) , being a vertex of $N(S_i)$ lying below (d, d) , satisfies $a_i > d$ and $b_i < d$ as needed. Using the last paragraph, (a_i, b_i) is the lower vertex of a compact edge e_i of $N(S_i)$ with slope $-\frac{1}{N_i}$ which either contains (d, d) or is below the edge containing (d, d) . Thus the intersection of the bisectrix with the line containing e_i is at (d, d) or below. But (a_i, b_i) is on this edge, so the intersection occurs at $(\frac{a_i + N_i b_i}{1 + N_i}, \frac{a_i + N_i b_i}{1 + N_i})$. Hence $\frac{a_i + N_i b_i}{1 + N_i} \leq d$ as needed. This completes the proof of part a).

Next we suppose we are in the setting of part b). In this case, the initial term of $g_i(x)$ is $p_i x^{l_i}$ for some $l_i > m$ since $N(S_0)$ has no edge with equation of the form $a + mb = \alpha$. As a result, the coordinate change $\phi_i(x, y)$ will not alter the fact that the bisectrix intersects the Newton polygon at (d, d) . Furthermore, one will still be in superadapted coordinates; if e is an edge of $N(S_0)$ containing (d, d) , then either $(S_0)_e(x, y)$ is unchanged by the coordinate change, or $(S_0)_e(1, y)$ becomes $(S_i)_e(1, y) = (S_0)_e(1, \pm y - p_i)$ and $(S_0)_e(-1, y)$ becomes $(S_i)_e(-1, y) = (S_0)_e(-1, \pm y - p_i)$ like above. In either event $S_i(x, y)$ will still be in superadapted coordinates. By the last paragraph, (a_i, b_i) is either (d, d) or a lower vertex. In the former case, $\frac{a_i + N_i b_i}{1 + N_i} = d$, and in the latter case, exactly as in part a) we have $\frac{a_i + N_i b_i}{1 + N_i} \leq d$ as needed. This completes the proof of part b).

Suppose we are in the setting of part c). Then the bisectrix intersects $N(S_0)$ either in the interior of the horizontal or vertical ray. Suppose it is the horizontal ray. Then $N(S_0)$ has a vertex (c, d) for some $c < d$. In this case no further subdivisions are necessary; we can take $S_i = S_0$ and let F_i be all of $\{(x, y) \in D : 0 < y < x^m\}$. Part c) of Theorem 3.1 automatically holds. So suppose the bisectrix intersects $N(S_0)$ in the interior of the vertical ray. In this case, the highest vertex of $N(S_0)$ is (d, c) for some $c < d$. Since any coordinate change ϕ_i fixes the highest vertex of $N(S_0)$, the highest vertex of $N(S_i)$ is (d, c) as well. Thus S_i is in superadapted coordinates with the bisectrix intersecting $N(S_i)$ inside its vertical ray. Since (a_i, b_i) is either the vertex (d, c) or a lower one, we have $a_i \geq d$ and $b_i \leq c < d$. Lastly, since the bisectrix intersects $N(S_i)$ inside the vertical ray, the ordinate of the intersection of any supporting line of $N(S_i)$ containing (a_i, b_i) with $N(S_i)$ is less than d . Hence $\frac{a_i + N_i b_i}{1 + N_i} < d$ and we are done.

5. The Main Argument.

We work in the setting of section 4. For some fixed value of t , let $R_i = S_i + t f_i$. We will proceed along the lines outlined in section 1, dividing a given F_i into finitely many pieces and show that, excluding finitely many values of t , the contribution to $M_{R_i, F_i}(\epsilon)$ of

each piece satisfies the upper bounds given by Theorem 1.1 or 1.2. We start this as follows. Since $S_i(x^{M'}, y)$ and $f_i(x^{M'}, y)$ are smooth functions, where M is as in Theorem 3.1, by Corollary 3.2, we can apply the resolution of singularities algorithm to $R_i(x, y)$. We now do so, but focus our attention only on the first stage of the algorithm, dividing the upper right quadrant into the sets called T_j and U_j in the proof. To highlight their dependence on i here, we refer to them as T_{ij} and U_{ij} here. Clearly, we need only consider those T_{ij} and U_{ij} that intersect F_i . Each U_{ij} corresponds to an edge of $N(R_i)$ whose equation we write as $a + m_{ij}b = \alpha_{ij}$, while each T_{ij} corresponds to a vertex of $N(R_i)$.

If $N(R_i)$ has a compact edge whose upper vertex is on or below the line $y = b_i$ and has slope greater than $-\frac{1}{N_i}$, let $e_{ij'}$ be the uppermost amongst all such edges. If $e_{ij'}$ exists, let G_i denote the union of $U_{ij'}$ with any T_{ij} and U_{ij} corresponding to edges and vertices of $N(R_i)$ below $e_{ij'}$. If $e_{ij'}$ does not exist, simply define G_i to be the lowest T_{ij} . We will now find upper bounds for $M_{R_i, G_i}(\epsilon)$ that are as good as needed for Theorem 1.1 or 1.2.

Lemma 5.1. Under the assumptions of Theorem 1.1, we have $M_{R_i, G_i}(\epsilon) \leq C_S \epsilon^j |\ln \epsilon|^p$, and under the assumptions of Theorem 1.2 we have $M_{R_i, G_i}(\epsilon) \leq A \epsilon^j |\ln \epsilon|^p$ for some constant A .

Proof. Let (a', b') be the uppermost vertex of the union of all edges and vertices of $N(R_i)$ that correspond to a T_{ij} or U_{ij} included in G_i . Note that by definition of G_i , we have $b' \leq b_i$. Write the Taylor expansion of $R_i(x, y)$ at the origin as $\sum_{cd} r_{cd} x^c y^d$, so that $r_{a'b'} x^{a'} y^{b'}$ denotes the term corresponding to the vertex (a', b') . Note that this Taylor expansion may contain fractional powers of x , but not y . By Lemma 2.4 c), on G_i we have

$$|\partial_y^{b'} R_i(x, y)| > \frac{b'!}{2} |r_{a'b'} x^{a'}| \quad (5.1)$$

Thus by Lemma 2.2 we have

$$\begin{aligned} M_{R_i, G_i}(\epsilon) &= |\{(x, y) \in G_i : |R_i(x, y)| < \epsilon\}| \\ &\leq 4 |\{(x, y) \in G_i : \frac{1}{2} |r_{a'b'} x^{a'} y^{b'}| < \epsilon\}| \end{aligned} \quad (5.2)$$

The right-hand side of (5.2) may be estimated using Lemma 2.3. We break into cases, depending on whether $b' > a'$, $b' = a'$, or $b' < a'$. If $b' > a'$, the lemma says that the right-hand side of (5.2) is bounded by $C r^\eta \epsilon^{\frac{1}{b'}}$ where r is the radius of D . Since $b' \leq b_i \leq d$, d the Newton distance of S , this can be made less than $\epsilon^{\frac{1}{d}}$ by shrinking the radius of original disk D enough. Since the growth index of $S(x, y)$ is given by $\frac{1}{d}$ in superadapted coordinates, this is at least as good as the estimate we need.

If $b' = a'$, then Lemma 2.3 tells us that

$$|\{(x, y) \in G_i : \frac{1}{2} |r_{b'b'} x^{b'} y^{b'}| < \epsilon\}| < C \epsilon^{\frac{1}{b'}} |\ln \epsilon| \quad (5.3)$$

Here the constant C depends on lower bounds for $|r_{b'b'}|$ as well as the set A_1 . Since $b' \leq b_i \leq d$, this is better than the estimate we seek unless $b' = b_i = d$. So we suppose $b' = b_i = d$. Since $N(S_i) \subset N(R_i)$, we must have $a_i \geq a' = d$. By Theorem 4.1, the only way one can have $a_i \geq d$ and $b_i = d$ is for $(a_i, b_i) = (d, d)$. Hence we have $(a_i, b_i) = (a', b') = (d, d)$. In view of Theorem 3.1c), F_i contains a set of the form $\{(x, y) : 0 < x < \eta, x^{m_1} < y < x^{m_2}\}$ on which $S_i(x, y) \sim x^d y^d$, and so by Lemma 2.3b) one has that $M_{S_i, G_i}(\epsilon) > c\epsilon^{\frac{1}{d}} |\ln \epsilon|$. Hence we must be in the situation where the growth index $\frac{1}{d}$ of $S(x, y)$ has multiplicity 1, and (5.2) and (5.3) give the desired estimate for Theorem 1.2. Next, note that

$$|\{(x, y) \in G_i : \frac{1}{2}|r_{dd}x^d y^d| < \epsilon\}| = |\{(x, y) \in G_i : x^d y^d < \frac{2\epsilon}{r_{dd}}\}| \quad (5.4)$$

In the setting of Theorem 1.1, by making δ sufficiently small we can ensure that $r_{dd} < 2s_{dd}$, where $s_{dd}x^d y^d$ denotes the term of the Taylor expansion of $S_i(x, y)$. Hence, using Lemma 2.3b) on (5.2) and (5.4), we see that $M_{R_i, G_i}(\epsilon) < C_S \epsilon^{\frac{1}{d}} |\ln \epsilon|$, the desired estimate for Theorem 1.1.

We now turn to the case where $b' < a'$. The definition of G_i is such that G_i is contained in a set of the form $\{(x, y) : 0 < x < \eta, 0 < y < x^m\}$ for some $m > N_i$. We apply part c) of Lemma 2.3, which says that

$$|\{(x, y) \in G_i : \frac{1}{2}|r_{a'b'}x^{a'} y^{b'}| < \epsilon\}| \leq C\epsilon^{\frac{m+1}{a'+mb'}} \quad (5.5a)$$

In view of (5.2), we have

$$M_{R_i, G_i}(\epsilon) = |\{(x, y) \in G_i : |R_i(x, y)| < \epsilon\}| < C'\epsilon^{\frac{m+1}{a'+mb'}} \quad (5.5b)$$

Like above, the constant C' depends on lower bounds for $R_{a'b'}$ (as well as (a', b')). Note that $\frac{a'+mb'}{m+1}$ is the ordinate of the intersection of the line of slope $-\frac{1}{m}$ containing (a', b') with the bisectrix. Since $a' > b'$ and $m > N_i$, this is strictly less than the corresponding ordinate for the line of slope $-\frac{1}{N_i}$ containing (a', b') . Since $N(R_i)$ contains $N(S_i)$, this will be at most the ordinate of the intersection of the bisectrix with line of the same slope $-\frac{1}{N_i}$ containing (a_i, b_i) , given by $\frac{a_i+b_i N_i}{1+N_i}$. This is at most d by Lemma 4.1. Hence by (5.5b), we have $M_{R_i, G_i}(\epsilon) < C\epsilon^{\frac{1}{d}+\eta}$ for some positive η , a better estimate than we need and we are done.

The next step is to prove upper bounds of Theorem 1.1 and 1.2 for the remaining $M_{R_i, T_{ij}}(\epsilon)$:

The upper bounds for $M_{R_i, T_{ij}}(\epsilon)$ when T_{ij} is not a subset of G_i .

Each such T_{ij} corresponds to some vertex of $N(R_i)$, which we denote by (p, q) . T_{ij} is of the form $\{(x, y) \in F_i : \frac{1}{\xi}x^{\frac{1}{m_2}} < y < \xi x^{m_1}\}$. Although we only need to bound $M_{R_i, T_{ij}}(\epsilon)$ for T_{ij} intersecting F_i , the following argument works for all T_{ij} . We will analyze

$R_i(x, y)$ on T_{ij} similarly to the way $R(x, y)$ was analyzed in (2.8). As above, we write the Taylor series of $R_i(x, y)$ at the origin as $\sum_{ab} r_{ab} x^a y^b$. We similarly write the Taylor series of $S_i(x, y)$ at the origin as $\sum_{ab} s_{ab} x^a y^b$. Note that

$$\begin{aligned} S_i(x, y) - s_{pq} x^p y^q = & \sum_{p \leq a < M, q \leq b < M, (a,b) \neq (p,q)} s_{ab} x^a y^b \\ + & \sum_{a < p, q < b < M, a+m_1 b \geq \alpha_1} s_{ab} x^a y^b + \sum_{p < a < M, b < q, am_2 + b \geq \alpha_2} s_{ab} x^a y^b + E_M(x, y) \end{aligned} \quad (5.6)$$

Here $a + m_1 b = \alpha_1$ and $m_2 a + b = \alpha_2$ are equations of the edges e_1 and e_2 of $N(R_i)$ meeting at (p, q) . In the case that e_1 doesn't exist, the second sum above disappears, and if e_2 doesn't exist then the third sum disappears.

First suppose (p, q) is not a vertex of $N(S)$, so that $s_{pq} = 0$. By definition, r_{pq} is nonzero. As in (2.8), for any $\delta > 0$ we can make each of the terms of (5.6) bounded by $\delta |r_{pq}| x^p y^q$ by choosing ξ small enough. In particular, we can make the absolute value of the entire right hand side of (5.6) less than $\frac{1}{4} |r_{pq}| x^p y^q$ on T_{ij} . Since $s_{pq} = 0$ this means $|S_i(x, y)| < \frac{1}{4} |r_{pq}| x^p y^q$ on T_{ij} . Similarly, by choosing ξ small enough, we can assume that the right-hand side of the analogue to (5.6) with S_i replaced by R_i is also less than $\frac{1}{4} |r_{pq}| x^p y^q$ on T_{ij} . Hence on T_{ij} we have

$$|R_i(x, y)| > \frac{3}{4} |r_{pq}| x^p y^q > 3 |S_i(x, y)| \quad (5.7)$$

As a result, we have

$$\begin{aligned} |\{(x, y) \in T_{ij} : |R_i(x, y)| < \epsilon\}| & \leq |\{(x, y) \in T_{ij} : |S_i(x, y)| < 3\epsilon\}| \\ & \leq C_S \epsilon^j |\ln \epsilon|^p \end{aligned}$$

This is at least as strong as the estimate we need.

Now suppose that $s_{pq} \neq 0$. Since (p, q) is a vertex of $N(R_i)$ and $N(S) \subset N(R_i)$, this means (p, q) is a vertex of $N(S)$ as well. Like above, by shrinking ξ enough one can assume that the right-hand side of (5.6) and its analogue with S_i replaced by R_i are less than $\frac{1}{2} |s_{pq}|$ and $\frac{1}{2} |r_{pq}|$ respectively. As a result, on T_{ij} we have

$$|S_i(x, y)| < \frac{3}{2} |s_{pq}| x^p y^q, \quad |R_i(x, y)| > \frac{1}{2} |r_{pq}| x^p y^q$$

Thus we have

$$|\{(x, y) \in T_{ij} : |R_i(x, y)| < \epsilon\}| < |\{(x, y) \in T_{ij} : |S_i(x, y)| < 3 \frac{|s_{pq}|}{|r_{pq}|} \epsilon\}| \quad (5.8)$$

The right-hand side of (5.8) is at most $C\epsilon^j |\ln \epsilon|^p$, the desired estimate in the setting of Theorem 1.2. As for Theorem 1.1, we may assume the δ of that Theorem is small enough so that $\frac{|s_{pq}|}{|r_{pq}|} < 2$. In this case, we have

$$\begin{aligned} |\{(x, y) \in T_{ij} : |S_i(x, y)| < 3 \frac{|r_{pq}|}{|s_{pq}|} \epsilon\}| &< |\{(x, y) \in T_{ij} : |S_i(x, y)| < 6\epsilon\}| \\ &< C_S \epsilon^j |\ln \epsilon|^p \end{aligned} \quad (5.9)$$

Combining (5.8) and (5.9) gives the desired estimates for Theorem 1.1 and we are done.

It remains to bound $M_{R_i, U_{ij}}(\epsilon)$ for the U_{ij} intersecting F_i that are not contained in G_i . Each U_{ij} corresponds to an edge e_{ij} of $N(R_i)$. There are only finitely many possible such e_{ij} for *any* R_i ; that is, there are only finitely many pairs of vertices that can be the endpoints of an edge corresponding to some such U_{ij} for any possible $S_i + tf_i$, regardless of what f_i is. To see why this is true, write the equation of e_{ij} as $a + m_{ij}b = \alpha_{ij}$, and its upper vertex as (a, b) . Since U_{ij} is a subset of F_i , we automatically have $m_{ij} \leq N_i$. We separately consider the cases $b < b_i$, $b = b_i$, and $b > b_i$, and show in each case that there are finitely many possibilities for e_{ij} .

We start with the case where $b < b_i$. In this case G_i takes care of all U_{ij} with $m_{ij} > N_i$, so we are left with the case when $m_{ij} = N_i$. There are only finitely many possibilities with $a \leq a_i$, so we may assume that $a > a_i$. Note that since (a_i, b_i) is in $N(S_i)$, it is also in $N(R_i)$. Since $a > a_i$ this means (a, b) can not be on a the vertical ray of $N(R_i)$. Instead it is the lower vertex of a compact edge e' of $N(R_i)$ of slope at least $\frac{b-b_i}{a-a_i} \geq -\frac{1}{a-a_i}$. Since the slope of e_{ij} is $-\frac{1}{N_i}$ and e_{ij} lies below e' , we must have that $-\frac{1}{N_i} < -\frac{1}{a-a_i}$. In other words we have $a < a_i + N_i$. Since it also true that $b < b_i$, there are finitely many possibilities for this to occur.

Next, consider the case where $b = b_i$. Here since $N(R_i)$ contains $N(S_i)$ and (a_i, b_i) is a vertex of $N(S_i)$, $a \leq a_i$ and there are finitely many possibilities for a . Since once again G_i takes care of all U_{ij} with $m_{ij} > N_i$, we have $m_{ij} = N_i$ and once again there are a finite number of possibilities for e_{ij} .

Lastly, we consider the situation where $b > b_i$. Then since (a, b) lies above the vertex (a_i, b_i) of $N(S_i)$, we have $a < a_i$. Since $N(R_i)$ contains $N(S_i)$, $(a_i, b_i) \in N(R_i)$. Hence the slope of e_{ij} is at most $\frac{b_i-b}{a_i-a} \leq b_i - b$. It is also greater than or equal to $-\frac{1}{N_i}$ because $U_{ij} \subset F_i$. Thus we have $-\frac{1}{N_i} \leq b_i - b$ or $b \leq b_i + \frac{1}{N_i}$. Coupled with the condition that $a < a_i$, once again we have finitely many possibilities for (a, b) and we are done.

Thus for our future arguments, it suffices to fix a single e_{ij} and prove the desired upper bounds for the U_{ij} associated with e_{ij} . Recall that for a fixed x the vertical cross-section of U_{ij} has width $(\frac{1}{\xi} - \xi)x^{m_{ij}}$. If one now applies the next step of the resolution of singularities algorithm of Theorem 3.1 (to $R_i(x, y)$), one divides U_{ij} into pieces V_{ijk} and W_{ijk} , where each V_{ijk} is of the form $\{(x, y) : (r_{ijk} - \xi)x^{m_{ij}} < y < (r_{ijk} + \xi)x^{m_{ij}} \text{ for a root}$

r_{ijk} of the polynomial $(S_i)_{e_{ij}}(1, y)$. On each W_{ijk} one has $R_i(x, y) \sim x^{\alpha_{ij}}$ where e_{ij} has equation $a + m_{ij}b = \alpha_{ij}$. We also need to split each V_{ijk} into two pieces V_{ijk}^1 , and V_{ijk}^2 , where V_{ijk}^2 is the portion where $|y - r_{ijk}x^{m_{ij}}| < x^{m_{ij}+\eta}$, and V_{ijk}^1 is the rest. η here is a small positive constant which must be sufficiently small for our arguments to work. Note that since there are at most boundedly many r_{ijk} for a given e_{ij} and boundedly many e_{ij} for our fixed $S(x, y)$, the total number of V_{ijk}^1, V_{ijk}^2 , and U_{ijk} is uniformly bounded given $S(x, y)$.

Our analysis will proceed as follows. We will first show that each $M_{R_i, W_{ijk}}(\epsilon)$ satisfies the bounds required of Theorems 1.1 or 1.2. Then we will show that if $\eta > 0$ is sufficiently small, a small additional argument will show that each $M_{R_i, V_{ijk}^1}(\epsilon)$ also satisfies the bounds. Afterwards, a separate argument will be used to show that for each V_{ijk}^2 intersecting F_i , $M_{R_i, V_{ijk}^2}(\epsilon)$ satisfies bounds better than the ones we need. We will do this by applying the full resolution of singularities algorithm on $f_i(x, y)$ on each V_{ijk}^2 . We will obtain the corresponding sets called D_i in Theorem 3.1, and show that for each D amongst them $M_{R_i, D}(\epsilon)$ satisfies better estimates than what we need.

The analysis of $M_{R_i, W_{ijk}}(\epsilon)$.

Since $N(R_i)$ contains $N(S_i)$ and e_{ij} has equation $a + m_{ij}b = \alpha_{ij}$, (a_i, b_i) is on or above the line containing e_{ij} and hence $a_i + m_{ij}b_i \geq \alpha_{ij}$. Since on W_{ijk} we have $R_i(x, y) \sim x^{\alpha_{ij}}$, we have $R_i(x, y) \geq C_1 x^{a_i + m_{ij}b_i}$. Since for fixed x the set W_{ijk} has cross-section $\sim x^{m_{ij}}$, we conclude that

$$|\{(x, y) \in W_{ijk} : |R_i(x, y)| < \epsilon\}| < C_2 \epsilon^{\frac{1+m_{ij}}{a_i+m_{ij}b_i}} \quad (5.10a)$$

If (a_i, b_i) is strictly above the line, then for some $\zeta > 0$ we have the even better estimate

$$|\{(x, y) \in W_{ijk} : |R_i(x, y)| < \epsilon\}| < C_3 \epsilon^{\frac{1+m_{ij}}{a_i+m_{ij}b_i} + \zeta} \quad (5.10b)$$

For now at least, we have no information concerning the constants C_2, C_3 of (5.10a) – (5.10b). Note that the exponent $\frac{1+m_{ij}}{a_i+m_{ij}b_i}$ is the reciprocal of the ordinate of the intersection of the bisectrix with the line of slope $-\frac{1}{m_{ij}}$ containing (a_i, b_i) . Also note that since $S_i(x, y) \sim x^{a_i+m_{ij}b_i}$ on W_{ijk} , we have

$$|\{(x, y) \in W_{ijk} : |S_i(x, y)| < \epsilon\}| > C_4 \epsilon^{\frac{1+m_{ij}}{a_i+m_{ij}b_i}} \quad (5.11)$$

Hence one has

$$\begin{aligned} |\{(x, y) \in W_{ijk} : |R_i(x, y)| < \epsilon\}| &< C_5 |\{(x, y) \in W_{ijk} : |S_i(x, y)| < \epsilon\}| \\ &< C_5 M_{S, D}(\epsilon) \\ &\leq C_6 \epsilon^{\frac{1}{d}} |\ln \epsilon|^p \end{aligned} \quad (5.12)$$

This gives the desired estimates for Theorem 1.2, which are also the desired estimates for Theorem 1.1 other than the constants, which will take some more work and which we now focus our attention on.

Note that if (a_i, b_i) lies strictly above the line containing e_{ij} , the added ζ in (5.10b) gives us any constant C_6 we want for ϵ small enough, thereby implying the estimate needed for Theorem 1.1. Thus we need only consider the case where (a_i, b_i) is actually on the line containing e_{ij} . Furthermore, in any situation in which $\frac{1+m_{ij}}{a_i+m_{ij}b_i}$ is strictly greater than $\frac{1}{d}$, we could once again make C_6 arbitrarily small. So it makes sense to analyze when we have do not have this strict inequality; we will see momentarily that this only happens in special situations.

Consider the case when the bisectrix intersects the interior of a compact edge of $N(S_i)$. In this case, by Lemma 4.1a) we have $b_i < a_i$. Hence if m_{ij} is greater than the minimal possible value on F_i , given by N_i , then we have $\frac{1+m_{ij}}{a_i+m_{ij}b_i} > \frac{1+N_i}{a_i+N_i b_i}$, which in turn is at least $\frac{1}{d}$. Hence when the bisectrix intersects the interior of a compact edge of $N(S_i)$, equality can only occur if $m_{ij} = N_i$.

Next, consider the case when the bisectrix intersects the vertex (d, d) of $N(S_i)$. In this case, by Lemma 4.1b) either $(a_i, b_i) = (d, d)$ or $a_i > d, b_i < d$, and $\frac{1+N_i}{a_i+N_i b_i} \geq \frac{1}{d}$. In the latter case, if $m_{ij} > N_i$ then like above we have $\frac{1+m_{ij}}{a_i+m_{ij}b_i} > \frac{1+N_i}{a_i+N_i b_i} \geq \frac{1}{d}$, and equality can occur only when $m_{ij} = N_i$.

Lastly, consider the case where the bisectrix intersects $N(S_i)$ in the interior of one of the rays. In this case Lemma 4.1c) applies and either $(a_i, b_i) = (c, d)$ for some $c < d$, or (a_i, b_i) satisfies $a_i \geq d, b_i < d$ and $\frac{1+N_i}{a_i+N_i b_i} > \frac{1}{d}$. In the latter case since $m_{ij} \geq N_i$ we automatically have $\frac{1+m_{ij}}{a_i+m_{ij}b_i} > \frac{1}{d}$ and equality does not occur. In the former case, since $c < d$ we also have $\frac{1+m_{ij}}{c+m_{ij}d} > \frac{1}{d}$ and equality also does not occur.

In summary, in bounding $M_{R_i, W_{ijk}}(\epsilon)$ we have already covered all possible cases except the following two situations. First, we can have $m_{ij} = N_i, a_i > d$, and $b_i < d$. Secondly, the bisectrix may intersect $N(S_i)$ at the vertex (d, d) with $(a_i, b_i) = (d, d)$. Furthermore, as mentioned above we only need to consider the situation when (a_i, b_i) is on the line containing the edge e_{ij} . The argument we will use for these two situations actually will give the needed bounds for all of $M_{R_i, V_{ijk}}(\epsilon)$ in those situations as well. In fact, recalling that $U_{ij} = \cup_k V_{ijk} \cup \cup_k W_{ijk}$, what we will do is prove upper bounds for all of $M_{R_i, U_{ij}}(\epsilon)$ in both situations.

Bounding $M_{R_i, U_{ij}}(\epsilon)$ in the two exceptional situations.

We start with the first situation, where $m_{ij} = N_i, a_i > d, b_i < d$, and (a_i, b_i) is on the line containing e_{ij} . We write the Taylor expansions of $R_i(x, y)$ and $S_i(x, y)$ at the origin as $\sum_{a,b} r_{ab} x^a y^b$ and $\sum_{a,b} s_{ab} x^a y^b$. Here the b 's are all integers but a may be a

nonintegral positive rational number. We write

$$R_i(x, y) = \sum_{a+N_i b=\alpha_{ij}} r_{ab} x^a y^b + \sum_{a+N_i b>\alpha_{ij}, a, b<M} r_{ab} x^a y^b + O(|x|^M + |y|^M) \quad (5.13a)$$

$$S_i(x, y) = \sum_{a+N_i b=\alpha_{ij}} s_{ab} x^a y^b + \sum_{a+N_i b>\alpha_{ij}, a, b<M} s_{ab} x^a y^b + O(|x|^M + |y|^M) \quad (5.13b)$$

Here M is some large integer and α_{ij} is such that e_{ij} has equation $a + N_i b = \alpha_{ij}$. Since for fixed x , the y cross-section of U_{ij} is contained in some $[c_{ij}x^{N_i}, A_i x^{N_i}]$ where $c_{ij}, A_i > 0$ and A_i depends only on S_i , it makes sense to look at $R_i(x, x^{N_i}y)$ and $S_i(x, x^{N_i}y)$, given by

$$R_i(x, x^{N_i}y) = x^{\alpha_{ij}} (R_i)_{e_{ij}}(1, y) + x^{\alpha_{ij}+\zeta} P(x, y) + O(|x|^{M'}) \quad (5.14a)$$

$$S_i(x, x^{N_i}y) = x^{\alpha_{ij}} (S_i)_{e_{ij}}(1, y) + x^{\alpha_{ij}+\zeta} Q(x, y) + O(|x|^{M'}) \quad (5.14b)$$

Here $\zeta > 0$, P and Q are polynomials in y and a fractional power of x , and M' is a large integer that grows linearly with M . Similarly, we may look at the b_i th y derivatives of R_i and S_i and get the following expressions

$$(\partial_y^{b_i} R_i)(x, x^{N_i}y) = x^{\alpha_{ij}-b_i N_i} \partial_y^{b_i} ((R_i)_{e_{ij}}(1, y)) + x^{\alpha_{ij}-b_i N_i+\zeta} \tilde{P}(x, y) + O(|x|^{M'}) \quad (5.15a)$$

$$(\partial_y^{b_i} S_i)(x, x^{N_i}y) = x^{\alpha_{ij}-b_i N_i} \partial_y^{b_i} ((S_i)_{e_{ij}}(1, y)) + x^{\alpha_{ij}-b_i N_i+\zeta} \tilde{Q}(x, y) + O(|x|^{M'}) \quad (5.15b)$$

Part b) or c) of Theorem 3.1 says that $\partial_y^{b_i} S_i(x, y) > Cx^{\alpha_i} = Cx^{\alpha_{ij}-b_i N_i}$ on the F_i which contains the set U_{ij} under consideration. Hence letting $x \rightarrow 0$ in (5.15b), we may conclude that $\partial_y^{b_i} ((S_i)_{e_{ij}}(1, y))$ has no zeroes on $[0, A_i]$. Hence there is a C_S such that for the (x, y) being considered we have

$$|\partial_y^{b_i} ((S_i)_{e_{ij}}(1, y)) x^{\alpha_{ij}-b_i N_i}| > C_S x^{\alpha_{ij}-b_i N_i} \quad (5.16)$$

Suppose now we are working under the hypotheses of Theorem 1.1. Then by shrinking δ sufficiently if necessary, we can assume each coefficient of each term of $(\partial_y^{b_i} f_i)_{e_{ij}}(1, y)$ is as small as we want. In particular, we can assume that on $[c_{ij}, A_i]$ we have

$$|\partial_y^{b_i} ((f_i)_{e_{ij}}(1, y)) x^{\alpha_{ij}-b_i N_i}| < \frac{1}{2} C_S x^{\alpha_{ij}-b_i N_i} \quad (5.17a)$$

Given that $R_i = S_i + t f_i$, for $|t| < 1$ this means that

$$|\partial_y^{b_i} ((R_i)_{e_{ij}}(1, y)) x^{\alpha_{ij}-b_i N_i}| > \frac{1}{2} C_S x^{\alpha_{ij}-b_i N_i} \quad (5.17b)$$

In view of (5.15a), this means that if x is sufficiently small, which we may assume, then we have

$$|(\partial_y^{b_i} R_i)(x, x^{N_i}y)| > \frac{1}{4} C_S x^{\alpha_{ij}-b_i N_i} \quad (5.18a)$$

Translating this back into the original coordinates, this means that on U_{ij} we have

$$|\partial_y^{b_i} R_i(x, y)| > \frac{1}{4} C_S x^{\alpha_{ij} - b_i N_i} \quad (5.18b)$$

Hence by Lemma 2.2, for a given x we have

$$|\{y : |R_i(x, y)| < \epsilon\}| < 4 |\{y : \frac{1}{4b_i!} C_S x^{\alpha_{ij} - b_i N_i} y^{b_i} < \epsilon\}| \quad (5.19)$$

Integrating in x , we obtain

$$|\{(x, y) \in U_{ij} : |R_i(x, y)| < \epsilon\}| < 4 |\{(x, y) \in U_{ij} : \frac{1}{4b_i!} C_S x^{\alpha_{ij} - b_i N_i} y^{b_i} < \epsilon\}| \quad (5.20)$$

We bound the right-hand side of (5.20) using Lemma 2.3. Since $\alpha_{ij} = a_i + N_i b_i$ and $b_i < a_i$, we have $b_i < \alpha_{ij} - b_i N_i$ and part c) of Lemma 2.3 applies. We obtain that

$$|\{(x, y) \in U_{ij} : |R_i(x, y)| < \epsilon\}| < C'_S \epsilon^{\frac{N_i + 1}{\alpha_{ij}}} \quad (5.21)$$

Since $\alpha_{ij} = a_i + N_i b_i$, by Lemma 4.1a) the exponent $\frac{N_i + 1}{\alpha_{ij}}$ is at most $\frac{1}{d}$, and (5.21) gives us the estimate we need. Thus we are done in the setting of Theorem 1.1.

Suppose now that we are in the setting of Theorem 1.2. Since $(a_i, b_i) \in e_{ij}$ and S_i is in superadapted coordinates, any zero of $(S_i)_{e_{ij}}(1, y)$ is of order less than d . As a result, no matter what f_i is, there are at most finitely many t for which $(R_i)_{e_{ij}}(1, y) = (S_i)_{e_{ij}}(1, y) + t(f_i)_{e_{ij}}(1, y)$ has a zero of order d or greater on $[0, A_i]$. (This can be proven by an elementary argument.) Hence excluding those t , for a given t we can divide $[0, N_i]$ into closed intervals B_1, \dots, B_m such that on each B_k , $\partial_y^l (R_i)_{e_{ij}}(1, y)$ is nonvanishing for some $0 \leq l < d$. We then apply the above argument for each B_k , replacing b_i by l . Thus for each k the corresponding set of points where $|R_i(x, y)| < \epsilon$ has measure less than $C|\epsilon|^{\frac{1}{d}}$. Adding over all k we get the upper bounds required by Theorem 1.2. Thus we have proven the desired bounds for $M_{R_i, U_{ij}}(\epsilon)$ in the case that the bisectrix intersects $N(S_i)$ in the interior of a compact edge.

We now turn to the case where the bisectrix intersects $N(S_i)$ at a vertex (d, d) with $(a_i, b_i) = (d, d)$. For fixed x , the y -cross-section of U_{ij} is contained in $[c_{ij}x^{m_{ij}}, C_{ij}x^{m_{ij}}]$ for some c_{ij} and C_{ij} which depend on the function $R_i(x, y)$. We write $[c_{ij}x^{m_{ij}}, C_{ij}x^{m_{ij}}]$ as the union of $[c_{ij}x^{m_{ij}}, x^{m_{ij}}]$ and $[x^{m_{ij}}, C_{ij}x^{m_{ij}}]$, and correspondingly write $U_{ij} = U_{ij}^1 \cup U_{ij}^2$. We focus our attention on U_{ij}^1 only, as U_{ij}^2 is done analogously with the roles of the two axes reversed. One technical point here is worth mentioning. Since $(a_i, b_i) = (d, d)$ and $S_0(x, y)$ was in superadapted coordinates, the algorithm of Theorem 3.1 is such that $\phi_i(x, y)$ is the identity; F_i is carved out of the original disk and there is no coordinate change. This is relevant here because it implies fractional powers of x do not appear; one can switch the x and y axes without any issues caused by fractional powers arising.

The argument is basically the same as that of (5.13) – (5.21) so we will be brief. This time we write

$$R_i(x, y) = \sum_{a+m_{ij}b=\alpha_{ij}} r_{ab}x^a y^b + \sum_{a+m_{ij}b>\alpha_{ij}, a,b<M} r_{ab}x^a y^b + O(|x|^M + |y|^M) \quad (5.22a)$$

$$S_i(x, y) = \sum_{a+m_{ij}b=\alpha_{ij}} s_{ab}x^a y^b + \sum_{a+m_{ij}b>\alpha_{ij}, a,b<M} s_{ab}x^a y^b + O(|x|^M + |y|^M) \quad (5.22b)$$

The analogue to (5.15a) – (5.15b) is

$$(\partial_y^d R_i)(x, x^{m_{ij}}y) = x^{\alpha_{ij}-dm_{ij}} \partial_y^d ((R_i)_{e_{ij}}(1, y)) + x^{\alpha_{ij}-dm_{ij}+\zeta} \tilde{P}(x, y) + O(|x|^{M'}) \quad (5.23a)$$

$$(\partial_y^d S_i)(x, x^{m_{ij}}y) = x^{\alpha_{ij}-dm_{ij}} \partial_y^d ((S_i)_{e_{ij}}(1, y)) + x^{\alpha_{ij}-dm_{ij}+\zeta} \tilde{Q}(x, y) + O(|x|^{M'}) \quad (5.23b)$$

Note that since (d, d) is on e_{ij} with equation $a + m_{ij}b = \alpha_{ij}$, the exponent $\alpha_{ij} - dm_{ij}$ in (5.23a) – (5.23b) is equal to d . Hence in the setting of Theorem 1.1, if the δ of that theorem is sufficiently small, the analogue to (5.18b) here becomes

$$(\partial_y^d R_i)(x, y) > \frac{1}{4} C_S'' x^d \quad (5.24)$$

Using Lemma 2.2, the analogue to (5.20) here is

$$|\{(x, y) \in U_{ij} : |R_i(x, y)| < \epsilon\}| < 4 |\{(x, y) \in U_{ij} : \frac{1}{4d!} C_S'' x^d y^d < \epsilon\}| \quad (5.25)$$

Hence using Lemma 2.3b) now, we get that

$$|\{(x, y) \in U_{ij} : |R_i(x, y)| < \epsilon\}| < C_S''' \epsilon^{\frac{1}{d}} |\ln \epsilon| \quad (5.26)$$

This is the desired estimate in the setting of Theorem 1.1. As for Theorem 1.2, we make modifications analogous to the ones before. Specifically, we again can exclude finitely many values of t for which $(R_i)_{e_{ij}}(1, y)$ has zeroes of order greater than d on $[c_{ij}, 1]$ and assume $(R_i)_{e_{ij}}(1, y)$ has no zeroes of order greater than d . One then proceeds as in the previous case, dividing $[c_{ij}, 1]$ into intervals on which either $(R_i)_{e_{ij}}(1, y)$ is nonvanishing or $\partial_y^k((R_i)_{e_{ij}}(1, y))$ is nonvanishing for some $k \leq d$. This completes the arguments for the $M_{R_i, U_{ij}}(\epsilon)$ in the exceptional cases.

The analysis of $M_{R_i, V_{ijk}^1}(\epsilon)$.

Analogous to (5.14a), for every i and j we have that

$$R_i(x, x^{m_{ij}}y) = x^{\alpha_{ij}} (R_i)_{e_{ij}}(1, y) + x^{\alpha_{ij}+\zeta} P(x, y) + O(|x|^{M'})$$

There is a zero r of $(R_i)_{e_{ij}}(1, y)$ such that on V_{ijk}^1 we have $|y - rx^{m_{ij}}| > x^{m_{ij}+\eta}$. Hence in the above equation, $|y - r| > x^\eta$. For any $\eta' > 0$, we can choose η so that $|y - r| > x^{\eta'}$

implies that $(R_i)_{e_{ij}}(1, y) > Cx^{\eta'}$ for the (x, y) being considered. Hence for any $\eta' > 0$ we can assume that

$$x^{\alpha_{ij}}(R_i)_{e_{ij}}(1, y) > Cx^{\alpha_{ij}+\eta'}$$

As long as $\eta' < \zeta$ and M' is sufficiently large, then we therefore have

$$|R_i(x, x^{m_{ij}}y)| > C'x^{\alpha_{ij}+\eta'} \quad (5.27)$$

Since V_{ijk}^1 is between the curves $y = c_1x^{m_{ij}}$ and $c_2x^{m_{ij}}$ for some c_1 and c_2 , we conclude that

$$|\{(x, y) \in V_{ijk}^1 : |R_i(x, y)| < \epsilon\}| < C''\epsilon^{\frac{1+m_{ij}}{\alpha_{ij}+\eta'}} \quad (5.28)$$

Thus for any (i, j) for which the exponent $\frac{1+m_{ij}}{\alpha_{ij}}$ is larger than what we need, by shrinking η' enough, (5.28) gives that $M_{R_i, V_{ijk}^1}(\epsilon)$ satisfies a better estimate than what we need. However, we saw that the only situations in which this exponent is not better than what we need are the exceptional cases discussed above. But these are exactly the cases in which we proved the desired upper bounds for all of $M_{R_i, U_{ij}}(\epsilon)$. Hence by shrinking η enough, (5.28) gives better than the needed estimates for all remaining V_{ijk}^1 and we are done.

The analysis of $M_{R_i, V_{ijk}^2}(\epsilon)$.

We now focus our attention on some fixed V_{ijk}^2 , which consists of the points in U_{ij} for which $|y - rx^{m_{ij}}| < x^{m_{ij}+\eta}$ for some $r \in \mathbf{R}$. The exact value of r will not be important in what follows. We apply the resolution of singularities algorithm of Theorem 3.1 to $f_i(x, y)$ on F_i , and consider the sets called D_i in that theorem that intersect V_{ijk} . Since we are already using the index i , we refer to them as D_l here. To each D_l there is a coordinate change ϕ_l such that $f_i \circ \phi_l$ is comparable to a monomial on D_l in the sense of Theorem 3.1 b) or c). Since we consider only those D_l intersecting V_{ijk} , the function ϕ_l is such that $|y| < x^{m_{ij}+\eta}$ on $\phi_l^{-1}D_l$. Write $S'_i(x, y) = S_i \circ \phi_l(x, y)$. Our first task will be to understand $S'_i(x, y)$'s behavior on the set $\phi_l^{-1}D_l$, which we denote by E_l .

To this end, note that the domain F_i of $S_i(x, y)$ has upper boundary given by $y = A_ix^{N_i} +$ higher order terms, and lower boundary given by either $y = 0$ or $y = a_ix^{n_i} +$ higher order terms, where $n_i > N_i$. Since $S_i(x, y) \sim x^{a_i}y^{b_i}$ on F_i , if $n_i > m_{ij} > N_i$ then on $N(S_i)$ the linear function $a + m_{ij}b$ is minimized at exactly one point, (a_i, b_i) . If $m_{ij} = N_i$ or n_i , then $a + m_{ij}b$ will be minimized at (a_i, b_i) and possibly also at other points in $N(S_i)$.

Next, suppose $f(x)$ is any function such that $f(x^K)$ is smooth for some K and such that the Taylor expansion of $f(x)$ has initial term $cx^{m_{ij}}$ for some c . Also suppose that $n_i > m_{ij} > N_i$. Then the Newton polygon of $S_i(x, y - f(x))$ will have an edge of slope $-\frac{1}{m_{ij}}$ containing the point $(a_i + m_{ij}b_i, 0)$ and no edges with lesser slope (i.e. no more horizontal edges). This fact implies that $S_i(x, y - f(x)) \sim x^{a_i+m_{ij}b_i}$ on V_{ijk}^2 , since $|y| < |x|^{m_{ij}+\eta}$ on V_{ijk}^2 . Note that the same is true if $m_{ij} = N_i$, if c is small enough, or if $m_{ij} = n_i$, if c is large enough. For in these cases the vertices other than (a_i, b_i) minimizing

$a + m_{ij}b$ do not interfere. Consequently, we may assume that $S'_i(x, y) = S_i \circ \phi_l(x, y)$ and its Newton polygon has the above properties. For $S'_i = S_i \circ \phi_l(x, y)$ is either of the form $S_i(x, y - f(x))$ or $S_i(x, -y - f(x))$ for an $f(x)$ of this type. (In the case where $m_{ij} = n_i$ or N_i , if the quantity called ξ in the proof of the resolution of singularities algorithm that was originally applied to $S_i(x, y)$ was small enough, which we may assume, then the coefficient a_i will be large enough and the coefficient A_i will be small enough for this to work.)

Let $f'_i = f_i \circ \phi_l$ and $R'_i = R_i \circ \phi_l$, so that $R'_i = S'_i + tf'_i$. We now will estimate the various $M_{R'_i, E_l}(\epsilon)$ and show they satisfy estimates better than those that we need. We start with E_l satisfying part c) of Theorem 3.1; that is, when $f'_i(x, y) \sim x^{\alpha_i} y^{\beta_i}$ with $\beta_i > 0$. Define X_l to be the set of points in E_l for which $|R'_i(x, y)| > \frac{1}{2}|S'_i(x, y)|$, and let $Y_l = E_l - X_l$. Then we have

$$\begin{aligned} |\{(x, y) \in X_l : |R'_i(x, y)| < \epsilon\}| &< |\{(x, y) \in X_l : |S'_i(x, y)| < 2\epsilon\}| \\ &\leq M_{S'_i, E_l}(\epsilon) \end{aligned} \quad (5.29)$$

This is better than the estimate we need, so we focus our attention on Y_l . Note that on Y_l we have

$$\frac{1}{2}|S'_i(x, y)| \leq |tf'_i(x, y)| \leq \frac{3}{2}|S'_i(x, y)| \quad (5.30)$$

Part c) of Theorem 3.1 says that, modulo a function vanishing to infinite order at $(0, 0)$, $f'_i(x, y) \sim x^{\alpha_i} y^{\beta_i}$ on E_l , while by the above discussion $S'_i(x, y) \sim x^{a_i + m_{ij}b_i}$ on E_l . Thus when (5.30) holds we have $x^{\alpha_i} y^{\beta_i} \sim x^{a_i + m_{ij}b_i}$. Next, note that by Theorem 3.1c), for any K one has

$$\begin{aligned} |\partial_y f'_i(x, y)| &> C_0 x^{\alpha_i} y^{\beta_i - 1} - O(x^K) > C_1 x^{a_i + m_{ij}b_i} y^{-1} - O(x^K) \\ &> C_2 x^{a_i + m_{ij}b_i - m_{ij} - \eta} \end{aligned} \quad (5.31)$$

The last inequality follows from the fact that $|y| < |x|^{m_{ij} + \eta}$ on E_l . On the other hand, since $S'_i(x, y)$'s Newton polygon has an edge of slope $-\frac{1}{m_{ij}}$ containing $(a_i + m_{ij}b_i, 0)$, and no other edges more horizontal than this one, we also have

$$|\partial_y S'_i(x, y)| < C_3 x^{a_i + m_{ij}b_i - m_{ij}} \quad (5.32)$$

Thus in taking the y derivative of $R'_i(x, y) = S'_i(x, y) + tf'_i(x, y)$, the derivative of the second term dominates and (for $t \neq 0$) we have

$$|\partial_y R'_i(x, y)| > C_4 x^{a_i + m_{ij}b_i - m_{ij} - \eta} \quad (5.33)$$

Next, we apply Theorem 2.2 and for some $x_0 > 0$ get that

$$\begin{aligned} |\{(x, y) \in Y_l : |R'_i(x, y)| < \epsilon\}| &\leq 4|\{(x, y) \in Y_l : C_4 x^{a_i + m_{ij}b_i - m_{ij} - \eta} y < \epsilon\}| \\ &\leq |\{(x, y) : 0 < x < x_0, |y| < x^{m_{ij} + \eta}, C_5 x^{a_i + m_{ij}b_i - m_{ij} - \eta} y < \epsilon\}| \end{aligned} \quad (5.34)$$

The last inequality again uses that $|y| < |x|^{m_{ij}+\eta}$ on E_l . Equation (5.34) can be bounded with the help of Lemma 2.3. If part a) or b) applies, it is bounded by $C_6\epsilon|\ln \epsilon|$, better than the estimate we need since we are assuming $d > 1$. If part c) applies, we get

$$|\{(x, y) \in Y_l : |R'_i(x, y)| < \epsilon\}| < C_7\epsilon^{\frac{m_{ij}+\eta+1}{a_i+m_{ij}b_i}} \quad (5.35)$$

Since $\eta > 0$, this exponent is better than $\frac{m_{ij}+1}{a_i+m_{ij}b_i}$, the reciprocal of the ordinate of the intersection of the bisectrix with the line of slope $-\frac{1}{m_{ij}}$ containing (a_i, b_i) . By (5.11), this ordinate is at most d . Hence the exponent on the right-hand side of (5.35) is greater than $\frac{1}{d}$, better than the estimate that we need.

We now move to the case where part b) of Theorem 3.1 is satisfied; that is, we assume $\beta_l = 0$ and thus $f'_i(x, y) \sim x^{\alpha_l}$ on E_l . If α_l were less than $a_i + m_{ij}b_i$, for $t \neq 0$ we would have $R'_i(x, y) = S'_i(x, y) + tf'_i(x, y) \sim x^{\alpha_l}$ on E_l as well. Thus for small enough x (which we may assume by making the radius of the original disk D sufficiently small) we would have $|R'_i(x, y)| \geq |S'_i(x, y)|$, and thus $M_{R'_i, E_l}(\epsilon) \leq M_{S'_i, E_l}(\epsilon)$, better than the estimate that we need. Similarly, if α_l were greater than $a_i + m_{ij}b_i$, $f'_i(x)$ would be small compared to $S'_i(x, y)$ and thus when x is sufficiently small we have $|R'_i(x, y)| \geq \frac{1}{2}|S'_i(x, y)|$, once again giving an estimate better than the one we need. Hence in the following we assume that $\alpha_l = a_i + m_{ij}b_i$.

Next, note that since we are in the setting of part b) of Theorem 3.1, the upper boundary of E_l has equation $y = bx^p +$ higher order terms, where $p \geq m_{ij} + \eta$. The lower boundary of E_l is the x -axis. Since $f'_i(x, y) \sim x^{a_i+m_{ij}b_i}$ on all of E_l , the Newton polygon of f'_i contains the vertex $(a_i + m_{ij}b_i, 0)$, and has no edge of slope greater than $-\frac{1}{p}$.

We now look at $S'_i(x, x^p y)$ and $f'_i(x, x^p y)$. First, in analogy with (5.13a) – (5.13b) we write

$$f'_i(x, y) = \sum_{a+pb=a_i+m_{ij}b_i} f'_{ab}x^a y^b + \sum_{a+pb>a_i+m_{ij}b_i, a,b<M} f'_{ab}x^a y^b + O(|x|^M + |y|^M) \quad (5.36a)$$

$$S'_i(x, y) = \sum_{a+pb=a_i+m_{ij}b_i} s'_{ab}x^a y^b + \sum_{a+pb>a_i+m_{ij}b_i, a,b<M} s'_{ab}x^a y^b + O(|x|^M + |y|^M) \quad (5.36b)$$

It is worth pointing out that since $N(S'_i)$ has no edge of slope greater than $-\frac{1}{m_{ij}}$, the first sum in (5.36b) has only one term. Next, in analogy with (5.14a) – (5.14b), we have

$$f'_i(x, x^p y) = x^{a_i+m_{ij}b_i} (f'_i)_{e_p}(1, y) + x^{a_i+m_{ij}b_i+\zeta} P'(x, y) + O(|x|^{M'}) \quad (5.37a)$$

$$S'_i(x, x^p y) = x^{a_i+m_{ij}b_i} (S'_i)_{e_p}(1, y) + x^{a_i+m_{ij}b_i+\zeta} Q'(x, y) + O(|x|^{M'}) \quad (5.37b)$$

Here $(f'_i)_{e_p}(x, y)$ denotes the sum of all terms $f'_{ab}x^a y^b$ of the Taylor expansion of $f'_i(x, y)$ at the origin with $a + pb = a_i + m_{ij}b_i$ (the minimal possible value of $a + pb$), and similarly

for $(S'_i)_{e_p}(x, y)$. Note that $(S'_i)_{e_p}(1, y)$ is constant here. Hence for all but finitely many t , $(S'_i)_{e_p}(1, y) + t(f'_i)_{e_p}(1, y)$ has no zeroes of order greater than one (this is actually true in more general scenarios as well.) Hence excluding these values of t , we write

$$R'_i(x, x^p y) = x^{a_i+m_{ij}b_i}(R'_i)_{e_p}(1, y) + x^{a_i+m_{ij}b_i+\zeta}R'(x, y) + O(|x|^{M'}) \quad (5.38)$$

Here $(R'_i)_{e_p}(1, y)$ has no zeroes of order greater than one. The y -range in (5.38) is contained in $[0, b + \epsilon]$ for any $\epsilon > 0$. (Recall b is such that the upper boundary of E_l is given by $y = bx^p + \dots$). We write $[0, b + \epsilon]$ as a union $B_1 \cup \dots \cup B_k$, where on each B_k either $(R'_i)_{e_p}(1, y)$ is nonvanishing, or $\partial_y(R'_i)_{e_p}(1, y)$ is nonvanishing.

Consider the case of a B_k for which $(R'_i)_{e_p}(1, y)$ is nonvanishing. Then on the domain of (5.38), we have $R'_i(x, x^p y) > C_1 x^{a_i+m_{ij}b_i}$. Translating back into the coordinates of E_l , we have $R'_i(x, y) > C_1 x^{a_i+m_{ij}b_i}$ on a subset A_k of $\{(x, y) : 0 < x < x_0, 0 < y < (b+1)x^p\}$ for some $x_0 > 0$. Thus we have

$$M_{R'_i, A_k}(\epsilon) = |\{(x, y) \in A_k : |R'_i(x, y)| < \epsilon\}| < C_2 \epsilon^{\frac{1+p}{a_i+m_{ij}b_i}} \quad (5.39)$$

Since $p > m_{ij}$, this exponent is better than $\frac{1+m_{ij}}{a_i+m_{ij}b_i} \geq \frac{1}{d}$. Thus (5.39) is better than the exponent we need. Now consider the case of a B_k for which $\partial_y((R'_i)_{e_p}(1, y))$ is nonvanishing. In this case, the relevant analogue of (5.38) is

$$(\partial_y R'_i)(x, x^p y) = x^{a_i+m_{ij}b_i-p} \partial_y((R'_i)_{e_p}(1, y)) + x^{a_i+m_{ij}b_i-p+\zeta} R''(x, y) + O(|x|^{M'}) \quad (5.40)$$

Hence on the domain of (5.40) we have $|(\partial_y R'_i)(x, x^p y)| > C_3 x^{a_i+m_{ij}b_i-p}$ if x is sufficiently small, which we may assume. Translating back into the original coordinates of E_l , this time we get that $\partial_y R'_i(x, y) > C_1 x^{a_i+m_{ij}b_i-p}$ on a subset A_k of $\{0 < x < x_0, 0 < y < (b+1)x^p\}$. Thus we may apply Lemma 2.2 and say that

$$\begin{aligned} M_{R'_i, A_k}(\epsilon) &= |\{(x, y) \in A_k : |R'_i(x, y)| < \epsilon\}| < 4 |\{(x, y) \in A_k : C_4 x^{a_i+m_{ij}b_i-p} y < \epsilon\}| \\ &\leq 4 |\{(x, y) \in E_l : C_4 x^{a_i+m_{ij}b_i-p} y < \epsilon\}| \end{aligned} \quad (5.41)$$

Since E_l is a subset of $\{(x, y) : 0 < x < x_0, 0 < y < (b+1)x^p\}$ for some $x_0 > 0$, we may use Lemma 2.3 to estimate the right-hand side of (5.41). If part a) or b) of the lemma applies, we get that $|M_{R'_i, A_k}(\epsilon)| < C\epsilon |\ln \epsilon|$, better than the estimate we need. If part c) applies, we get

$$M_{R'_i, A_k}(\epsilon) < C_5 \epsilon^{\frac{1+p}{a_i+m_{ij}b_i}} \quad (5.42)$$

This exponent is the same as that of (5.39), which we saw is better than what we need. This completes the proofs of Theorem 1.1 and 1.2.

6. References.

[AGV] V. Arnold, S Gusein-Zade, A Varchenko, *Singularities of differentiable maps*, Volume II, Birkhauser, Basel, 1988.

- [CaCWr] A. Carbery, M. Christ, J. Wright, *Multidimensional van der Corput and sublevel set estimates.*, J. Amer. Math. Soc. **12** (1999), no. 4, 981-1015.
- [G1] M. Greenblatt, *The asymptotic behavior of degenerate oscillatory integrals in two dimensions*, to appear, J. Funct. Anal.
- [G2] M. Greenblatt, *Resolution of singularities, asymptotic expansions of integrals over sublevel sets, and applications*, submitted.
- [G3] M. Greenblatt, *A direct resolution of singularities for functions of two variables with applications to analysis*, J. Anal. Math. **92** (2004), 233-257.
- [G4] M. Greenblatt, *Sharp L^2 estimates for one-dimensional oscillatory integral operators with C^∞ phase*. Amer. J. Math. **127** (2005), no. 3, 659-695.
- [H] L. Hörmander, *The analysis of linear partial differential operators. I. Distribution theory and Fourier analysis*, 2nd ed. Springer-Verlag, Berlin, (1990). xii+440 pp.
- [IKeM] I. Ikromov, M. Kempe, and D. Müller, *Sharp L^p estimates for maximal operators associated to hypersurfaces in \mathbf{R}^3 for $p > 2$* , preprint.
- [IoSa] A. Iosevich, E. Sawyer, *Oscillatory integrals and maximal averages over homogeneous surfaces*, Duke Math. J. **82** no. 1 (1996), 103-141.
- [K1] V. N. Karpushkin, *A theorem concerning uniform estimates of oscillatory integrals when the phase is a function of two variables*, J. Soviet Math. **35** (1986), 2809-2826.
- [K2] V. N. Karpushkin, *Uniform estimates of oscillatory integrals with parabolic or hyperbolic phases*, J. Soviet Math. **33** (1986), 1159-1188.
- [K3] V. N. Karpushkin, *Uniform estimates for volumes*, Tr. Math. Inst. Steklova **221** (1998), 225-231.
- [PS] D. H. Phong, E. M. Stein, *The Newton polyhedron and oscillatory integral operators*, Acta Mathematica **179** (1997), 107-152.
- [PSSSt] D. H. Phong, E. M. Stein, J. Sturm, *On the growth and stability of real-analytic functions*, Amer. J. Math. **121** (1999), no. 3, 519-554.
- [R] K. M. Rogers, *Sharp Van der Corput estimates and minimal divided differences*, Proc. Amer. Math. Soc. **133** (2005), no. 12, 3543-3550.
- [V] A. N. Varchenko, *Newton polyhedra and estimates of oscillatory integrals*, Functional Anal. Appl. **18** (1976), no. 3, 175-196.

Department of Mathematics, Statistics, and Computer Science
 University of Illinois at Chicago
 322 Science and Engineering Offices
 851 S. Morgan Street
 Chicago, IL 60607-7045