

1           **JOINT GEOMETRY/FREQUENCY ANALYTICITY OF FIELDS**  
2           **SCATTERED BY PERIODIC LAYERED MEDIA\***

3                           MATTHEW KEHOE   AND DAVID P. NICHOLLS †

4           **Abstract.** The scattering of linear waves by periodic structures is a crucial phenomena in many  
5 branches of applied physics and engineering. In this paper we establish rigorous analytic results neces-  
6 sary for the proper numerical analysis of a class of High-Order Perturbation of Surfaces/Asymptotic  
7 Waveform Evaluation (HOPS/AWE) methods for numerically simulating scattering returns from  
8 periodic diffraction gratings. More specifically, we prove a theorem on existence and uniqueness of  
9 solutions to a system of partial differential equations which model the interaction of linear waves with  
10 a periodic two-layer structure. Furthermore, we establish joint analyticity of these solutions with  
11 respect to both geometry and frequency perturbations. This result provides hypotheses under which  
12 a rigorous numerical analysis could be conducted on our recently developed HOPS/AWE algorithm.

13           **Key words.** High-Order Perturbation of Surfaces Methods; Layered media; Linear wave scat-  
14 tering; Helmholtz equation; Diffraction gratings.

15           **AMS subject classifications.** 65N35, 78A45, 78B22

16           **1. Introduction.** The scattering of linear waves by periodic structures is a cen-  
17 tral model in many problems of scientific and engineering interest. Examples arise in  
18 areas such as geophysics [64, 8], imaging [48], materials science [28], nanoplasmonics  
19 [61, 44, 24], and oceanography [10]. In the case of nanoplasmonics there are many  
20 such topics, for instance, extraordinary optical transmission [23], surface enhanced  
21 spectroscopy [47], and surface plasmon resonance (SPR) biosensing [31, 33, 42, 35].  
22 In all of these physical problems it is necessary to approximate scattering returns in  
23 a fast, robust, and highly accurate fashion.

24           The most popular approaches to solving these problems numerically in the en-  
25 gineering literature are *volumetric* methods. These include formulations based on  
26 the Finite Difference [40], Finite Element [34], Discontinuous Galerkin [30], Spectral  
27 Element [20], and Spectral Methods [29, 9, 63]. However, these methods suffer from  
28 the requirement that they discretize the full volume of the problem domain which  
29 results in an unnecessarily large number of degrees of freedom for a periodic *layered*  
30 structure. There is also the additional difficulty of approximating far-field boundary  
31 conditions explicitly [7].

32           For these reasons, *surface* methods are an appealing alternative, and we advocate  
33 the use of Boundary Integral Methods (BIM) [17, 37, 62] or High-Order Perturbation  
34 of Surfaces (HOPS) Methods [45, 46, 11, 12, 13, 54, 56]. Regarding the latter, we  
35 mention the classical Methods of Operator Expansions [45, 46] and Field Expansions  
36 [11, 12, 13], as well as the stabilized Method of Transformed Field Expansions [54, 56].  
37 All of these surface methods are greatly advantaged over the volumetric algorithms  
38 discussed above primarily due to the greatly reduced number of degrees of freedom  
39 that they require. Additionally the *exact* enforcement of the far-field boundary condi-  
40 tions is assured for both BIM and HOPS approaches. Consequently, these approaches  
41 are a favorable alternative and are becoming more widely used by practitioners.

42           There has been a large amount of not only rigorous analysis of systems of partial  
43 differential equations which model these scattering phenomena, but also careful design

---

\*D.P.N. gratefully acknowledges support from the National Science Foundation through Grants No. DMS-1813033 and DMS-2111283.

†Department of Mathematics, Statistics, and Computer Science, University of Illinois at Chicago, Chicago, IL 60607 ([mkehoe5@uic.edu](mailto:mkehoe5@uic.edu), [davidn@uic.edu](mailto:davidn@uic.edu))

44 of numerical schemes to simulate solutions of these. Most of these results utilize either  
 45 Integral Equation techniques or weak formulations of the volumetric problem, each  
 46 of which lead to a variety of natural numerical implementations. We recommend  
 47 the Habilitationsschrift of T. Arens [3] as a definitive reference for periodic layered  
 48 media problems in two and three dimensions. In particular, we refer the interested  
 49 reader to Chapter 1 which discusses in great detail the state-of-the-art in uniqueness  
 50 and existence results for scattering problems on biperiodic structures. For the two  
 51 dimensional problem we further refer the reader to the work of Petit [59]; Bao, Cowsar,  
 52 and Masters [5]; and Wilcox [65]. In three dimensions, results on the Helmholtz  
 53 equation can be found in Abboud and Nedelec [1]; Bao [4]; Bao, Dobson, and Cox  
 54 [6]; and Dobson [22]. In the context of Maxwell’s equations, we point out the work  
 55 of Chen and Friedman [16], and Dobson and Friedman [21]. Of course the field has  
 56 progressed from these classical contributions in a number of directions, and survey  
 57 volumes like [5] give further details.

58 Oftentimes in applications it is important to consider families of gratings interro-  
 59 gated over a range of illumination frequencies. An example of this is the computation  
 60 of the Reflectivity Map,  $R$ , which records the energy scattered by a layered structure  
 61 with interface shaped by  $z = g(x)$  and illuminated by radiation of frequency  $\omega$  (see,  
 62 e.g., [39]). Taking the point of view that this configuration is simply one in a family  
 63 with interface

$$64 \quad z = \varepsilon f(x), \quad \varepsilon \in \mathbf{R}, \quad \varepsilon \ll 1,$$

65 illuminated by radiation of frequency

$$66 \quad \omega = \underline{\omega} + \delta \underline{\omega}, \quad \delta \in \mathbf{R}, \quad \delta \ll 1,$$

67 where  $\underline{\omega}$  is a distinguished frequency of interest, our novel High-Order Perturbation  
 68 of Surfaces/Asymptotic Waveform Evaluation (HOPS/AWE) method [50, 36] is a  
 69 compelling numerical algorithm. In short, this scheme studies a *joint* Taylor expansion  
 70 of the solutions of the scattering problem in both  $\varepsilon$  and  $\delta$ . Upon insertion of this  
 71 expansion into relevant governing equations, the resulting recursions can be solved  
 72 up to a prescribed number of Taylor orders *once* and then simply summed for  $(\varepsilon, \delta)$   
 73 many times. Clearly, this is a most efficient and accurate method for approximating  
 74  $R = R(\varepsilon, \delta)$ , as we have demonstrated in our previous work [50, 36], provided that this  
 75 joint expansion can be justified. The point of the current contribution is to provide  
 76 this justification in the language of rigorous analysis (see Theorem 4.6). Not only is  
 77 this of intrinsic interest, but it also provides hypotheses and estimates as the starting  
 78 point for a rigorous numerical analysis of our HOPS/AWE scheme (see, e.g., [57] for  
 79 a possible path) for this problem.

80 The paper is organized as follows: In Section 2 we summarize the equations which  
 81 govern the propagation of linear waves in a two-dimensional periodic structure, and  
 82 in Section 2.1 we discuss how the outgoing wave conditions can be exactly enforced  
 83 through the use of Transparent Boundary Conditions. Then in Section 3 we restate  
 84 our governing equations in terms of interfacial quantities via a Non-Overlapping Do-  
 85 main Decomposition phrased in terms of Dirichlet-Neumann Operators (DNOs). In  
 86 Section 4 we discuss our analyticity result with a general theory in Section 4.1 and  
 87 our specific result in Section 4.2. This requires a study of analyticity of the data in  
 88 Section 4.3 and an investigation of the flat-interface situation in Section 4.4. We con-  
 89 clude with the final piece required for the general theory: The analyticity of Dirichlet-  
 90 Neumann Operators (Section 6). We accomplish this by first establishing analyticity

91 of the underlying fields (Section 5) requiring a special change of variables specified  
 92 in Section 5.1. With this we demonstrate the analyticity of the scattered field in  
 93 Sections 5.2 and 5.3. Given these theorems, we prove the analyticity of the DNOs in  
 94 Section 6.

95 **2. The Governing Equations.** An example of the geometry we consider is  
 displayed in Figure 1: a  $y$ -invariant, doubly layered structure with a periodic interface




Fig. 1: A two-layer structure with a periodic interface,  $z = g(x)$ , separating two material layers,  $S^{(u)}$  and  $S^{(w)}$ , illuminated by plane-wave incidence.

96 separating the two materials. The interface is specified by the graph of the function  
 97  $z = g(x)$  which is  $d$ -periodic so that  $g(x+d) = g(x)$ . Dielectrics occupy both domains  
 98 where an insulator (with refractive index  $n^u$ ) fills the region above the graph  $z = g(x)$   
 99

$$100 \quad S^{(u)} := \{z > g(x)\},$$

101 and a second material (with index of refraction  $n^w$ ) occupies

$$102 \quad S^{(w)} := \{z < g(x)\}.$$

103 The superscripts are chosen to conform to the notation of the authors in previous  
 104 work [49, 52]. The structure is illuminated from above by monochromatic plane-wave  
 105 incident radiation of frequency  $\omega$  and wavenumber  $k^u = n^u\omega/c_0 = \omega/c^u$  ( $c_0$  is the  
 106 speed of light) aligned with the grooves

$$107 \quad \underline{\mathbf{E}}^i(x, z, t) = \mathbf{A}e^{-i\omega t + i\alpha x - i\gamma^u z}, \quad \underline{\mathbf{H}}^i(x, z, t) = \mathbf{B}e^{-i\omega t + i\alpha x - i\gamma^u z},$$

$$108 \quad \alpha := k^u \sin(\theta), \quad \gamma^u := k^u \cos(\theta).$$

110 We consider the reduced incident fields

$$111 \quad \mathbf{E}^i(x, z) = e^{i\omega t} \underline{\mathbf{E}}^i(x, z, t), \quad \mathbf{H}^i(x, z) = e^{i\omega t} \underline{\mathbf{H}}^i(x, z, t),$$

112 where the time dependence  $\exp(-i\omega t)$  has been factored out. As shown in [59],  
 113 the reduced electric and magnetic fields, like the reduced scattered fields, are  $\alpha$ -  
 114 quasiperiodic due to the incident radiation. To close the problem, we specify that  
 115 the scattered radiation is “outgoing,” upward propagating in  $S^{(u)}$  and downward  
 116 propagating in  $S^{(w)}$ .

117 It is well known (see, e.g., Petit [59]) that in this two-dimensional setting, the  
 118 time-harmonic Maxwell equations decouple into two scalar Helmholtz problems which  
 119 govern the Transverse Electric (TE) and Transverse Magnetic (TM) polarizations.  
 120 We define the invariant ( $y$ ) direction of the scattered (electric or magnetic) field by  
 121  $\tilde{u} = \tilde{u}(x, z)$  and  $\tilde{w} = \tilde{w}(x, z)$  in  $S^{(u)}$  and  $S^{(w)}$ , respectively. The incident radiation in  
 122 the upper field is denoted by  $\tilde{u}^i(x, z)$ .

123 Following our previous work [50] we further factor out the phase  $\exp(i\alpha x)$  from  
 124 the fields  $\tilde{u}$  and  $\tilde{w}$

$$125 \quad u(x, z) = e^{-i\alpha x} \tilde{u}(x, z), \quad w(x, z) = e^{-i\alpha x} \tilde{w}(x, z),$$

126 which, we note, are  $d$ -periodic. In light of all of this, we are led to seek outgoing,  
 127  $d$ -periodic solutions of

$$128 \quad (2.1a) \quad \Delta u + 2i\alpha \partial_x u + (\gamma^u)^2 u = 0, \quad z > g(x),$$

$$129 \quad (2.1b) \quad \Delta w + 2i\alpha \partial_x w + (\gamma^w)^2 w = 0, \quad z < g(x),$$

$$130 \quad (2.1c) \quad u - w = \zeta, \quad z = g(x),$$

$$131 \quad (2.1d) \quad \partial_N u - i\alpha(\partial_x g)u - \tau^2 [\partial_N w - i\alpha(\partial_x g)w] = \psi, \quad z = g(x),$$

133 where  $N := (-\partial_x g, 1)^T$ . The Dirichlet and Neumann data are

$$134 \quad (2.1e) \quad \zeta(x) := -e^{-i\gamma^u g(x)},$$

$$135 \quad (2.1f) \quad \psi(x) := (i\gamma^u + i\alpha(\partial_x g))e^{-i\gamma^u g(x)},$$

137 and

$$138 \quad \tau^2 = \begin{cases} 1, & \text{TE,} \\ (k^u/k^w)^2 = (n^u/n^w)^2, & \text{TM,} \end{cases}$$

139 where  $k^w = n^w \omega / c_0 = \omega / c^w$  and  $\gamma^w = k^w \cos(\theta)$ .

140 **2.1. Transparent Boundary Conditions.** The Rayleigh expansions, which  
 141 are derived through separation of variables [59], are the periodic, upward/downward  
 142 propagating solutions of (2.1a) and (2.1b). In order to truncate the bi-infinite problem  
 143 domain to one of finite size we use these to define Transparent Boundary Conditions.  
 144 For this we choose values  $a$  and  $b$  such that

$$145 \quad a > |g|_\infty, \quad -b < -|g|_\infty,$$

146 and define the artificial boundaries  $\{z = a\}$  and  $\{z = -b\}$ . In  $\{z > a\}$  the Rayleigh  
 147 expansions tell us that upward propagating solutions of (2.1a) are

$$148 \quad (2.2) \quad u(x, z) = \sum_{p=-\infty}^{\infty} \hat{a}_p e^{i\tilde{p}x + i\gamma_p^u z},$$

149 while downward propagating solutions of (2.1b) in  $\{z < -b\}$  can be expressed as

$$150 \quad w(x, z) = \sum_{p=-\infty}^{\infty} \hat{d}_p e^{i\tilde{p}x - i\gamma_p^w z},$$

151 where, for  $p \in \mathbf{Z}$  and  $q \in \{u, w\}$ ,

$$152 \quad (2.3) \quad \tilde{p} := \frac{2\pi p}{d}, \quad \alpha_p := \alpha + \tilde{p}, \quad \gamma_p^q := \begin{cases} \sqrt{(k^q)^2 - \alpha_p^2}, & p \in \mathcal{U}^q, \\ i\sqrt{\alpha_p^2 - (k^q)^2}, & p \notin \mathcal{U}^q, \end{cases}$$

153 and

$$154 \quad \mathcal{U}^q := \{p \in \mathbf{Z} \mid \alpha_p^2 < (k^q)^2\},$$

155 which are the propagating modes in the upper and lower layers. With these we can  
156 define the Transparent Boundary Conditions in the following way: we first rewrite  
157 (2.2) as

$$158 \quad u(x, z) = \sum_{p=-\infty}^{\infty} \left( \hat{a}_p e^{i\gamma_p^u a} \right) e^{i\tilde{p}x + i\gamma_p^u (z-a)} = \sum_{p=-\infty}^{\infty} \hat{\xi}_p e^{i\tilde{p}x + i\gamma_p^u (z-a)},$$

159 and observe that,

$$160 \quad u(x, a) = \sum_{p=-\infty}^{\infty} \hat{\xi}_p e^{i\tilde{p}x} =: \xi(x),$$

161 and

$$162 \quad \partial_z u(x, a) = \sum_{p=-\infty}^{\infty} (i\gamma_p^u) \hat{\xi}_p e^{i\tilde{p}x} =: T^u[\xi(x)],$$

163 which defines the order-one Fourier multiplier  $T^u$ . From this we state that upward-  
164 propagating solutions of (2.1a) satisfy the Transparent Boundary Condition at  $z = a$

$$165 \quad (2.4) \quad \partial_z u(x, a) - T^u[u(x, a)] = 0, \quad z = a.$$

166 A similar calculation leads to the Transparent Boundary Condition at  $z = -b$

$$167 \quad (2.5) \quad \partial_z w(x, -b) - T^w[w(x, -b)] = 0, \quad z = -b,$$

168 where

$$169 \quad T^w[\psi(x)] := \sum_{p=-\infty}^{\infty} (-i\gamma_p^w) \hat{\psi}_p e^{i\tilde{p}x}.$$

170 We note that these conditions enforce the Upward and Downward Propagating Con-  
171 ditions described by Arens [3].

172 With these we now state the full set of governing equations as

$$173 \quad (2.6a) \quad \Delta u + 2i\alpha\partial_x u + (\gamma^u)^2 u = 0, \quad z > g(x),$$

$$174 \quad (2.6b) \quad \Delta w + 2i\alpha\partial_x w + (\gamma^w)^2 w = 0, \quad z < g(x),$$

$$175 \quad (2.6c) \quad u - w = \zeta, \quad z = g(x),$$

$$176 \quad (2.6d) \quad \partial_N u - i\alpha(\partial_x g)u - \tau^2 [\partial_N w - i\alpha(\partial_x g)w] = \psi, \quad z = g(x),$$

$$177 \quad (2.6e) \quad \partial_z u(x, a) - T^u[u(x, a)] = 0, \quad z = a,$$

$$178 \quad (2.6f) \quad \partial_z w(x, -b) - T^w[w(x, -b)] = 0, \quad z = -b,$$

$$179 \quad (2.6g) \quad u(x + d, z) = u(x, z),$$

$$180 \quad (2.6h) \quad w(x + d, z) = w(x, z).$$

182 **3. A Non-Overlapping Domain Decomposition Method.** We now rewrite  
183 our governing equations (2.6) in terms of *surface* quantities via a Non-Overlapping  
184 Domain Decomposition Method [43, 19, 18]. For this we define

$$185 \quad U(x) := u(x, g(x)), \quad \tilde{U}(x) := -\partial_N u(x, g(x)),$$

$$186 \quad W(x) := w(x, g(x)), \quad \tilde{W}(x) := \partial_N w(x, g(x)),$$

188 where  $u$  is a  $d$ -periodic solution of (2.6a) and (2.6e), and  $w$  is a  $d$ -periodic solution of  
189 (2.6b) and (2.6f). In terms of these, our full governing equations (2.6) are equivalent  
190 to the pair of boundary conditions, (2.6c) and (2.6d),

$$191 \quad (3.1a) \quad U - W = \zeta,$$

$$192 \quad (3.1b) \quad -\tilde{U} - (i\alpha)(\partial_x g)U - \tau^2 [\tilde{W} - (i\alpha)(\partial_x g)W] = \psi.$$

194 This set of two equations and four unknowns can be closed by noting that the pairs  
195  $\{U, \tilde{U}\}$  and  $\{W, \tilde{W}\}$  are connected, e.g., by Dirichlet-Neumann Operators (DNOs),  
196 which [56] showed are well-defined under the hypotheses presently listed.

197 **DEFINITION 3.1.** *Given an integer  $s \geq 0$ , if  $g \in C^{s+2}$  then the unique solution of*  
198

$$199 \quad (3.2a) \quad \Delta u + 2i\alpha\partial_x u + (\gamma^u)^2 u = 0, \quad z > g(x),$$

$$200 \quad (3.2b) \quad u = U, \quad z = g(x),$$

$$201 \quad (3.2c) \quad \partial_z u(x, a) - T^u[u(x, a)] = 0, \quad z = a,$$

$$202 \quad (3.2d) \quad u(x + d, z) = u(x, z),$$

204 *defines the upper layer DNO*

$$205 \quad (3.3) \quad G : U \rightarrow \tilde{U}.$$

206 **DEFINITION 3.2.** *Given an integer  $s \geq 0$ , if  $g \in C^{s+2}$  then the unique solution of*  
207

$$208 \quad (3.4a) \quad \Delta w + 2i\alpha\partial_x w + (\gamma^w)^2 w = 0, \quad z < g(x),$$

$$209 \quad (3.4b) \quad w = W, \quad z = g(x),$$

$$210 \quad (3.4c) \quad \partial_z w(x, -b) - T^w[w(x, -b)] = 0, \quad z = -b,$$

$$211 \quad (3.4d) \quad w(x + d, z) = w(x, z).$$

213 *defines the lower layer DNO*

$$214 \quad (3.5) \quad J : W \rightarrow \tilde{W}.$$

215 The interfacial reformulation of our governing equations (3.1) now becomes

$$216 \quad (3.6) \quad \mathbf{A}\mathbf{V} = \mathbf{R},$$

217 where

$$218 \quad (3.7) \quad \mathbf{A} = \begin{pmatrix} I & -I \\ G + (\partial_x g)(i\alpha) & \tau^2 J - \tau^2 (\partial_x g)(i\alpha) \end{pmatrix}, \quad \mathbf{V} = \begin{pmatrix} U \\ W \end{pmatrix}, \quad \mathbf{R} = \begin{pmatrix} \zeta \\ -\psi \end{pmatrix}.$$

219 **4. Joint Analyticity of Solutions.** There are many possible ways to analyze  
220 (3.6) rigorously. Following our recent work [36], we select a jointly perturbative ap-  
221 proach based on two smallness assumptions:

- 222 1. Boundary Perturbation:  $g(x) = \varepsilon f(x)$ ,  $\varepsilon \in \mathbf{R}$ ,  $\varepsilon \ll 1$ ,
- 223 2. Frequency Perturbation:  $\omega = (1 + \delta)\underline{\omega} = \underline{\omega} + \delta\underline{\omega}$ ,  $\delta \in \mathbf{R}$ ,  $\delta \ll 1$ .

224 We point out that possibly one or both of these smallness requirements can be relaxed,  
225 provided that the parameters ( $\varepsilon$  and/or  $\delta$ ) are real (c.f., [55, 58]). The frequency  
226 perturbation has the following important consequences

$$227 \quad k^q = \omega/c^q = (1 + \delta)\underline{\omega}/c^q =: (1 + \delta)\underline{k}^q = \underline{k}^q + \delta\underline{k}^q, \quad q \in \{u, w\},$$

$$228 \quad \alpha = k^u \sin(\theta) = (1 + \delta)\underline{k}^u \sin(\theta) =: (1 + \delta)\underline{\alpha} = \underline{\alpha} + \delta\underline{\alpha},$$

$$229 \quad \gamma^q = k^q \cos(\theta) = (1 + \delta)\underline{\gamma}^q \cos(\theta) =: (1 + \delta)\underline{\gamma}^q = \underline{\gamma}^q + \delta\underline{\gamma}^q, \quad q \in \{u, w\}.$$

231 This, in turn, delivers

$$232 \quad \alpha_p = \alpha + \tilde{p} = \underline{\alpha} + \delta\underline{\alpha} + \tilde{p} =: \underline{\alpha}_p + \delta\underline{\alpha}.$$

233 We now pursue this perturbative approach to establish the existence, uniqueness,  
234 and analyticity of solutions to (3.6). To accomplish this we will presently show the  
235 joint analytic dependence of  $\mathbf{A} = \mathbf{A}(\varepsilon, \delta)$  and  $\mathbf{R} = \mathbf{R}(\varepsilon, \delta)$  upon  $\varepsilon$  and  $\delta$ , and then  
236 appeal to the regular perturbation theory for linear systems of equations outlined in  
237 [51] to discover the analyticity of the unique solution  $\mathbf{V} = \mathbf{V}(\varepsilon, \delta)$ . More precisely,  
238 we view (3.6) as

$$239 \quad \mathbf{A}(\varepsilon, \delta)\mathbf{V}(\varepsilon, \delta) = \mathbf{R}(\varepsilon, \delta),$$

240 establish the analyticity of  $\mathbf{A}$  and  $\mathbf{R}$  so that

$$241 \quad (4.1) \quad \{\mathbf{A}, \mathbf{R}\}(\varepsilon, \delta) = \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \{\mathbf{A}_{n,m}, \mathbf{R}_{n,m}\} \varepsilon^n \delta^m,$$

242 and seek a solution of the form

$$243 \quad (4.2) \quad \mathbf{V}(\varepsilon, \delta) = \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \mathbf{V}_{n,m} \varepsilon^n \delta^m,$$

244 which we will show converges in a function space. To pursue this we insert (4.2) and  
245 (4.1) into (3.6) and find, at each perturbation order  $(n, m)$ , that we must solve

$$246 \quad \mathbf{A}_{0,0} \mathbf{V}_{n,m} = \mathbf{R}_{n,m} - \sum_{\ell=0}^{n-1} \mathbf{A}_{n-\ell,0} \mathbf{V}_{\ell,m} - \sum_{r=0}^{m-1} \mathbf{A}_{0,m-r} \mathbf{V}_{n,r}$$

$$247 \quad (4.3) \quad - \sum_{\ell=0}^{n-1} \sum_{r=0}^{m-1} \mathbf{A}_{n-\ell,m-r} \mathbf{V}_{\ell,r}.$$

248

249 A brief inspection of the formulas for  $\mathbf{A}$  and  $\mathbf{R}$ , (3.7), reveals that

$$250 \quad (4.4a) \quad \mathbf{A}_{0,0} = \begin{pmatrix} I & -I \\ G_{0,0} & \tau^2 J_{0,0} \end{pmatrix},$$

$$251 \quad \mathbf{A}_{n,m} = \begin{pmatrix} 0 & 0 \\ G_{n,m} & \tau^2 J_{n,m} \end{pmatrix}$$

$$252 \quad (4.4b) \quad + \delta_{n,1} \{1 + \delta_{m,1}\} (\partial_x f)(i\alpha) \begin{pmatrix} 0 & 0 \\ 1 & -\tau^2 \end{pmatrix}, \quad n \neq 0 \text{ or } m \neq 0,$$

$$253 \quad (4.4c) \quad \mathbf{R}_{n,m} = \begin{pmatrix} \zeta_{n,m} \\ -\psi_{n,m} \end{pmatrix},$$

254

255 where  $\delta_{p,q}$  is the Kronecker delta function. Formulas for the terms  $\{\zeta_{n,m}, \psi_{n,m}\}$  can  
 256 be found in [36] or by using the recursions described in Section 4.3. The terms  $G_{n,m}$   
 257 and  $J_{n,m}$  are the  $(n, m)$ -th corrections of the DNOs  $G$  and  $J$ , respectively, in a Taylor  
 258 series expansion of each jointly in  $\varepsilon$  and  $\delta$ . This is explained in Section 6, together  
 259 with precise estimates of the coefficients,  $G_{n,m}$  and  $J_{n,m}$ , in the appropriate Sobolev  
 260 spaces. Finally, in Section 4.4 we utilize expressions for the flat-interface DNOs,  $G_{0,0}$   
 261 and  $J_{0,0}$ , to investigate the mapping properties of the linearized operator,  $\mathbf{A}_{0,0}$ , and  
 262 its inverse.

263 **4.1. A General Analyticity Theory.** Given these estimates, existence, unique-  
 264 ness, and analyticity of solutions can be deduced in a rather straightforward fashion  
 265 using the following result from one of the authors' previous papers [51] (Theorem 3.2).  
 266 This result uses multi-index notation [25], in particular

$$267 \quad \tilde{\varepsilon} := \begin{pmatrix} \varepsilon_1 \\ \vdots \\ \varepsilon_M \end{pmatrix}, \quad \tilde{n} := \begin{pmatrix} n_1 \\ \vdots \\ n_M \end{pmatrix},$$

268 and the convention

$$269 \quad \sum_{\tilde{n}=0}^{\infty} A_{\tilde{n}} \tilde{\varepsilon}^{\tilde{n}} = \sum_{n_1=0}^{\infty} \cdots \sum_{n_M=0}^{\infty} A_{n_1, \dots, n_M} \varepsilon_1^{n_1} \cdots \varepsilon_M^{n_M}.$$

270

271 **THEOREM 4.1.** *Given two Banach spaces,  $\tilde{X}$  and  $\tilde{Y}$ , suppose that:*

- 272 1.  $\mathbf{R}_{\tilde{n}} \in \tilde{Y}$  for all  $\tilde{n} \geq 0$ , and there exist multi-indexed constants  $C_R > 0$ ,  
 273  $B_R > 0$  such that

$$274 \quad \|\mathbf{R}_{\tilde{n}}\|_{\tilde{Y}} \leq C_R B_R^{\tilde{n}},$$

- 275 2.  $\mathbf{A}_{\tilde{n}} : \tilde{X} \rightarrow \tilde{Y}$  for all  $\tilde{n} \geq 0$ , and there exist multi-indexed constants  $C_A > 0$ ,  
 276  $B_A > 0$  such that

$$277 \quad \|\mathbf{A}_{\tilde{n}}\|_{\tilde{X} \rightarrow \tilde{Y}} \leq C_A B_A^{\tilde{n}},$$

- 278 3.  $\mathbf{A}_0^{-1} : \tilde{Y} \rightarrow \tilde{X}$ , and there exists a constant  $C_e > 0$  such that

$$279 \quad \|\mathbf{A}_0^{-1}\|_{\tilde{Y} \rightarrow \tilde{X}} \leq C_e.$$



280 Then the equation (3.6) has a unique solution,

$$281 \quad (4.5) \quad \mathbf{V}(\tilde{\varepsilon}) = \sum_{\tilde{n}=0}^{\infty} \mathbf{V}_{\tilde{n}} \tilde{\varepsilon}^{\tilde{n}},$$

282 and there exist multi-indexed constants  $C_V > 0$  and  $B_V > 0$  such that

$$283 \quad \|\mathbf{V}_{\tilde{n}}\|_{\tilde{X}} \leq C_V B_V^{\tilde{n}},$$

284 for all  $\tilde{n} \geq 0$  and any

$$285 \quad C_V \geq 2C_e C_R, \quad B_V \geq \max\{B_R, 2B_A, 4C_e C_A B_A\},$$

286 enforced componentwise. This implies that, for any multi-indexed constant  $0 \leq \tilde{\rho} < 1$ ,  
287 (4.5), converges for all  $\tilde{\varepsilon}$  such that  $B\tilde{\varepsilon} < \tilde{\rho}$ , i.e.,  $\tilde{\varepsilon} < \tilde{\rho}/B$ .

288 *Remark 4.2.* In the current context we will use this result in the case  $M = 2$  and

$$289 \quad \tilde{\varepsilon} = \begin{pmatrix} \varepsilon \\ \delta \end{pmatrix}, \quad \tilde{n} = \begin{pmatrix} n \\ m \end{pmatrix}, \quad \tilde{\rho} = \begin{pmatrix} \rho \\ \sigma \end{pmatrix}.$$

290 **4.2. Analyticity of Solutions to the Two-Layer Problem.** To state our  
291 theorem precisely we briefly define and recall classical properties of the  $L^2$ -based  
292 Sobolev spaces,  $H^s$ , of laterally periodic functions [37]. We know that any  $d$ -periodic  
293  $L^2$  function can be expressed in a Fourier series as

$$294 \quad \mu(x) = \sum_{p=-\infty}^{\infty} \hat{\mu}_p e^{i\tilde{p}x}, \quad \hat{\mu}_p = \frac{1}{d} \int_0^d \mu(x) e^{-i\tilde{p}x},$$

295 [37]. We define the symbol  $\langle \tilde{p} \rangle^2 := 1 + |\tilde{p}|^2$  so that laterally periodic norms for surface  
296 and volumetric functions are defined by

$$297 \quad \|\mu\|_{H^s}^2 := \sum_{p=-\infty}^{\infty} \langle \tilde{p} \rangle^{2s} |\hat{\mu}_p|^2,$$

298 and

$$299 \quad \|u\|_{H^s}^2 := \sum_{\ell=0}^s \sum_{p=-\infty}^{\infty} \langle \tilde{p} \rangle^{2(s-\ell)} \int_0^a |\hat{u}_p(z)|^2 dz = \sum_{\ell=0}^s \sum_{p=-\infty}^{\infty} \langle \tilde{p} \rangle^{2(s-\ell)} \|\hat{u}_p\|_{L^2(0,a)}^2,$$

300 respectively. With these we define the laterally  $d$ -periodic Sobolev spaces  $H^s$  as the  
301  $L^2$  functions for which  $\|\cdot\|_{H^s}$  is finite. For our present use we define the vector-valued  
302 spaces for  $s \geq 0$

$$303 \quad X^s := \left\{ \mathbf{V} = \begin{pmatrix} U \\ W \end{pmatrix} \middle| U, W \in H^{s+3/2}([0, d]) \right\},$$

304 and

$$305 \quad Y^s := \left\{ \mathbf{R} = \begin{pmatrix} \zeta \\ -\psi \end{pmatrix} \middle| \zeta \in H^{s+3/2}([0, d]), \psi \in H^{s+1/2}([0, d]) \right\}.$$

306 These have the norms

$$307 \quad \|\mathbf{V}\|_{X^s}^2 = \left\| \begin{pmatrix} U \\ W \end{pmatrix} \right\|_{X^s}^2 := \|U\|_{H^{s+3/2}}^2 + \|W\|_{H^{s+3/2}}^2,$$

$$308 \quad \|\mathbf{R}\|_{Y^s}^2 = \left\| \begin{pmatrix} \zeta \\ -\psi \end{pmatrix} \right\|_{Y^s}^2 := \|\zeta\|_{H^{s+3/2}}^2 + \|\psi\|_{H^{s+1/2}}^2.$$

309

310 In addition to these function spaces we also require the following three results from  
 311 the classical theory of Sobolev spaces [2, 41] and elliptic partial differential equations  
 312 [38, 26, 27, 25]. (See also [53, 32] in the context of HOPS methods.)

313 LEMMA 4.3. *Given an integer  $s \geq 0$  and any  $\eta > 0$ , there exists a constant*  
 314  *$\mathcal{M} = \mathcal{M}(s)$  such that if  $f \in C^s([0, d])$  and  $u \in H^s([0, d] \times [0, a])$  then*

$$315 \quad (4.6) \quad \|fu\|_{H^s} \leq \mathcal{M} |f|_{C^s} \|u\|_{H^s},$$

316 *and if  $\tilde{f} \in C^{s+1/2+\eta}([0, d])$  and  $\tilde{u} \in H^{s+1/2}([0, d])$  then*

$$317 \quad (4.7) \quad \|\tilde{f}\tilde{u}\|_{H^{s+1/2}} \leq \mathcal{M} |\tilde{f}|_{C^{s+1/2+\eta}} \|\tilde{u}\|_{H^{s+1/2}}.$$

318 THEOREM 4.4. *Given an integer  $s \geq 0$ , if  $F \in H^s([0, d] \times [0, a])$ ,  $U \in H^{s+3/2}([0, d])$ ,*  
 319  *$P \in H^{s+1/2}([0, d])$ , then the unique solution of*

$$320 \quad \Delta u(x, z) + 2i\underline{\alpha}\partial_x u(x, z) + (\underline{\gamma}^u)^2 u(x, z) = F(x, z), \quad 0 < z < a,$$

$$321 \quad u(x, 0) = U(x, 0), \quad z = 0,$$

$$322 \quad \partial_z u(x, a) - T^u[u(x, a)] = P(x), \quad z = a,$$

$$323 \quad u(x + d, z) = u(x, z),$$

324

325 *satisfies*

$$326 \quad (4.8) \quad \|u\|_{H^{s+2}} \leq C_e \{ \|F\|_{H^s} + \|U\|_{H^{s+3/2}} + \|P\|_{H^{s+1/2}} \},$$

327 *for some constant  $C_e > 0$ .*

328 LEMMA 4.5. *Given an integer  $s \geq 0$ , if  $F \in H^s([0, d] \times [0, a])$ , then  $(a - z)F \in$   
 329  $H^s([0, d] \times [0, a])$  and there exists a positive constant  $Z_a = Z_a(s)$  such that*

$$330 \quad \|(a - z)F\|_{H^s} \leq Z_a \|F\|_{H^s}.$$

331 We now state our main result.

332 THEOREM 4.6. *Given an integer  $s \geq 0$ , if  $f \in C^{s+2}([0, d])$  then the equation (3.6)  
 333 has a unique solution, (4.2), and there exist constants  $B, C, D > 0$  such that*

$$334 \quad \|\mathbf{V}_{n,m}\|_{X^s} \leq CB^n D^m,$$

335 *for all  $n, m \geq 0$ . This implies that for any  $0 \leq \rho, \sigma < 1$ , (4.2) converges for all  $\varepsilon$  such  
 336 that  $B\varepsilon < \rho$ , i.e.,  $\varepsilon < \rho/B$  and all  $\delta$  such that  $D\delta < \sigma$ , i.e.,  $\delta < \sigma/D$ .*

337 *Proof.* As mentioned above, our strategy is to invoke Theorem 4.1 and thus we  
 338 must verify its hypotheses. To begin, we consider the spaces

$$339 \quad \tilde{X} = X^s, \quad \tilde{Y} = Y^s.$$

340 In Section 4.3 we will show that the vector  $\mathbf{R}_{n,m}$ , consisting of  $\zeta_{n,m}$  and  $\psi_{n,m}$ , is  
 341 bounded in  $Y^s$  for any  $s \geq 0$  provided that  $f \in C^{s+2}([0, d])$ . (This implies that the  
 342  $\mathbf{R}_{n,m}$  satisfies the estimates of Item 1 in Theorem 4.1.)

343 Then in Section 6 we show that the operators  $G_{n,m}$  and  $J_{n,m}$  in the Taylor series  
 344 expansions of the DNOs satisfy appropriate bounds provided that  $f \in C^{s+2}([0, d])$ .  
 345 With this, it is clear that the  $\mathbf{A}_{n,m}$  satisfy the estimates of Item 2 in Theorem 4.1.

346 Finally, in Section 4.4 we show that the estimates and mapping properties of  $\mathbf{A}_{0,0}^{-1}$   
 347 for Item 3 in Theorem 4.1 hold.  $\square$

348 **4.3. Analyticity of the Surface Data.** To establish the analyticity of the  
 349 Dirichlet and Neumann data we begin by defining

$$350 \quad \mathcal{E}(x; \varepsilon, \delta) := e^{-i(1+\delta)\underline{\gamma}^u \varepsilon f(x)},$$

351 and note that we can write (2.1e) and (2.1f) as

$$352 \quad \zeta(x) = \zeta(x; \varepsilon, \delta) = -\mathcal{E}(x; \varepsilon, \delta),$$

$$353 \quad \psi(x) = \psi(x; \varepsilon, \delta) = \{i(1+\delta)\underline{\gamma}^u + i(1+\delta)\underline{\alpha}(\varepsilon \partial_x f)\} \mathcal{E}(x; \varepsilon, \delta).$$

355 We will now demonstrate that the function  $\mathcal{E}$  is jointly analytic in  $\varepsilon$  and  $\delta$ , which  
 356 clearly demonstrates the joint analytic dependence of the data,  $\zeta(x; \varepsilon, \delta)$  and  $\psi(x; \varepsilon, \delta)$ .

357 **LEMMA 4.7.** *Given any integer  $s \geq 0$ , if  $f \in C^{s+2}([0, d])$  then the function*  
 358  *$\mathcal{E}(x; \varepsilon, \delta)$  is jointly analytic in  $\varepsilon$  and  $\delta$ . Therefore*

$$359 \quad (4.9) \quad \mathcal{E}(x; \varepsilon, \delta) = \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \mathcal{E}_{n,m}(x) \varepsilon^n \delta^m,$$

360 and, for constants  $C_{\mathcal{E}}, B_{\mathcal{E}}, D_{\mathcal{E}} > 0$ ,

$$361 \quad (4.10) \quad \|\mathcal{E}_{n,m}\|_{H^{s+3/2}} \leq C_{\mathcal{E}} B_{\mathcal{E}}^n D_{\mathcal{E}}^m,$$

362 for all  $n, m \geq 0$ .

363 *Proof.* By evaluating at  $\varepsilon = 0$  we find that

$$364 \quad \mathcal{E}(x; 0, \delta) = 1,$$

365 so that

$$366 \quad \mathcal{E}_{0,m}(x) = \begin{cases} 1, & m = 0, \\ 0, & m > 0. \end{cases}$$

367 For  $\varepsilon > 0$  we use the straightforward computation

$$368 \quad \partial_{\varepsilon} \mathcal{E} = \{-i(1+\delta)\underline{\gamma}^u f\} \mathcal{E},$$

369 and the expansion (4.9) to learn that, for  $m = 0$ ,

$$370 \quad (4.11) \quad \mathcal{E}_{n+1,0} = \left( \frac{-i\underline{\gamma}^u f}{n+1} \right) \mathcal{E}_{n,0},$$

371 and, for  $m > 0$ ,

$$372 \quad (4.12) \quad \mathcal{E}_{n+1,m} = \left( \frac{-i\underline{\gamma}^u f}{n+1} \right) \{\mathcal{E}_{n,m} + \mathcal{E}_{n,m-1}\}.$$

373 We work by induction in  $n$  and begin by establishing (4.10) at  $n = 0$  for all  $m \geq 0$ .  
 374 This is immediate as

$$375 \quad \|\mathcal{E}_{0,0}\|_{H^{s+3/2}} = 1, \quad \|\mathcal{E}_{0,m}\|_{H^{s+3/2}} = 0.$$

376 We now assume (4.10) for all  $n < \bar{n}$  and all  $m \geq 0$ , and seek this estimate in the case  
 377  $n = \bar{n}$  and all  $m \geq 0$ . For this we conduct another induction on  $m$ , and for  $m = 0$  we  
 378 use (4.11) (together with Lemma 4.3 with  $\tilde{s} = s + 1$ ) to discover

$$379 \quad \|\mathcal{E}_{\bar{n},0}\|_{H^{s+3/2}} \leq \mathcal{M} \left( \frac{|\underline{\gamma}^u| |f|_{C^{s+3/2+\eta}}}{\bar{n}} \right) \|\mathcal{E}_{\bar{n}-1,0}\|_{H^{s+3/2}} \\
 380 \quad \leq \mathcal{M} \left( \frac{|\underline{\gamma}^u| |f|_{C^{s+2}}}{\bar{n}} \right) C_{\mathcal{E}} B_{\mathcal{E}}^{\bar{n}-1} \leq C_{\mathcal{E}} B_{\mathcal{E}}^{\bar{n}}, \\
 381$$

382 provided that

$$383 \quad B_{\mathcal{E}} \geq \mathcal{M} |\underline{\gamma}^u| |f|_{C^{s+2}} \geq \mathcal{M} \left( \frac{|\underline{\gamma}^u| |f|_{C^{s+2}}}{\bar{n}} \right).$$

384 Finally, we assume the estimate (4.10) for  $n = \bar{n}$  and  $m < \bar{m}$ , and use (4.12) to learn  
 385 that

$$386 \quad \|\mathcal{E}_{\bar{n},\bar{m}}\|_{H^{s+3/2}} \leq \mathcal{M} \left( \frac{|\underline{\gamma}^u| |f|_{C^{s+3/2+\eta}}}{\bar{n}} \right) \{ \|\mathcal{E}_{\bar{n}-1,\bar{m}}\|_{H^{s+3/2}} + \|\mathcal{E}_{\bar{n}-1,\bar{m}-1}\|_{H^{s+3/2}} \} \\
 387 \quad \leq \mathcal{M} \left( \frac{|\underline{\gamma}^u| |f|_{C^{s+2}}}{\bar{n}} \right) C_{\mathcal{E}} \{ B_{\mathcal{E}}^{\bar{n}-1} D_{\mathcal{E}}^{\bar{m}} + B_{\mathcal{E}}^{\bar{n}-1} D_{\mathcal{E}}^{\bar{m}-1} \} \\
 388 \quad \leq C_{\mathcal{E}} B_{\mathcal{E}}^{\bar{n}} D_{\mathcal{E}}^{\bar{m}},$$

390 provided that

$$391 \quad \mathcal{M} \left( \frac{|\underline{\gamma}^u| |f|_{C^{s+2}}}{\bar{n}} \right) \leq \frac{B_{\mathcal{E}}}{2}, \quad \mathcal{M} \left( \frac{|\underline{\gamma}^u| |f|_{C^{s+2}}}{\bar{n}} \right) \leq \frac{B_{\mathcal{E}} D_{\mathcal{E}}}{2},$$

392 which can be accomplished, e.g., with

$$393 \quad B_{\mathcal{E}} \geq 2\mathcal{M} |\underline{\gamma}^u| |f|_{C^{s+2}} \geq 2\mathcal{M} \left( \frac{|\underline{\gamma}^u| |f|_{C^{s+2}}}{\bar{n}} \right), \quad D_{\mathcal{E}} \geq 1,$$

394 and we are done.  $\square$

395 With Lemma 4.7 it is straightforward to prove the following analyticity result for  
 396 the Dirichlet and Neumann data.

397 LEMMA 4.8. *Given any integer  $s \geq 0$ , if  $f \in C^{s+2}([0, d])$  then the functions  
 398  $\zeta(x; \varepsilon, \delta)$  and  $\psi(x; \varepsilon, \delta)$  are jointly analytic in  $\varepsilon$  and  $\delta$ . Therefore*

$$399 \quad (4.13) \quad \{\zeta, \psi\}(x; \varepsilon, \delta) = \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \{\zeta_{n,m}, \psi_{n,m}\}(x) \varepsilon^n \delta^m$$

400 and, for constants  $C_{\zeta}, B_{\zeta}, D_{\zeta} > 0$ , and  $C_{\psi}, B_{\psi}, D_{\psi} > 0$ ,

$$401 \quad (4.14) \quad \|\zeta_{n,m}\|_{H^{s+3/2}} \leq C_{\zeta} B_{\zeta}^n D_{\zeta}^m, \quad \|\psi_{n,m}\|_{H^{s+3/2}} \leq C_{\psi} B_{\psi}^n D_{\psi}^m,$$

402 for all  $n, m \geq 0$ .

403 **4.4. Invertibility of the Flat–Interface Operator.** The final hypothesis to  
 404 be verified in order to invoke Theorem 4.1 is the existence and mapping properties  
 405 of the linearized (flat–interface) operator  $\mathbf{A}_{0,0}$ . In our previous work [36] we showed  
 406 that

$$407 \quad (4.15) \quad \mathbf{A}_{0,0} = \begin{pmatrix} I & -I \\ G_{0,0} & \tau^2 J_{0,0} \end{pmatrix},$$

408 where

$$409 \quad (4.16) \quad G_{0,0} = -i\gamma_D^u, \quad J_{0,0} = -i\gamma_D^w,$$

410 are order–one Fourier multipliers defined by

$$411 \quad (4.17) \quad G_{0,0}[U] = \sum_{p=-\infty}^{\infty} (-i\gamma_p^u) \hat{U}_p e^{i\tilde{p}x}, \quad J_{0,0}[W] = \sum_{p=-\infty}^{\infty} (-i\gamma_p^w) \hat{W}_p e^{i\tilde{p}x}.$$

412 **LEMMA 4.9.** *The linear operator  $A_{0,0}$  maps  $X^s$  to  $Y^s$ , is invertible, and its inverse*  
 413 *maps  $Y^s$  to  $X^s$ .*

414 *Proof.* We begin by defining the operator

$$415 \quad \Delta := G_{0,0} + \tau^2 J_{0,0} = (-i\gamma_D^u) + \tau^2(-i\gamma_D^w),$$

416 which has Fourier symbol

$$417 \quad \hat{\Delta}_p = (-i\gamma_p^u) + \tau^2(-i\gamma_p^w),$$

418 and noting that there exist positive constants  $C_G$ ,  $C_J$ , and  $C_\Delta$  such that

$$419 \quad |-i\gamma_p^u| \leq C_G \langle \tilde{p} \rangle, \quad |-i\gamma_p^w| \leq C_J \langle \tilde{p} \rangle, \quad |\hat{\Delta}_p| \leq C_\Delta \langle \tilde{p} \rangle.$$

420 Importantly, provided that  $n^u \neq n^w$ , it is not difficult to establish that  $\hat{\Delta}_p \neq 0$ .

421 Finally, one can also find a positive constant  $C_{\Delta^{-1}}$  such that

$$422 \quad \left| \frac{1}{\hat{\Delta}_p} \right| \leq C_{\Delta^{-1}} \langle \tilde{p} \rangle^{-1}.$$

423 With this it is a simple matter to realize that  $\Delta^{-1}$  exists and that

$$424 \quad \Delta : H^{s+3/2} \rightarrow H^{s+1/2}, \quad \Delta^{-1} : H^{s+1/2} \rightarrow H^{s+3/2}.$$

425 Next, we write generic elements of  $X^s$  and  $Y^s$  as

$$426 \quad \mathbf{V} = \begin{pmatrix} U \\ W \end{pmatrix} \in X^s, \quad \mathbf{R} = \begin{pmatrix} \zeta \\ -\psi \end{pmatrix} \in Y^s.$$

427 Using the definitions of the norms of  $X^s$  and  $Y^s$  we find that

$$\begin{aligned} 428 \quad \|\mathbf{A}_{0,0}\mathbf{V}\|_{Y^s}^2 &= \|U - W\|_{H^{s+3/2}}^2 + \|G_{0,0}U + \tau^2 J_{0,0}W\|_{H^{s+1/2}}^2 \\ 429 \quad &\leq \|U\|_{H^{s+3/2}}^2 + \|W\|_{H^{s+3/2}}^2 + C_G^2 \|U\|_{H^{s+3/2}}^2 + C_J^2 \tau^4 \|W\|_{H^{s+3/2}}^2 \\ 430 \quad &\leq \max\{1, C_G^2, \tau^4 C_J^2\} \left( \|U\|_{H^{s+3/2}}^2 + \|W\|_{H^{s+3/2}}^2 \right) \\ 431 \quad &= \max\{1, C_G^2, \tau^4 C_J^2\} \|\mathbf{V}\|_{X^s}^2, \end{aligned}$$

433 so that  $\mathbf{A}_{0,0}$  does indeed map  $X^s$  to  $Y^s$ . We define the operator

$$434 \quad \mathbf{B} := \Delta^{-1} \begin{pmatrix} \tau^2 J_{0,0} & I \\ -G_{0,0} & I \end{pmatrix},$$

435 and note that

$$436 \quad \mathbf{B}\mathbf{A}_{0,0} = \mathbf{A}_{0,0}\mathbf{B} = \begin{pmatrix} I & 0 \\ 0 & I \end{pmatrix},$$

437 so that the inverse of  $\mathbf{A}_{0,0}$  exists and  $\mathbf{A}_{0,0}^{-1} = \mathbf{B}$ . Furthermore, as above,

$$\begin{aligned} 438 \quad \|\mathbf{A}_{0,0}^{-1}\mathbf{R}\|_{X^s}^2 &= \|\Delta^{-1}(\tau^2 J_{0,0}\zeta - \psi)\|_{H^{s+3/2}}^2 + \|\Delta^{-1}(-G_{0,0}\zeta - \psi)\|_{H^{s+3/2}}^2 \\ 439 \quad &\leq C_{\Delta^{-1}}\tau^4 C_J^2 \|\zeta\|_{H^{s+3/2}}^2 + C_{\Delta^{-1}} \|\psi\|_{H^{s+1/2}}^2 \\ 440 \quad &\quad + C_{\Delta^{-1}} C_G^2 \|\zeta\|_{H^{s+3/2}}^2 + C_{\Delta^{-1}} \|\psi\|_{H^{s+1/2}}^2 \\ 441 \quad &\leq C_{\Delta^{-1}} \max\{1, \tau^4 C_J^2, C_G^2\} \left( \|\zeta\|_{H^{s+3/2}}^2 + \|\psi\|_{H^{s+1/2}}^2 \right) \\ 442 \quad &= C_{\Delta^{-1}} \max\{1, \tau^4 C_J^2, C_G^2, \} \|\mathbf{R}\|_{Y^s}, \end{aligned}$$

444 and  $\mathbf{A}_{0,0}^{-1}$  maps  $Y^s$  to  $X^s$ . □

445 **5. Analyticity of the Scattered Fields.** At this point we establish the analy-  
446 ticity of the fields which define the DNOs,  $G$  and  $J$ , though, for brevity, we restrict  
447 our attention to the one in the upper layer,  $G$ , and note that the considerations for  
448 the lower layer DNO,  $J$ , are largely the same.

449 **5.1. Change of Variables and Formal Expansions.** For our rigorous demon-  
450 stration we appeal to the Method of Transformed Field Expansions (TFE) [53, 56]  
451 which begins with a domain-flattening change of variables (the  $\sigma$ -coordinates of  
452 oceanography [60] and the C-method of the dynamical theory of gratings [15, 14]) to  
453 the governing equations, (3.2),

$$454 \quad x' = x, \quad z' = a \left( \frac{z - g(x)}{a - g(x)} \right).$$

455 With this we can rewrite the DNO problem, (3.2), in terms of the transformed field

$$456 \quad u'(x', z') := u \left( x', \left( \frac{a - g(x')}{a} \right) z' + g(x') \right),$$

457 as (upon dropping primes)

$$458 \quad (5.1a) \quad \Delta u + 2i\alpha\partial_x u + (\gamma^u)^2 u = F(x, z), \quad 0 < z < a,$$

$$459 \quad (5.1b) \quad u(x, 0) = U(x), \quad z = 0,$$

$$460 \quad (5.1c) \quad \partial_z u(x, a) - T^u[u(x, a)] = P(x), \quad z = a,$$

$$461 \quad (5.1d) \quad u(x + d, z) = u(x, z),$$

463 and the DNO itself, (3.3), as

$$464 \quad (5.2) \quad G(g)[U] = -\partial_z u(x, 0) + H(x).$$

465 The forms for  $\{F, P, H\}$  have been derived and reported in [56] and, for brevity, we  
466 do not repeat them here.

467 Following our HOPS/AWE philosophy we assume the joint boundary/frequency  
468 perturbation

$$469 \quad g(x) = \varepsilon f(x), \quad \omega = \underline{\omega} + \delta \underline{\omega} = (1 + \delta) \underline{\omega},$$

470 and study the effect of this on (5.1) and (5.2). These become

$$471 \quad (5.3a) \quad \Delta u + 2i\underline{\alpha} \partial_x u + (\underline{\gamma}^u)^2 u = \tilde{F}(x, z), \quad 0 < z < a,$$

$$472 \quad (5.3b) \quad u(x, 0) = U(x), \quad z = 0,$$

$$473 \quad (5.3c) \quad \partial_z u(x, a) - T^u[u(x, a)] = \tilde{P}(x), \quad z = a,$$

$$474 \quad (5.3d) \quad u(x + d, z) = u(x, z),$$

476 and

$$477 \quad (5.4) \quad G(\varepsilon f)[U] = -\partial_z u(x, 0) + \tilde{H}(x),$$

478 where  $\tilde{F}, \tilde{P}, \tilde{H} = \mathcal{O}(\varepsilon) + \mathcal{O}(\delta)$ . More specifically,

$$479 \quad \begin{aligned} \tilde{F} = & -\varepsilon \operatorname{div} [A_1(f) \nabla u] - \varepsilon^2 \operatorname{div} [A_2(f) \nabla u] - \varepsilon B_1(f) \nabla u - \varepsilon^2 B_2(f) \nabla u \\ 480 & - 2i\underline{\alpha} \delta \partial_x u - \delta^2 (\underline{\gamma}^u)^2 u - 2\delta (\underline{\gamma}^u)^2 u \\ 481 & - 2i\varepsilon S_1(f) \underline{\alpha} \partial_x u - 2i\varepsilon S_1(f) \underline{\alpha} \delta \partial_x u - \varepsilon S_1(f) \delta^2 (\underline{\gamma}^u)^2 u \\ 482 & - 2\varepsilon S_1(f) \delta (\underline{\gamma}^u)^2 u - \varepsilon S_1(f) (\underline{\gamma}^u)^2 u \\ 483 & - 2i\varepsilon^2 S_2(f) \underline{\alpha} \partial_x u - 2i\varepsilon^2 S_2(f) \underline{\alpha} \delta \partial_x u - \varepsilon^2 S_2(f) \delta^2 (\underline{\gamma}^u)^2 u \\ 484 \quad (5.5) & - 2\varepsilon^2 S_2(f) \delta (\underline{\gamma}^u)^2 u - \varepsilon^2 S_2(f) (\underline{\gamma}^u)^2 u, \end{aligned}$$

486 and

$$487 \quad (5.6) \quad \tilde{P} = -\frac{1}{a} (\varepsilon f(x)) T^u [u(x, a)],$$

488 and

$$489 \quad (5.7) \quad \tilde{H} = \varepsilon (\partial_x f) \partial_x u(x, 0) + \varepsilon \frac{f}{a} G(\varepsilon f)[U] - \varepsilon^2 \frac{f(\partial_x f)}{a} \partial_x u(x, 0) - \varepsilon^2 (\partial_x f)^2 \partial_z u(x, 0).$$

490 It is not difficult to see that the forms for the  $A_j$ ,  $B_j$ , and  $S_j$  are

$$491 \quad (5.8a) \quad A_0 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix},$$

$$492 \quad (5.8b) \quad A_1(f) = \begin{pmatrix} A_1^{xx} & A_1^{xz} \\ A_1^{zx} & A_1^{zz} \end{pmatrix} = \frac{1}{a} \begin{pmatrix} -2f & -(a-z)(\partial_x f) \\ -(a-z)(\partial_x f) & 0 \end{pmatrix},$$

$$493 \quad (5.8c) \quad A_2(f) = \begin{pmatrix} A_2^{xx} & A_2^{xz} \\ A_2^{zx} & A_2^{zz} \end{pmatrix} = \frac{1}{a^2} \begin{pmatrix} f^2 & (a-z)f(\partial_x f) \\ (a-z)f(\partial_x f) & (a-z)^2(\partial_x f)^2 \end{pmatrix},$$

495 and

$$496 \quad (5.9) \quad B_1(f) = \begin{pmatrix} B_1^x \\ B_1^z \end{pmatrix} = \frac{1}{a} \begin{pmatrix} \partial_x f \\ 0 \end{pmatrix}, \quad B_2(f) = \begin{pmatrix} B_2^x \\ B_2^z \end{pmatrix} = \frac{1}{a^2} \begin{pmatrix} -f(\partial_x f) \\ -(a-z)(\partial_x f)^2 \end{pmatrix},$$

497 and

$$498 \quad (5.10) \quad S_0 = 1, \quad S_1(f) = -\frac{2}{a} f, \quad S_2(f) = \frac{1}{a^2} f^2.$$

499 At this point we posit the expansions

$$500 \quad u(x, z; \varepsilon, \delta) = \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} u_{n,m}(x, z) \varepsilon^n \delta^m, \quad G(\varepsilon, \delta) = \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} G_{n,m} \varepsilon^n \delta^m,$$

501 and, upon insertion into (5.3) and (5.4), we find

$$502 \quad (5.11a) \quad \Delta u_{n,m} + 2i\alpha \partial_x u_{n,m} + (\underline{\gamma}^u)^2 u_{n,m} = \tilde{F}_{n,m}(x, z), \quad 0 < z < a,$$

$$503 \quad (5.11b) \quad u_{n,m}(x, 0) = U_{n,m}(x), \quad z = 0,$$

$$504 \quad (5.11c) \quad \partial_z u_{n,m}(x, a) - T^u[u_{n,m}(x, a)] = \tilde{P}_{n,m}(x), \quad z = a,$$

$$505 \quad (5.11d) \quad u_{n,m}(x+d, z) = u_{n,m}(x, z),$$

507 and

$$508 \quad (5.12) \quad G_{n,m}(f) = -\partial_z u_{n,m}(x, 0) + \tilde{H}_{n,m}(x).$$

509 The formulas for  $\tilde{F}_{n,m}$ ,  $\tilde{P}_{n,m}$  and  $\tilde{H}_{n,m}$  can be readily derived from (5.5), (5.6), and  
510 (5.7) giving

$$\begin{aligned} 511 \quad \tilde{F}_{n,m} &= -\operatorname{div} [A_1(f) \nabla u_{n-1,m}] - \operatorname{div} [A_2(f) \nabla u_{n-2,m}] \\ 512 &\quad - B_1(f) \nabla u_{n-1,m} - B_2(f) \nabla u_{n-2,m} \\ 513 &\quad - 2i\alpha \partial_x u_{n,m-1} - (\underline{\gamma}^u)^2 u_{n,m-2} - 2(\underline{\gamma}^u)^2 u_{n,m-1} \\ 514 &\quad - 2iS_1(f) \alpha \partial_x u_{n-1,m} - 2iS_1(f) \alpha \partial_x u_{n-1,m-1} - S_1(f) (\underline{\gamma}^u)^2 u_{n-1,m-2} \\ 515 &\quad - 2S_1(f) (\underline{\gamma}^u)^2 u_{n-1,m-1} - S_1(f) (\underline{\gamma}^u)^2 u_{n-1,m} \\ 516 &\quad - 2iS_2(f) \alpha \partial_x u_{n-2,m} - 2iS_2(f) \alpha \partial_x u_{n-2,m-1} - S_2(f) (\underline{\gamma}^u)^2 u_{n-2,m-2} \\ 517 \quad (5.13) &\quad - 2S_2(f) (\underline{\gamma}^u)^2 u_{n-2,m-1} - S_2(f) (\underline{\gamma}^u)^2 u_{n-2,m}, \end{aligned}$$

519 and

$$520 \quad (5.14) \quad \tilde{P}_{n,m} = -\frac{1}{a} f(x) T^u [u_{n-1,m}(x, a)],$$

521 and

$$\begin{aligned} 522 \quad \tilde{H}_{n,m} &= (\partial_x f) \partial_x u_{n-1,m}(x, 0) + \frac{f}{a} G_{n-1,m}(f) [U] - \frac{f(\partial_x f)}{a} \partial_x u_{n-2,m}(x, 0) \\ 523 \quad (5.15) &\quad - (\partial_x f)^2 \partial_z u_{n-2,m}(x, 0). \end{aligned}$$

525 **5.2. Geometric Analyticity of the Upper Field.** To prove our joint analyt-  
526 icity result we begin by stating the single, geometric, analyticity result for the field  
527  $u$  under boundary perturbation,  $\varepsilon$ , alone. This was essentially established in [53] but  
528 we present it here for completeness.

529 **THEOREM 5.1.** *Given any integer  $s \geq 0$ , if  $f \in C^{s+2}([0, d])$  and  $U_{n,0} \in H^{s+3/2}([0, d])$*   
530 *such that*

$$531 \quad \|U_{n,0}\|_{H^{s+3/2}} \leq K_U B_U^n$$

532 *for constants  $K_U, B_U > 0$ , then  $u_{n,0} \in H^{s+2}([0, d] \times [0, a])$  and*

$$533 \quad (5.16) \quad \|u_{n,0}\|_{H^{s+2}} \leq K B^n,$$

534 *for constants  $K, B > 0$ .*



535 To establish this we work by induction and the key estimate is the following Lemma.

536 LEMMA 5.2. *Given an integer  $s \geq 0$ , if  $f \in C^{s+2}([0, d])$  and*

$$537 \quad (5.17) \quad \|u_{n,0}\|_{H^{s+2}} \leq KB^n, \quad \forall n < \bar{n},$$

538 *for constants  $K, B > 0$  then there exists a constant  $\bar{C} > 0$  such that*

$$539 \quad (5.18) \quad \max \left\{ \left\| \tilde{F}_{\bar{n},0} \right\|_{H^s}, \left\| \tilde{P}_{\bar{n},0} \right\|_{H^{s+1/2}} \right\} \leq K\bar{C} \left\{ |f|_{C^{s+2}} B^{\bar{n}-1} + |f|_{C^{s+2}}^2 B^{\bar{n}-2} \right\}.$$

540 *Proof.* [Lemma 5.2] We begin with  $\tilde{F}_{\bar{n},0}$  and note that from (5.13), (5.8), (5.9),  
541 and (5.10) we have

$$\begin{aligned} 542 \quad \|\tilde{F}_{\bar{n},0}\|_{H^s}^2 &\leq \|A_1^{xx} \partial_x u_{\bar{n}-1,0}\|_{H^{s+1}}^2 + \|A_1^{xz} \partial_z u_{\bar{n}-1,0}\|_{H^{s+1}}^2 + \|A_1^{zx} \partial_x u_{\bar{n}-1,0}\|_{H^{s+1}}^2 \\ 543 &\quad + \|A_1^{zz} \partial_z u_{\bar{n}-1,0}\|_{H^{s+1}}^2 + \|A_2^{xx} \partial_x u_{\bar{n}-2,0}\|_{H^{s+1}}^2 + \|A_2^{xz} \partial_z u_{\bar{n}-2,0}\|_{H^{s+1}}^2 \\ 544 &\quad + \|A_2^{zx} \partial_x u_{\bar{n}-2,0}\|_{H^{s+1}}^2 + \|A_2^{zz} \partial_z u_{\bar{n}-2,0}\|_{H^{s+1}}^2 + \|B_1^x \partial_x u_{\bar{n}-1,0}\|_{H^s}^2 \\ 545 &\quad + \|B_1^z \partial_z u_{\bar{n}-1,0}\|_{H^s}^2 + \|B_2^x \partial_x u_{\bar{n}-2,0}\|_{H^s}^2 + \|B_2^z \partial_z u_{\bar{n}-2,0}\|_{H^s}^2 \\ 546 &\quad + \|2S_1 i\alpha \partial_x u_{\bar{n}-1,0}\|_{H^s}^2 + \|S_1 (\underline{\gamma}^u)^2 u_{\bar{n}-1,0}\|_{H^s}^2 + \|2S_2 i\alpha \partial_x u_{\bar{n}-2,0}\|_{H^s}^2 \\ 547 &\quad + \|S_2 (\underline{\gamma}^u)^2 u_{\bar{n}-2,0}\|_{H^s}^2. \end{aligned}$$

549 We now estimate each of these by applying Lemmas 4.3 and 4.5. We begin with

$$\begin{aligned} 550 \quad \|A_1^{xx} \partial_x u_{\bar{n}-1,0}\|_{H^{s+1}} &= \|-(2/a) f \partial_x u_{\bar{n}-1,0}\|_{H^{s+1}} \\ 551 &\leq (2/a) \mathcal{M} |f|_{C^{s+1}} \|u_{\bar{n}-1,0}\|_{H^{s+2}} \\ 552 &\leq (2/a) \mathcal{M} |f|_{C^{s+1}} KB^{\bar{n}-1}, \end{aligned}$$

554 and in a similar fashion

$$\begin{aligned} 555 \quad \|A_1^{xz} \partial_z u_{\bar{n}-1,0}\|_{H^{s+1}} &= \| -((a-z)/a) (\partial_x f) \partial_z u_{\bar{n}-1,0} \|_{H^{s+1}} \\ 556 &\leq (Z_a/a) \mathcal{M} |\partial_x f|_{C^{s+1}} \|u_{\bar{n}-1,0}\|_{H^{s+2}} \\ 557 &\leq (Z_a/a) \mathcal{M} |f|_{C^{s+2}} KB^{\bar{n}-1}. \end{aligned}$$

559 Also,

$$\begin{aligned} 560 \quad \|A_1^{zx} \partial_x u_{\bar{n}-1,0}\|_{H^{s+1}} &= \| -((a-z)/a) (\partial_x f) \partial_x u_{\bar{n}-1,0} \|_{H^{s+1}} \\ 561 &\leq (Z_a/a) \mathcal{M} |\partial_x f|_{C^{s+1}} \|u_{\bar{n}-1,0}\|_{H^{s+2}} \\ 562 &\leq (Z_a/a) \mathcal{M} |f|_{C^{s+2}} KB^{\bar{n}-1}, \end{aligned}$$

564 and we recall that  $A_1^{zz} \equiv 0$ . Moving to the second order

$$\begin{aligned} 565 \quad \|A_2^{xx} \partial_x u_{\bar{n}-2,0}\|_{H^{s+1}} &= \|(1/a^2) f^2 \partial_x u_{\bar{n}-2,0}\|_{H^{s+1}} \\ 566 &\leq (1/a^2) \mathcal{M}^2 |f|_{C^{s+1}}^2 \|u_{\bar{n}-2,0}\|_{H^{s+2}} \\ 567 &\leq (1/a^2) \mathcal{M}^2 |f|_{C^{s+1}}^2 KB^{\bar{n}-2}. \end{aligned}$$

569 Also,

$$\begin{aligned} 570 \quad \|A_2^{zz} \partial_z u_{\bar{n}-2,0}\|_{H^{s+1}} &= \|((a-z)/a^2) f (\partial_x f) \partial_x u_{\bar{n}-2,0}\|_{H^{s+1}} \\ 571 &\leq (Z_a/a^2) \mathcal{M}^2 |f|_{C^{s+1}} |\partial_x f|_{C^{s+1}} \|u_{\bar{n}-2,0}\|_{H^{s+2}} \\ 572 &\leq (Z_a/a^2) \mathcal{M}^2 |f|_{C^{s+2}}^2 KB^{\bar{n}-2}, \end{aligned}$$

574 and

$$\begin{aligned}
575 \quad \|A_2^{zx} \partial_x u_{\bar{n}-2,0}\|_{H^{s+1}} &= \|((a-z)/a^2) f(\partial_x f) \partial_z u_{\bar{n}-2,0}\|_{H^{s+1}} \\
576 \quad &\leq (Z_a/a^2) \mathcal{M}^2 |f|_{C^{s+1}} |\partial_x f|_{C^{s+1}} \|u_{\bar{n}-2,0}\|_{H^{s+2}} \\
577 \quad &\leq (Z_a/a^2) \mathcal{M}^2 |f|_{C^{s+2}}^2 K B^{\bar{n}-2},
\end{aligned}$$

579 and

$$\begin{aligned}
580 \quad \|A_2^{zz} \partial_z u_{\bar{n}-2,0}\|_{H^{s+1}} &= \|((a-z)^2/a^2) (\partial_x f)^2 \partial_z u_{\bar{n}-2,0}\|_{H^{s+1}} \\
581 \quad &\leq (Z_a^2/a^2) \mathcal{M}^2 |\partial_x f|_{C^{s+1}}^2 \|u_{\bar{n}-2,0}\|_{H^{s+2}} \\
582 \quad &\leq (Z_a^2/a^2) \mathcal{M}^2 |f|_{C^{s+2}}^2 K B^{\bar{n}-2}.
\end{aligned}$$

584 Next for the  $B_1$  terms

$$\begin{aligned}
585 \quad \|B_1^x \partial_x u_{\bar{n}-1,0}\|_{H^s} &= \|(1/a) (\partial_x f) \partial_x u_{\bar{n}-1,0}\|_{H^s} \\
586 \quad &\leq (1/a) \mathcal{M} |\partial_x f|_{C^{s+1}} \|u_{\bar{n}-1,0}\|_{H^s} \\
587 \quad &\leq (1/a) \mathcal{M} |f|_{C^{s+2}} K B^{\bar{n}-1},
\end{aligned}$$

589 and  $B_1^z \equiv 0$ . Moving to the second order

$$\begin{aligned}
590 \quad \|B_2^x \partial_x u_{\bar{n}-2,0}\|_{H^s} &= \|(-1/a^2) f(\partial_x f) \partial_x u_{\bar{n}-2,0}\|_{H^s} \\
591 \quad &\leq (1/a^2) \mathcal{M}^2 |f|_{C^{s+1}} |\partial_x f|_{C^{s+1}} \|u_{\bar{n}-2,0}\|_{H^s} \\
592 \quad &\leq (1/a^2) \mathcal{M}^2 |f|_{C^{s+2}}^2 K B^{\bar{n}-2},
\end{aligned}$$

594 and

$$\begin{aligned}
595 \quad \|B_2^z \partial_z u_{\bar{n}-2,0}\|_{H^s} &= \|(-1/a^2) (a-z) (\partial_x f)^2 \partial_z u_{\bar{n}-2,0}\|_{H^s} \\
596 \quad &\leq (Z_a/a^2) \mathcal{M}^2 |\partial_x f|_{C^{s+1}} \|u_{\bar{n}-2,0}\|_{H^s} \\
597 \quad &\leq (Z_a/a^2) \mathcal{M}^2 |f|_{C^{s+2}}^2 K B^{\bar{n}-2}.
\end{aligned}$$

599 To address the  $S_0, S_1, S_2$  terms we have

$$\begin{aligned}
600 \quad \|2S_1 i \underline{\alpha} \partial_x u_{\bar{n}-1,0}\|_{H^s} &= \|(-4/a) i \underline{\alpha} f \partial_x u_{\bar{n}-1,0}\|_{H^s} \\
601 \quad &\leq (4/a) \underline{\alpha} \mathcal{M} |f|_{C^s} \|u_{\bar{n}-1,0}\|_{H^{s+1}} \\
602 \quad &\leq (4/a) \underline{\alpha} \mathcal{M} |f|_{C^s} K B^{\bar{n}-1},
\end{aligned}$$

604 and

$$\begin{aligned}
605 \quad \|S_1 (\underline{\gamma}^u)^2 u_{\bar{n}-1,0}\|_{H^s} &= \|(-2/a) (\underline{\gamma}^u)^2 f u_{\bar{n}-1,0}\|_{H^s} \\
606 \quad &\leq (2/a) (\underline{\gamma}^u)^2 \mathcal{M} |f|_{C^s} \|u_{\bar{n}-1,0}\|_{H^s} \\
607 \quad &\leq (2/a) (\underline{\gamma}^u)^2 \mathcal{M} |f|_{C^s} K B^{\bar{n}-1},
\end{aligned}$$

609 and

$$\begin{aligned}
610 \quad \|2S_2 i \underline{\alpha} \partial_x u_{\bar{n}-2,0}\|_{H^s} &= \|(2/a^2) i \underline{\alpha} f^2 \partial_x u_{\bar{n}-2,0}\|_{H^s} \\
611 \quad &\leq (2/a^2) \underline{\alpha} \mathcal{M}^2 |f|_{C^s}^2 \|u_{\bar{n}-2,0}\|_{H^{s+1}} \\
612 \quad &\leq (2/a^2) \underline{\alpha} \mathcal{M}^2 |f|_{C^s}^2 K B^{\bar{n}-2},
\end{aligned}$$

614 and

$$\begin{aligned}
615 \quad \|S_2(\underline{\gamma}^u)^2 u_{\bar{n}-2,0}\|_{H^s} &= \|(1/a^2)(\underline{\gamma}^u)^2 f^2 u_{\bar{n}-2,0}\|_{H^s} \\
616 \quad &\leq (1/a^2)(\underline{\gamma}^u)^2 \mathcal{M}^2 |f|_{C^s}^2 \|u_{\bar{n}-2,0}\|_{H^s} \\
617 \quad &\leq (1/a^2)(\underline{\gamma}^u)^2 \mathcal{M}^2 |f|_{C^s}^2 K B^{\bar{n}-2}.
\end{aligned}$$

619 We satisfy the estimate for  $\|\tilde{F}_{\bar{n},0}\|_{H^s}$  provided that we choose

$$620 \quad \bar{C} > \max \left\{ \left( \frac{3 + 2Z_a + 4\underline{\alpha} + 2(\underline{\gamma}^u)^2}{a} \right) \mathcal{M}, \left( \frac{2 + 3Z_a + Z_a^2 + 2\underline{\alpha} + (\underline{\gamma}^u)^2}{a^2} \right) \mathcal{M}^2 \right\}.$$

621 The estimate for  $\tilde{P}_{\bar{n},0}$  follows from an elementary estimate on the order-one Fourier  
622 multiplier  $T^u$

$$\begin{aligned}
623 \quad \|\tilde{P}_{\bar{n},0}\|_{H^{s+1/2}} &= \|(1/a) f T^u [u_{\bar{n}-1,0}]\|_{H^{s+1/2}} \\
624 \quad &\leq (1/a) \mathcal{M} |f|_{C^{s+1/2+\eta}} \|T^u [u_{\bar{n}-1,0}]\|_{H^{s+1/2}} \\
625 \quad &\leq (1/a) \mathcal{M} |f|_{C^{s+1/2+\eta}} C_{T^u} \|u_{\bar{n}-1,0}\|_{H^{s+3/2}} \\
626 \quad &\leq (1/a) \mathcal{M} |f|_{C^{s+1/2+\eta}} C_{T^u} K B^{\bar{n}-1},
\end{aligned}$$

628 and provided that

$$629 \quad \bar{C} > (1/a) \mathcal{M} C_{T^u},$$

630 we are done.  $\square$

631 With this information, we can now prove Theorem 5.1.

632 *Proof.* [Theorem 5.1] We proceed by induction and at order  $n = 0$  and  $m = 0$   
633 Theorem 4.4 guarantees a unique solution such that

$$634 \quad \|u_{0,0}\|_{H^{s+2}} \leq C_e \|U_{0,0}\|_{H^{s+3/2}}.$$

635 So we choose  $K \geq C_e \|U_{0,0}\|_{H^{s+3/2}}$ . We now assume the estimate (5.16) for all  $n < \bar{n}$   
636 and study  $u_{\bar{n},0}$ . From Theorem 4.4 we have a unique solution satisfying

$$637 \quad \|u_{\bar{n},0}\|_{H^{s+2}} \leq C_e \{ \|\tilde{F}_{\bar{n},0}\|_{H^s} + \|U_{\bar{n},0}\|_{H^{s+3/2}} + \|\tilde{P}_{\bar{n},0}\|_{H^{s+1/2}} \},$$

638 and appealing to Lemmas 4.8 and 5.2 we find

$$639 \quad \|u_{\bar{n},0}\|_{H^{s+2}} \leq C_e \{ K_U B_U^{\bar{n}} + 2K\bar{C} [|f|_{C^{s+2}} B^{\bar{n}-1} + |f|_{C^{s+2}}^2 B^{\bar{n}-2}] \}.$$

640 We are done provided we choose  $K \geq 3C_e K_U$  and

$$641 \quad B > \max \left\{ B_\zeta, 6C_e \bar{C} |f|_{C^{s+2}}, \sqrt{6C_e \bar{C}} |f|_{C^{s+2}} \right\}. \quad \square$$

643 Analogous results hold in the lower field which we record here for completeness.

644 **THEOREM 5.3.** *Given any integer  $s \geq 0$ , if  $f \in C^{s+2}([0, d])$  and  $W_{n,0} \in H^{s+3/2}([0, d])$*   
645 *such that*

$$646 \quad \|W_{n,0}\|_{H^{s+3/2}} \leq K_W B_W^n$$

647 for constants  $K_W, B_W > 0$ , then  $w_{n,0} \in H^{s+2}([0, d] \times [-b, 0])$  and

$$648 \quad \|w_{n,0}\|_{H^{s+2}} \leq KB^n,$$

649 for constants  $K, B > 0$ .

650 **5.3. Joint Analyticity of the Upper Field.** We can now proceed to prove  
651 our main result concerning joint analyticity of the transformed field.

652 **THEOREM 5.4.** *Given any integer  $s \geq 0$ , if  $f \in C^{s+2}([0, d])$  and  $U_{n,m} \in H^{s+3/2}([0, d])$*   
653 *such that*

$$654 \quad \|U_{n,m}\|_{H^{s+3/2}} \leq K_U B_U^n D_U^m,$$

655 for constants  $K_U, B_U, D_U > 0$ , then  $u_{n,m} \in H^{s+2}([0, d] \times [0, a])$  and

$$656 \quad (5.19) \quad \|u_{n,m}\|_{H^{s+2}} \leq KB^n D^m,$$

657 for constants  $K, B, D > 0$ .

658 As before, we establish this result by induction.

659 **LEMMA 5.5.** *Given an integer  $s \geq 0$ , if  $f \in C^{s+2}([0, d])$  and*

$$660 \quad (5.20) \quad \|u_{n,m}\|_{H^{s+2}} \leq KB^n D^m, \quad \forall n \geq 0, m < \bar{m},$$

661 for constants  $K, B, D > 0$  then there exists a constant  $\bar{C} > 0$  such that

$$662 \quad \max\{\|\tilde{F}_{n,\bar{m}}\|_{H^s}, \|\tilde{P}_{n,\bar{m}}\|_{H^{s+1/2}}\} \leq K\bar{C} \left\{ \underline{\alpha}(\underline{\gamma}^u)^2 B^n D^{\bar{m}-1} + (\underline{\gamma}^u)^2 B^n D^{\bar{m}-2} \right. \\ 663 \quad \quad \quad + \underline{\alpha}(\underline{\gamma}^u)^2 |f|_{C^{s+2}} B^{n-1} D^{\bar{m}} + \underline{\alpha}(\underline{\gamma}^u)^2 |f|_{C^{s+2}} B^{n-1} D^{\bar{m}-1} \\ 664 \quad \quad \quad + (\underline{\gamma}^u)^2 |f|_{C^{s+2}} B^{n-1} D^{\bar{m}-2} + \underline{\alpha}(\underline{\gamma}^u)^2 |f|_{C^{s+2}}^2 B^{n-2} D^{\bar{m}} \\ 665 \quad \quad \quad \left. + \underline{\alpha}(\underline{\gamma}^u)^2 |f|_{C^{s+2}}^2 B^{n-2} D^{\bar{m}-1} + (\underline{\gamma}^u)^2 |f|_{C^{s+2}}^2 B^{n-2} D^{\bar{m}-2} \right\}.$$

667 *Proof.* [Lemma 5.5] We begin with  $\tilde{F}_{n,\bar{m}}$  and note that from (5.13), (5.8), (5.9),  
668 and (5.10) we have

$$669 \quad \|\tilde{F}_{n,\bar{m}}\|_{H^s}^2 \leq \|A_1^{xx} \partial_x u_{n-1,\bar{m}}\|_{H^{s+1}}^2 + \|A_1^{xz} \partial_z u_{n-1,\bar{m}}\|_{H^{s+1}}^2 + \|A_1^{zx} \partial_x u_{n-1,\bar{m}}\|_{H^{s+1}}^2 \\ 670 \quad \quad \quad + \|A_1^{zz} \partial_z u_{n-1,\bar{m}}\|_{H^{s+1}}^2 + \|A_2^{xx} \partial_x u_{n-2,\bar{m}}\|_{H^{s+1}}^2 + \|A_2^{xz} \partial_z u_{n-2,\bar{m}}\|_{H^{s+1}}^2 \\ 671 \quad \quad \quad + \|A_2^{zx} \partial_x u_{n-2,\bar{m}}\|_{H^{s+1}}^2 + \|A_2^{zz} \partial_z u_{n-2,\bar{m}}\|_{H^{s+1}}^2 + \|B_1^x \partial_x u_{n-1,\bar{m}}\|_{H^s}^2 \\ 672 \quad \quad \quad + \|B_1^z \partial_z u_{n-1,\bar{m}}\|_{H^s}^2 + \|B_2^x \partial_x u_{n-2,\bar{m}}\|_{H^s}^2 + \|B_2^z \partial_z u_{n-2,\bar{m}}\|_{H^s}^2 \\ 673 \quad \quad \quad + \|2i\underline{\alpha} \partial_x u_{n,\bar{m}-1}\|_{H^s}^2 + \|(\underline{\gamma}^u)^2 u_{n,\bar{m}-2}\|_{H^s}^2 + \|2(\underline{\gamma}^u)^2 u_{n,\bar{m}-1}\|_{H^s}^2 \\ 674 \quad \quad \quad + \|2S_1 i\underline{\alpha} \partial_x u_{n-1,\bar{m}}\|_{H^s}^2 + \|2S_1 i\underline{\alpha} \partial_x u_{n-1,\bar{m}-1}\|_{H^s}^2 + \|S_1 (\underline{\gamma}^u)^2 u_{n-1,\bar{m}-2}\|_{H^s}^2 \\ 675 \quad \quad \quad + \|2S_1 (\underline{\gamma}^u)^2 u_{n-1,\bar{m}-1}\|_{H^s}^2 + \|S_1 (\underline{\gamma}^u)^2 u_{n-1,\bar{m}}\|_{H^s}^2 + \|2S_2 i\underline{\alpha} \partial_x u_{n-2,\bar{m}}\|_{H^s}^2 \\ 676 \quad \quad \quad + \|2S_2 i\underline{\alpha} \partial_x u_{n-2,\bar{m}-1}\|_{H^s}^2 + \|S_2 (\underline{\gamma}^u)^2 u_{n-2,\bar{m}-2}\|_{H^s}^2 \\ 677 \quad \quad \quad + \|2S_2 (\underline{\gamma}^u)^2 u_{n-2,\bar{m}-1}\|_{H^s}^2 + \|S_2 (\underline{\gamma}^u)^2 u_{n-2,\bar{m}}\|_{H^s}^2.$$

679 We now estimate each of these by applying Lemmas 4.3 and 4.5. We begin with

$$680 \quad \|A_1^{xx} \partial_x u_{n-1,\bar{m}}\|_{H^{s+1}} = \|-(2/a) f \partial_x u_{n-1,\bar{m}}\|_{H^{s+1}} \\ 681 \quad \quad \quad \leq (2/a) \mathcal{M} |f|_{C^{s+1}} \|u_{n-1,\bar{m}}\|_{H^{s+2}} \\ 682 \quad \quad \quad \leq (2/a) \mathcal{M} |f|_{C^{s+1}} KB^{n-1} D^{\bar{m}},$$

684 and in a similar fashion

$$\begin{aligned}
685 \quad \|A_1^{xz} \partial_z u_{n-1, \bar{m}}\|_{H^{s+1}} &= \| -((a-z)/a)(\partial_x f) \partial_z u_{n-1, \bar{m}} \|_{H^{s+1}} \\
686 \quad &\leq (Z_a/a) \mathcal{M} |\partial_x f|_{C^{s+1}} \|u_{n-1, \bar{m}}\|_{H^{s+2}} \\
687 \quad &\leq (Z_a/a) \mathcal{M} |f|_{C^{s+2}} K B^{n-1} D^{\bar{m}}.
\end{aligned}$$

689 Also,

$$\begin{aligned}
690 \quad \|A_1^{zx} \partial_x u_{n-1, \bar{m}}\|_{H^{s+1}} &= \| -((a-z)/a)(\partial_x f) \partial_x u_{n-1, \bar{m}} \|_{H^{s+1}} \\
691 \quad &\leq (Z_a/a) \mathcal{M} |\partial_x f|_{C^{s+1}} \|u_{n-1, \bar{m}}\|_{H^{s+2}} \\
692 \quad &\leq (Z_a/a) \mathcal{M} |f|_{C^{s+2}} K B^{n-1} D^{\bar{m}},
\end{aligned}$$

694 and we recall that  $A_1^{zz} \equiv 0$ . Moving to the second order

$$\begin{aligned}
695 \quad \|A_2^{xx} \partial_x u_{n-2, \bar{m}}\|_{H^{s+1}} &= \|(1/a^2) f^2 \partial_x u_{n-2, \bar{m}}\|_{H^{s+1}} \\
696 \quad &\leq (1/a^2) \mathcal{M}^2 |f|_{C^{s+1}}^2 \|u_{n-2, \bar{m}}\|_{H^{s+2}} \\
697 \quad &\leq (1/a^2) \mathcal{M}^2 |f|_{C^{s+1}}^2 K B^{n-2} D^{\bar{m}}.
\end{aligned}$$

699 Also,

$$\begin{aligned}
700 \quad \|A_2^{xz} \partial_z u_{n-2, \bar{m}}\|_{H^{s+1}} &= \|((a-z)/a^2) f (\partial_x f) \partial_x u_{n-2, \bar{m}}\|_{H^{s+1}} \\
701 \quad &\leq (Z_a/a^2) \mathcal{M}^2 |f|_{C^{s+1}} |\partial_x f|_{C^{s+1}} \|u_{n-2, \bar{m}}\|_{H^{s+2}} \\
702 \quad &\leq (Z_a/a^2) \mathcal{M}^2 |f|_{C^{s+2}}^2 K B^{n-2} D^{\bar{m}},
\end{aligned}$$

704 and

$$\begin{aligned}
705 \quad \|A_2^{zx} \partial_x u_{n-2, \bar{m}}\|_{H^{s+1}} &= \|((a-z)/a^2) f (\partial_x f) \partial_z u_{n-2, \bar{m}}\|_{H^{s+1}} \\
706 \quad &\leq (Z_a/a^2) \mathcal{M}^2 |f|_{C^{s+1}} |\partial_x f|_{C^{s+1}} \|u_{n-2, \bar{m}}\|_{H^{s+2}} \\
707 \quad &\leq (Z_a/a^2) \mathcal{M}^2 |f|_{C^{s+2}}^2 K B^{n-2} D^{\bar{m}},
\end{aligned}$$

709 and

$$\begin{aligned}
710 \quad \|A_2^{zz} \partial_z u_{n-2, \bar{m}}\|_{H^{s+1}} &= \|((a-z)^2/a^2) (\partial_x f)^2 \partial_z u_{n-2, \bar{m}}\|_{H^{s+1}} \\
711 \quad &\leq (Z_a^2/a^2) \mathcal{M}^2 |\partial_x f|_{C^{s+1}}^2 \|u_{n-2, \bar{m}}\|_{H^{s+2}} \\
712 \quad &\leq (Z_a^2/a^2) \mathcal{M}^2 |f|_{C^{s+2}}^2 K B^{n-2} D^{\bar{m}}.
\end{aligned}$$

714 Next for the  $B_1$  terms

$$\begin{aligned}
715 \quad \|B_1^x \partial_x u_{n-1, \bar{m}}\|_{H^s} &= \|(1/a) (\partial_x f) \partial_x u_{n-1, \bar{m}}\|_{H^s} \\
716 \quad &\leq (1/a) \mathcal{M} |\partial_x f|_{C^{s+1}} \|u_{n-1, \bar{m}}\|_{H^s} \\
717 \quad &\leq (1/a) \mathcal{M} |f|_{C^{s+2}} K B^{n-1} D^{\bar{m}},
\end{aligned}$$

719 and  $B_1^z \equiv 0$ . Moving to the second order

$$\begin{aligned}
720 \quad \|B_2^x \partial_x u_{n-2, \bar{m}}\|_{H^s} &= \|(-1/a^2) f (\partial_x f) \partial_x u_{n-2, \bar{m}}\|_{H^s} \\
721 \quad &\leq (1/a^2) \mathcal{M}^2 |f|_{C^{s+1}} |\partial_x f|_{C^{s+1}} \|u_{n-2, \bar{m}}\|_{H^s} \\
722 \quad &\leq (1/a^2) \mathcal{M}^2 |f|_{C^{s+2}}^2 K B^{n-2} D^{\bar{m}},
\end{aligned}$$

724 and

$$\begin{aligned}
725 \quad \|B_2^z \partial_z u_{n-2, \bar{m}}\|_{H^s} &= \|(-1/a^2)(a-z)(\partial_x f)^2 \partial_z u_{n-2, \bar{m}}\|_{H^s} \\
726 \quad &\leq (Z_a/a^2) \mathcal{M}^2 |\partial_x f|_{C^{s+1}} \|u_{n-2, \bar{m}}\|_{H^s} \\
727 \quad &\leq (Z_a/a^2) \mathcal{M}^2 |f|_{C^{s+2}}^2 K B^{n-2} D^{\bar{m}}.
\end{aligned}$$

729 To address the  $S_0, S_1, S_2$  terms we have

$$\begin{aligned}
730 \quad \|2i\alpha \partial_x u_{n, \bar{m}-1}\|_{H^s} &\leq 2\alpha \|u_{n, \bar{m}-1}\|_{H^{s+1}} \\
731 \quad &\leq 2\alpha K B^n D^{\bar{m}-1},
\end{aligned}$$

733 and

$$\begin{aligned}
734 \quad \|(\underline{\gamma}^u)^2 u_{n, \bar{m}-2}\|_{H^s} &\leq (\underline{\gamma}^u)^2 \|u_{n, \bar{m}-2}\|_{H^s} \\
735 \quad &\leq (\underline{\gamma}^u)^2 K B^n D^{\bar{m}-2},
\end{aligned}$$

737 and

$$\begin{aligned}
738 \quad \|2(\underline{\gamma}^u)^2 u_{n, \bar{m}-1}\|_{H^s} &\leq 2(\underline{\gamma}^u)^2 \|u_{n, \bar{m}-1}\|_{H^s} \\
739 \quad &\leq 2(\underline{\gamma}^u)^2 K B^n D^{\bar{m}-1},
\end{aligned}$$

741 and

$$\begin{aligned}
742 \quad \|2S_1 i\alpha \partial_x u_{n-1, \bar{m}}\|_{H^s} &= \|(-4/a) i\alpha f \partial_x u_{n-1, \bar{m}}\|_{H^s} \\
743 \quad &\leq (4/a) \alpha \mathcal{M} |f|_{C^s} \|u_{n-1, \bar{m}}\|_{H^{s+1}} \\
744 \quad &\leq (4/a) \alpha \mathcal{M} |f|_{C^s} K B^{n-1} D^{\bar{m}},
\end{aligned}$$

746 and

$$\begin{aligned}
747 \quad \|2S_1 i\alpha \partial_x u_{n-1, \bar{m}-1}\|_{H^s} &= \|(-4/a) i\alpha f \partial_x u_{n-1, \bar{m}-1}\|_{H^s} \\
748 \quad &\leq (4/a) \alpha \mathcal{M} |f|_{C^s} \|u_{n-1, \bar{m}-1}\|_{H^{s+1}} \\
749 \quad &\leq (4/a) \alpha \mathcal{M} |f|_{C^s} K B^{n-1} D^{\bar{m}-1},
\end{aligned}$$

751 and

$$\begin{aligned}
752 \quad \|S_1 (\underline{\gamma}^u)^2 u_{n-1, \bar{m}-2}\|_{H^s} &= \|(-2/a) (\underline{\gamma}^u)^2 f u_{n-1, \bar{m}-2}\|_{H^s} \\
753 \quad &\leq (2/a) (\underline{\gamma}^u)^2 \mathcal{M} |f|_{C^s} \|u_{n-1, \bar{m}-2}\|_{H^s} \\
754 \quad &\leq (2/a) (\underline{\gamma}^u)^2 \mathcal{M} |f|_{C^s} K B^{n-1} D^{\bar{m}-2},
\end{aligned}$$

756 and

$$\begin{aligned}
757 \quad \|2S_1 (\underline{\gamma}^u)^2 u_{n-1, \bar{m}-1}\|_{H^s} &= \|(-4/a) (\underline{\gamma}^u)^2 f u_{n-1, \bar{m}-1}\|_{H^s} \\
758 \quad &\leq (4/a) (\underline{\gamma}^u)^2 \mathcal{M} |f|_{C^s} \|u_{n-1, \bar{m}-1}\|_{H^s} \\
759 \quad &\leq (4/a) (\underline{\gamma}^u)^2 \mathcal{M} |f|_{C^s} K B^{n-1} D^{\bar{m}-1},
\end{aligned}$$

761 and

$$\begin{aligned}
762 \quad \|S_1 (\underline{\gamma}^u)^2 u_{n-1, \bar{m}}\|_{H^s} &= \|(-2/a) (\underline{\gamma}^u)^2 f u_{n-1, \bar{m}}\|_{H^s} \\
763 \quad &\leq (2/a) (\underline{\gamma}^u)^2 \mathcal{M} |f|_{C^s} \|u_{n-1, \bar{m}}\|_{H^s} \\
764 \quad &\leq (2/a) (\underline{\gamma}^u)^2 \mathcal{M} |f|_{C^s} K B^{n-1} D^{\bar{m}},
\end{aligned}$$

766 and

$$\begin{aligned}
767 \quad & \|2S_2 i \underline{\alpha} \partial_x u_{n-2, \bar{m}}\|_{H^s} = \|(2/a^2) i \underline{\alpha} f^2 \partial_x u_{n-2, \bar{m}}\|_{H^s} \\
768 \quad & \leq (2/a^2) \underline{\alpha} \mathcal{M}^2 |f|_{C^s}^2 \|u_{n-2, \bar{m}}\|_{H^{s+1}} \\
769 \quad & \leq (2/a^2) \underline{\alpha} \mathcal{M}^2 |f|_{C^s}^2 K B^{n-2} D^{\bar{m}},
\end{aligned}$$

771 and

$$\begin{aligned}
772 \quad & \|2S_2 i \underline{\alpha} \partial_x u_{n-2, \bar{m}-1}\|_{H^s} = \|(2/a^2) i \underline{\alpha} f^2 \partial_x u_{n-2, \bar{m}-1}\|_{H^s} \\
773 \quad & \leq (2/a^2) \underline{\alpha} \mathcal{M}^2 |f|_{C^s}^2 \|u_{n-2, \bar{m}-1}\|_{H^{s+1}} \\
774 \quad & \leq (2/a^2) \underline{\alpha} \mathcal{M}^2 |f|_{C^s}^2 K B^{n-2} D^{\bar{m}-1},
\end{aligned}$$

776 and

$$\begin{aligned}
777 \quad & \|S_2 (\underline{\gamma}^u)^2 u_{n-2, \bar{m}-2}\|_{H^s} = \|(1/a^2) (\underline{\gamma}^u)^2 f^2 u_{n-2, \bar{m}-2}\|_{H^s} \\
778 \quad & \leq (1/a^2) (\underline{\gamma}^u)^2 \mathcal{M}^2 |f|_{C^s}^2 \|u_{n-2, \bar{m}-2}\|_{H^s} \\
779 \quad & \leq (1/a^2) (\underline{\gamma}^u)^2 \mathcal{M}^2 |f|_{C^s}^2 K B^{n-2} D^{\bar{m}-2},
\end{aligned}$$

781 and

$$\begin{aligned}
782 \quad & \|2S_2 (\underline{\gamma}^u)^2 u_{n-2, \bar{m}-1}\|_{H^s} = \|(2/a^2) (\underline{\gamma}^u)^2 f^2 u_{n-2, \bar{m}-1}\|_{H^s} \\
783 \quad & \leq (2/a^2) (\underline{\gamma}^u)^2 \mathcal{M}^2 |f|_{C^s}^2 \|u_{n-2, \bar{m}-1}\|_{H^s} \\
784 \quad & \leq (2/a^2) (\underline{\gamma}^u)^2 \mathcal{M}^2 |f|_{C^s}^2 K B^{n-2} D^{\bar{m}-1},
\end{aligned}$$

786 and

$$\begin{aligned}
787 \quad & \|S_2 (\underline{\gamma}^u)^2 u_{n-2, \bar{m}}\|_{H^s} = \|(1/a^2) (\underline{\gamma}^u)^2 f^2 u_{n-2, \bar{m}}\|_{H^s} \\
788 \quad & \leq (1/a^2) (\underline{\gamma}^u)^2 \mathcal{M}^2 |f|_{C^s}^2 \|u_{n-2, \bar{m}}\|_{H^s} \\
789 \quad & \leq (1/a^2) (\underline{\gamma}^u)^2 \mathcal{M}^2 |f|_{C^s}^2 K B^{n-2} D^{\bar{m}}.
\end{aligned}$$

791 We satisfy the estimate for  $\|\tilde{F}_{n, \bar{m}}\|_{H^s}$  provided that we choose

$$\begin{aligned}
792 \quad & \bar{C} > \max \left\{ \left( 2\underline{\alpha} + 3(\underline{\gamma}^u)^2 \right), \left( \frac{3 + 2Z_a + 8\underline{\alpha} + 8(\underline{\gamma}^u)^2}{a} \right) \mathcal{M}, \right. \\
793 \quad & \left. \left( \frac{2 + 3Z_a + Z_a^2 + 4\underline{\alpha} + 4(\underline{\gamma}^u)^2}{a^2} \right) \mathcal{M}^2 \right\}. \\
794
\end{aligned}$$

795 The estimate for  $\tilde{P}_{n, \bar{m}}$  follows from the mapping properties of  $T^u$ ,

$$\begin{aligned}
796 \quad & \|\tilde{P}_{n, \bar{m}}\|_{H^{s+1/2}} = \|(1/a) f T^u [u_{n-1, \bar{m}}]\|_{H^{s+1/2}} \\
797 \quad & \leq (1/a) \mathcal{M} |f|_{C^{s+1/2+\eta}} \|T^u [u_{n-1, \bar{m}}]\|_{H^{s+1/2}} \\
798 \quad & \leq (1/a) \mathcal{M} |f|_{C^{s+1/2+\eta}} C_{T^u} \|u_{n-1, \bar{m}}\|_{H^{s+3/2}} \\
799 \quad & \leq (1/a) \mathcal{M} |f|_{C^{s+1/2+\eta}} C_{T^u} K B^{n-1} D^{\bar{m}}, \\
800
\end{aligned}$$

801 and provided that

$$802 \quad \bar{C} > (1/a) \mathcal{M} C_{T^u},$$

803 we are done. □

804 With this information, we can now prove Theorem 5.4.

805 *Proof.* [Theorem 5.4] We proceed by induction and at order  $m = 0$  Theorem 5.1  
806 guarantees a unique solution such that

$$807 \quad \|u_{n,0}\|_{H^{s+2}} \leq KB^n, \quad \forall n \geq 0.$$

808 We now assume the estimate (5.19) for all  $n, m < \bar{m}$  and study  $u_{n,\bar{m}}$ . From Theorem  
809 4.4 we have a unique solution satisfying

$$810 \quad \|u_{n,\bar{m}}\|_{H^{s+2}} \leq C_e \{ \|\tilde{F}_{n,\bar{m}}\|_{H^s} + \|U_{n,\bar{m}}\|_{H^{s+3/2}} + \|\tilde{P}_{n,\bar{m}}\|_{H^{s+1/2}} \},$$

811 and appealing to Lemmas 4.8 and 5.5 we find

$$812 \quad \|u_{n,\bar{m}}\|_{H^{s+2}} \leq C_e \left\{ K_U B_U^n D_U^{\bar{m}} + 2K\bar{C} \left( \underline{\alpha}(\underline{\gamma}^u)^2 B^n D^{\bar{m}-1} + (\underline{\gamma}^u)^2 B^n D^{\bar{m}-2} \right. \right. \\ 813 \quad \quad \quad \left. \left. + \underline{\alpha}(\underline{\gamma}^u)^2 |f|_{C^{s+2}} B^{n-1} D^{\bar{m}} + \underline{\alpha}(\underline{\gamma}^u)^2 |f|_{C^{s+2}} B^{n-1} D^{\bar{m}-1} \right. \right. \\ 814 \quad \quad \quad \left. \left. + (\underline{\gamma}^u)^2 |f|_{C^{s+2}} B^{n-1} D^{\bar{m}-2} + \underline{\alpha}(\underline{\gamma}^u)^2 |f|_{C^{s+2}}^2 B^{n-2} D^{\bar{m}} \right. \right. \\ 815 \quad \quad \quad \left. \left. + \underline{\alpha}(\underline{\gamma}^u)^2 |f|_{C^{s+2}}^2 B^{n-2} D^{\bar{m}-1} + (\underline{\gamma}^u)^2 |f|_{C^{s+2}}^2 B^{n-2} D^{\bar{m}-2} \right) \right\}. \\ 816$$

817 We are done provided we choose  $K \geq 9C_e K_U$  and

$$818 \quad B > \max \left\{ B_U, 18C_e \bar{C} \underline{\alpha}(\underline{\gamma}^u)^2 |f|_{C^{s+2}}, 18C_e \bar{C} (\underline{\gamma}^u)^2 |f|_{C^{s+2}}, \sqrt{18C_e \bar{C} \underline{\alpha}(\underline{\gamma}^u)^2 |f|_{C^{s+2}}}, \right. \\ 819 \quad \quad \left. \sqrt{18C_e \bar{C} (\underline{\gamma}^u)^2 |f|_{C^{s+2}}} \right\},$$

$$820 \quad D > \max \left\{ 1, D_U, 18C_e \bar{C} \underline{\alpha}(\underline{\gamma}^u)^2, \sqrt{18C_e \bar{C} (\underline{\gamma}^u)^2} \right\}. \\ 821$$

822 These inequalities are obtained from the bounds

$$823 \quad B > \max \left\{ B_U, 18C_e \bar{C} \underline{\alpha}(\underline{\gamma}^u)^2 |f|_{C^{s+2}}, \sqrt{18C_e \bar{C} \underline{\alpha}(\underline{\gamma}^u)^2 |f|_{C^{s+2}}} \right\},$$

$$824 \quad D > \max \left\{ D_U, 18C_e \bar{C} \underline{\alpha}(\underline{\gamma}^u)^2, \sqrt{18C_e \bar{C} (\underline{\gamma}^u)^2} \right\},$$

$$825 \quad BD > 18C_e \bar{C} \underline{\alpha}(\underline{\gamma}^u)^2 |f|_{C^{s+2}}, \quad BD^2 > 18C_e \bar{C} (\underline{\gamma}^u)^2 |f|_{C^{s+2}},$$

$$826 \quad B^2 D > 18C_e \bar{C} \underline{\alpha}(\underline{\gamma}^u)^2 |f|_{C^{s+2}}^2, \quad B^2 D^2 > 18C_e \bar{C} (\underline{\gamma}^u)^2 |f|_{C^{s+2}}^2. \quad \square \\ 827$$

828 As before, a similar analysis will establish the joint analyticity of the lower field  
829 which we now record.

830 **THEOREM 5.6.** *Given any integer  $s \geq 0$ , if  $f \in C^{s+2}([0, d])$  and  $W_{n,m} \in H^{s+3/2}([0, d])$*   
831 *such that*

$$832 \quad \|W_{n,m}\|_{H^{s+3/2}} \leq K_W B_W^n D_W^m,$$

833 *for constants  $K_W, B_W, D_W > 0$ , then  $w_{n,m} \in H^{s+2}([0, d] \times [-b, 0])$  and*

$$834 \quad \|w_{n,m}\|_{H^{s+2}} \leq KB^n D^m,$$

835 *for constants  $K, B, D > 0$ .*



836 **6. Analyticity of the Dirichlet–Neumann Operators.** Now that we have  
 837 established the joint analyticity of the upper field  $u$  we move to establishing the  
 838 analyticity of the upper layer DNO,  $G(g) = G(\varepsilon f)$ . To begin we give a recursive  
 839 estimate of the  $\tilde{H}_{n,m}$  appearing in (5.15).

840 LEMMA 6.1. *Given an integer  $s \geq 0$ , if  $f \in C^{s+2}([0, d])$  and*

$$841 \quad (6.1) \quad \|u_{n,m}\|_{H^{s+2}} \leq KB^n D^m, \quad \|G_{n,m}\|_{H^{s+1/2}} \leq \tilde{K} \tilde{B}^n \tilde{D}^m, \quad \forall n < \bar{n}, m,$$

842 *for constants  $K, B, D, \tilde{K}, \tilde{B}, \tilde{D} > 0$  where  $\tilde{K} \geq K, \tilde{B} \geq B, \tilde{D} \geq D$ , then there exists a*  
 843 *constant  $\tilde{C} > 0$  such that*

$$844 \quad (6.2) \quad \|\tilde{H}_{\bar{n},m}\|_{H^{s+1/2}} \leq \tilde{K} \tilde{C} \left\{ |f|_{C^{s+2}} \tilde{B}^{\bar{n}-1} \tilde{D}^m + |f|_{C^{s+2}}^2 \tilde{B}^{\bar{n}-2} \tilde{D}^m \right\}.$$

845 *Proof.* [Lemma 6.1] From (5.15) we estimate

$$\begin{aligned} 846 \quad \|\tilde{H}_{\bar{n},m}\|_{H^{s+1/2}} &\leq \mathcal{M} |\partial_x f|_{C^{s+1/2+\eta}} \|\partial_x u_{\bar{n}-1,m}(x, 0)\|_{H^{s+1/2}} \\ 847 \quad &+ \frac{1}{a} \mathcal{M} |f|_{C^{s+1/2+\eta}} \|G_{\bar{n}-1,m}(f)[U]\|_{H^{s+1/2}} \\ 848 \quad &+ \frac{1}{a} \mathcal{M}^2 |f|_{C^{s+1/2+\eta}} |\partial_x f|_{C^{s+1/2+\eta}} \|\partial_x u_{\bar{n}-2,m}(x, 0)\|_{H^{s+1/2}} \\ 849 \quad &+ \mathcal{M}^2 |\partial_x f|_{C^{s+1/2+\eta}}^2 \|\partial_z u_{\bar{n}-2,m}(x, 0)\|_{H^{s+1/2}}. \end{aligned}$$

851 This gives

$$\begin{aligned} 852 \quad \|\tilde{H}_{\bar{n},m}\|_{H^{s+1/2}} &\leq \tilde{K} \left\{ \mathcal{M} |f|_{C^{s+2}} \tilde{B}^{\bar{n}-1} \tilde{D}^m + \frac{1}{a} \mathcal{M} |f|_{C^{s+2}} \tilde{B}^{\bar{n}-1} \tilde{D}^m \right. \\ 853 \quad &+ \left. \frac{1}{a} \mathcal{M}^2 |f|_{C^{s+2}}^2 \tilde{B}^{\bar{n}-2} \tilde{D}^m + \mathcal{M}^2 |f|_{C^{s+2}}^2 \tilde{B}^{\bar{n}-2} \tilde{D}^m \right\}, \\ 854 \end{aligned}$$

855 and we are done provided

$$856 \quad \tilde{C} \geq \left(1 + \frac{1}{a}\right) \max\{\mathcal{M}, \mathcal{M}^2\}. \quad \square$$

858 We now have everything we need to prove the analyticity of the upper layer DNO.

859 THEOREM 6.2. *Given any integer  $s \geq 0$ , if  $f \in C^{s+2}([0, d])$  and  $U_{n,m} \in H^{s+3/2}([0, d])$  ■*  
 860 *such that*

$$861 \quad \|U_{n,m}\|_{H^{s+3/2}} \leq K_U B_U^n D_U^m,$$

862 *for constants  $K_U, B_U, D_U > 0$ , then  $G_{n,m} \in H^{s+1/2}([0, d])$  and*

$$863 \quad (6.3) \quad \|G_{n,m}\|_{H^{s+1/2}} \leq \tilde{K} \tilde{B}^n \tilde{D}^m,$$

864 *for constants  $\tilde{K}, \tilde{B}, \tilde{D} > 0$ .*

865 *Proof.* [Theorem 6.2] As before, we work by induction. At  $n = 0$  we have from  
 866 (5.12) that

$$867 \quad G_{0,m} = -\partial_z u_{0,m}(x, 0),$$

868 and from Theorem 5.4 we have

$$869 \quad \|G_{0,m}\|_{H^{s+1/2}} = \|\partial_z u_{0,m}(x, 0)\|_{H^{s+1/2}} \leq \|u_{0,m}\|_{H^{s+2}} \leq KD^m.$$

870 So we choose  $\tilde{K} \geq K$  and  $\tilde{D} \geq D$ . We now assume  $\tilde{B} \geq B$  and the estimate (6.3) for  
871 all  $n < \bar{n}$ ; from (5.12) we have

$$872 \quad \|G_{\bar{n},m}(f)[U]\|_{H^{s+1/2}} \leq \|\partial_z u_{\bar{n},m}(x,0)\|_{H^{s+1/2}} + \|\tilde{H}_{\bar{n},m}(x)\|_{H^{s+1/2}}.$$

873 Using the inductive hypothesis, Lemma 6.1, and Theorem 5.4 we have

$$874 \quad \|G_{\bar{n},m}(f)[U]\|_{H^{s+1/2}} \leq KB\bar{n}D^m + \tilde{K}\tilde{C} \left\{ |f|_{C^{s+2}}\tilde{B}^{\bar{n}-1}\tilde{D}^m + |f|_{C^{s+2}}^2\tilde{B}^{\bar{n}-2}\tilde{D}^m \right\}.$$

875 We are done provided  $\tilde{K} \geq 2K$ ,  $\tilde{D} \geq D$ , and

$$876 \quad \tilde{B} \geq \max \left\{ B, 4\tilde{C}|f|_{C^{s+2}}, 2\sqrt{\tilde{C}}|f|_{C^{s+2}} \right\}. \quad \square$$

877 Finally, a similar approach will give the joint analyticity of the DNO in the lower  
878 field.

879 **THEOREM 6.3.** *Given any integer  $s \geq 0$ , if  $f \in C^{s+2}([0, d])$  and  $W_{n,m} \in H^{s+3/2}([0, d])$*  ■  
880 *such that*

$$881 \quad \|W_{n,m}\|_{H^{s+3/2}} \leq K_W B_W^n D_W^m,$$

882 *for constants  $K_W, B_W, D_W > 0$ , then  $J_{n,m} \in H^{s+1/2}([0, d])$  and*

$$883 \quad (6.4) \quad \|J_{n,m}\|_{H^{s+1/2}} \leq \tilde{K}\tilde{B}^n\tilde{D}^m,$$

884 *for constants  $\tilde{K}, \tilde{B}, \tilde{D} > 0$ .*

885 **Acknowledgments.** D.P.N. gratefully acknowledges support from the National  
886 Science Foundation through grants No. DMS-1813033 and No. DMS-2111283.

## 887 REFERENCES

- 888 [1] T. ABBOUD AND J. C. NEDELEC, *Electromagnetic waves in an inhomogeneous medium*, J. Math.  
889 Anal. Appl., 164 (1992), pp. 40–58.
- 890 [2] R. A. ADAMS, *Sobolev spaces*, Academic Press [A subsidiary of Harcourt Brace Jovanovich,  
891 Publishers], New York-London, 1975. Pure and Applied Mathematics, Vol. 65.
- 892 [3] T. ARENS, *Scattering by Biperiodic Layered Media: The Integral Equation Approach*, habilita-  
893 tionsschrift, Karlsruhe Institute of Technology, 2009.
- 894 [4] G. BAO, *Finite element approximation of time harmonic waves in periodic structures*, SIAM  
895 J. Numer. Anal., 32 (1995), pp. 1155–1169.
- 896 [5] G. BAO, L. COWSAR, AND W. MASTERS, *Mathematical Modeling in Optical Science*, SIAM,  
897 Philadelphia, 2001.
- 898 [6] G. BAO, D. C. DOBSON, AND J. A. COX, *Mathematical studies in rigorous grating theory*, J.  
899 Opt. Soc. Amer. A, 12 (1995), pp. 1029–1042.
- 900 [7] J.-P. BÉRENGER, *A perfectly matched layer for the absorption of electromagnetic waves*, J.  
901 Comput. Phys., 114 (1994), pp. 185–200.
- 902 [8] F. BLEIBINHAUS AND S. RONDENAY, *Effects of surface scattering in full-waveform inversion*,  
903 Geophysics, 74 (2009), pp. WCC69–WCC77.
- 904 [9] J. P. BOYD, *Chebyshev and Fourier spectral methods*, Dover Publications Inc., Mineola, NY,  
905 second ed., 2001.
- 906 [10] L. M. BREKHOVSKIKH AND Y. P. LYSANOV, *Fundamentals of Ocean Acoustics*, Springer-Verlag,  
907 Berlin, 1982.
- 908 [11] O. BRUNO AND F. REITICH, *Numerical solution of diffraction problems: A method of variation*  
909 *of boundaries*, J. Opt. Soc. Am. A, 10 (1993), pp. 1168–1175.
- 910 [12] O. BRUNO AND F. REITICH, *Numerical solution of diffraction problems: A method of variation*  
911 *of boundaries. II. Finitely conducting gratings, Padé approximants, and singularities*, J.  
912 Opt. Soc. Am. A, 10 (1993), pp. 2307–2316.

- 913 [13] O. BRUNO AND F. REITICH, *Numerical solution of diffraction problems: A method of variation*  
914 *of boundaries. III. Doubly periodic gratings*, J. Opt. Soc. Am. A, 10 (1993), pp. 2551–2562.
- 915 [14] J. CHANDEZON, M. DUPUIS, G. CORNET, AND D. MAYSTRE, *Multicoated gratings: a differential*  
916 *formalism applicable in the entire optical region*, J. Opt. Soc. Amer., 72 (1982), p. 839.
- 917 [15] J. CHANDEZON, D. MAYSTRE, AND G. RAOULT, *A new theoretical method for diffraction gratings*  
918 *and its numerical application*, J. Opt., 11 (1980), pp. 235–241.
- 919 [16] X. CHEN AND A. FRIEDMAN, *Maxwell's equations in a periodic structure*, Trans. Amer. Math.  
920 Soc., 323 (1991), pp. 465–507.
- 921 [17] D. COLTON AND R. KRESS, *Inverse acoustic and electromagnetic scattering theory*, vol. 93 of  
922 Applied Mathematical Sciences, Springer, New York, third ed., 2013.
- 923 [18] B. DESPRÉS, *Domain decomposition method and the Helmholtz problem*, in Mathematical and  
924 numerical aspects of wave propagation phenomena (Strasbourg, 1991), SIAM, Philadelphia,  
925 PA, 1991, pp. 44–52.
- 926 [19] B. DESPRÉS, *Méthodes de décomposition de domaine pour les problèmes de propagation d'ondes*  
927 *en régime harmonique. Le théorème de Borg pour l'équation de Hill vectorielle*, Institut  
928 National de Recherche en Informatique et en Automatique (INRIA), Rocquencourt, 1991.  
929 Thèse, Université de Paris IX (Dauphine), Paris, 1991.
- 930 [20] M. O. DEVILLE, P. F. FISCHER, AND E. H. MUND, *High-order methods for incompressible*  
931 *fluid flow*, vol. 9 of Cambridge Monographs on Applied and Computational Mathematics,  
932 Cambridge University Press, Cambridge, 2002.
- 933 [21] D. DOBSON AND A. FRIEDMAN, *The time-harmonic Maxwell equations in a doubly periodic*  
934 *structure*, J. Math. Anal. Appl., 166 (1992), pp. 507–528.
- 935 [22] D. C. DOBSON, *A variational method for electromagnetic diffraction in biperiodic structures*,  
936 RAIRO Modél. Math. Anal. Numér., 28 (1994), pp. 419–439.
- 937 [23] T. W. EBBESEN, H. J. LEZEC, H. F. GHAEMI, T. THIO, AND P. A. WOLFF, *Extraordinary*  
938 *optical transmission through sub-wavelength hole arrays*, Nature, 391 (1998), pp. 667–669.
- 939 [24] S. ENOCH AND N. BONOD, *Plasmonics: From Basics to Advanced Topics*, Springer Series in  
940 Optical Sciences, Springer, New York, 2012.
- 941 [25] L. C. EVANS, *Partial differential equations*, American Mathematical Society, Providence, RI,  
942 second ed., 2010.
- 943 [26] G. B. FOLLAND, *Introduction to partial differential equations*, Princeton University Press,  
944 Princeton, N.J., 1976. Preliminary informal notes of university courses and seminars in  
945 mathematics, Mathematical Notes.
- 946 [27] D. GILBARG AND N. S. TRUDINGER, *Elliptic partial differential equations of second order*,  
947 Springer-Verlag, Berlin, second ed., 1983.
- 948 [28] *Solids far from equilibrium*, Cambridge University Press, Cambridge, 1992.
- 949 [29] D. GOTTLIEB AND S. A. ORSZAG, *Numerical analysis of spectral methods: theory and applica-*  
950 *tions*, Society for Industrial and Applied Mathematics, Philadelphia, Pa., 1977. CBMS-NSF  
951 Regional Conference Series in Applied Mathematics, No. 26.
- 952 [30] J. S. HESTHAVEN AND T. WARBURTON, *Nodal discontinuous Galerkin methods*, vol. 54 of Texts  
953 in Applied Mathematics, Springer, New York, 2008. Algorithms, analysis, and applications.
- 954 [31] J. HOMOLA, *Surface plasmon resonance sensors for detection of chemical and biological species*,  
955 Chemical Reviews, 108 (2008), pp. 462–493.
- 956 [32] Y. HONG AND D. P. NICHOLLS, *A rigorous numerical analysis of the transformed field expansion*  
957 *method for diffraction by periodic, layered structures*, SIAM Journal on Numerical Analysis,  
958 59 (2021), pp. 456–476.
- 959 [33] H. IM, S. H. LEE, N. J. WITTENBERG, T. W. JOHNSON, N. C. LINDQUIST, P. NAGPAL, D. J.  
960 NORRIS, AND S.-H. OH, *Template-stripped smooth Ag nanohole arrays with silica shells for*  
961 *surface plasmon resonance biosensing*, ACS Nano, 5 (2011), pp. 6244–6253.
- 962 [34] C. JOHNSON, *Numerical solution of partial differential equations by the finite element method*,  
963 Cambridge University Press, Cambridge, 1987.
- 964 [35] J. JOSE, L. R. JORDAN, T. W. JOHNSON, S. H. LEE, N. J. WITTENBERG, AND S.-H. OH,  
965 *Topographically flat substrates with embedded nanoplasmic devices for biosensing*, Adv  
966 Funct Mater, 23 (2013), pp. 2812–2820.
- 967 [36] M. KEHOE AND D. P. NICHOLLS, *A stable high-order perturbation of surfaces/asymptotic wave-*  
968 *form evaluation method for the numerical solution of grating scattering problems*, SIAM  
969 Journal on Scientific Computing (submitted), (2021).
- 970 [37] R. KRESS, *Linear integral equations*, Springer-Verlag, New York, third ed., 2014.
- 971 [38] O. A. LADYZHENSKAYA AND N. N. URAL'TSEVA, *Linear and quasilinear elliptic equations*, Aca-  
972 demic Press, New York, 1968.
- 973 [39] N. LASSALINE, R. BRECHBÜHLER, S. VONK, K. RIDDERBEEK, M. SPIESER, S. BISIG,  
974 B. LE FEBER, F. RABOUW, AND D. NORRIS, *Optical fourier surfaces*, Nature, 582 (2020),

- 975 pp. 506–510.
- 976 [40] R. J. LEVEQUE, *Finite difference methods for ordinary and partial differential equations*, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2007. Steady-state and time-dependent problems.
- 977
- 978
- 979 [41] E. H. LIEB AND M. LOSS, *Analysis*, vol. 14 of Graduate Studies in Mathematics, American Mathematical Society, Providence, RI, second ed., 2001.
- 980
- 981 [42] N. C. LINDQUIST, T. W. JOHNSON, J. JOSE, L. M. OTTO, AND S.-H. OH, *Ultrasmooth metallic films with buried nanostructures for backside reflection-mode plasmonic biosensing*, *Annalen der Physik*, 524 (2012), pp. 687–696.
- 982
- 983
- 984 [43] P.-L. LIONS, *On the Schwarz alternating method. III. A variant for nonoverlapping subdomains*, in Third International Symposium on Domain Decomposition Methods for Partial Differential Equations (Houston, TX, 1989), SIAM, Philadelphia, PA, 1990, pp. 202–223.
- 985
- 986
- 987 [44] S. A. MAIER, *Plasmonics: Fundamentals and Applications*, Springer, New York, 2007.
- 988 [45] D. M. MILDER, *An improved formalism for rough-surface scattering of acoustic and electromagnetic waves*, in Proceedings of SPIE - The International Society for Optical Engineering (San Diego, 1991), vol. 1558, Int. Soc. for Optical Engineering, Bellingham, WA, 1991, pp. 213–221.
- 991
- 992 [46] D. M. MILDER, *An improved formalism for wave scattering from rough surfaces*, *J. Acoust. Soc. Am.*, 89 (1991), pp. 529–541.
- 993
- 994 [47] M. MOSKOVITS, *Surface-enhanced spectroscopy*, *Reviews of Modern Physics*, 57 (1985), pp. 783–826.
- 995
- 996 [48] F. NATTERER AND F. WÜBBELING, *Mathematical methods in image reconstruction*, SIAM Monographs on Mathematical Modeling and Computation, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2001.
- 997
- 998
- 999 [49] D. P. NICHOLLS, *Three-dimensional acoustic scattering by layered media: A novel surface formulation with operator expansions implementation*, *Proceedings of the Royal Society of London, A*, 468 (2012), pp. 731–758.
- 1000
- 1001
- 1002 [50] D. P. NICHOLLS, *Numerical solution of diffraction problems: A high-order perturbation of surfaces/asymptotic waveform evaluation method*, *SIAM Journal on Numerical Analysis*, 55 (2017), pp. 144–167.
- 1003
- 1004
- 1005 [51] D. P. NICHOLLS, *On analyticity of linear waves scattered by a layered medium*, *Journal of Differential Equations*, 263 (2017), pp. 5042–5089.
- 1006
- 1007 [52] D. P. NICHOLLS, *Numerical simulation of grating structures incorporating two-dimensional materials: A high-order perturbation of surfaces framework*, *SIAM Journal on Applied Mathematics*, 78 (2018), pp. 19–44.
- 1008
- 1009
- 1010 [53] D. P. NICHOLLS AND F. REITICH, *A new approach to analyticity of Dirichlet-Neumann operators*, *Proc. Roy. Soc. Edinburgh Sect. A*, 131 (2001), pp. 1411–1433.
- 1011
- 1012 [54] D. P. NICHOLLS AND F. REITICH, *Stability of high-order perturbative methods for the computation of Dirichlet-Neumann operators*, *J. Comput. Phys.*, 170 (2001), pp. 276–298.
- 1013
- 1014 [55] D. P. NICHOLLS AND F. REITICH, *Analytic continuation of Dirichlet-Neumann operators*, *Numer. Math.*, 94 (2003), pp. 107–146.
- 1015
- 1016 [56] D. P. NICHOLLS AND F. REITICH, *Shape deformations in rough surface scattering: Improved algorithms*, *J. Opt. Soc. Am. A*, 21 (2004), pp. 606–621.
- 1017
- 1018 [57] D. P. NICHOLLS AND J. SHEN, *A rigorous numerical analysis of the transformed field expansion method*, *SIAM Journal on Numerical Analysis*, 47 (2009), pp. 2708–2734.
- 1019
- 1020 [58] D. P. NICHOLLS AND M. TABER, *Joint analyticity and analytic continuation for Dirichlet-Neumann operators on doubly perturbed domains*, *J. Math. Fluid Mech.*, 10 (2008), pp. 238–271.
- 1021
- 1022
- 1023 [59] *Electromagnetic theory of gratings*, Springer-Verlag, Berlin, 1980.
- 1024 [60] N. A. PHILLIPS, *A coordinate system having some special advantages for numerical forecasting*, *Journal of the Atmospheric Sciences*, 14 (1957), pp. 184–185.
- 1025
- 1026 [61] H. RAETHER, *Surface plasmons on smooth and rough surfaces and on gratings*, Springer, Berlin, 1988.
- 1027
- 1028 [62] S. A. SAUTER AND C. SCHWAB, *Boundary element methods*, vol. 39 of Springer Series in Computational Mathematics, Springer-Verlag, Berlin, 2011. Translated and expanded from the 2004 German original.
- 1029
- 1030
- 1031 [63] J. SHEN, T. TANG, AND L.-L. WANG, *Spectral methods*, vol. 41 of Springer Series in Computational Mathematics, Springer, Heidelberg, 2011. Algorithms, analysis and applications.
- 1032
- 1033 [64] J. VIRIEUX AND S. OPERTO, *An overview of full-waveform inversion in exploration geophysics*, *Geophysics*, 74 (2009), pp. WCC1–WCC26.
- 1034
- 1035 [65] C. H. WILCOX, *Scattering Theory for Diffraction Gratings*, Springer, Berlin, 1984.