

DISCONNECTED MODULI SPACES OF STABLE BUNDLES ON SURFACES

IZZET COSKUN, JACK HUIZENGA, AND JOHN KOPPER

ABSTRACT. We use hypersurfaces containing unexpected linear spaces to construct interesting vector bundles on complete intersection surfaces in projective space. We discover examples of moduli spaces of rank 2 stable bundles on surfaces of Picard rank one with arbitrarily many connected components.

CONTENTS

1. Introduction	1
2. Preliminaries	2
3. The construction	4
4. Examples of moduli spaces	8
References	9

1. INTRODUCTION

In this note we construct examples of moduli spaces of rank two bundles on surfaces of Picard rank one with arbitrarily many connected components.

On a smooth projective curve, the moduli space of semistable bundles of rank r and degree d is irreducible [HuL, Cor. 4.5.5]. In contrast, if $\dim(X) \geq 3$, then even the Hilbert scheme of points on X is reducible with components of different dimensions [Ia72]. There are many constructions of reducible and disconnected moduli spaces if $\dim(X) \geq 3$ (for example, see [Ei88]). Moduli spaces of sheaves on surfaces exhibit the most interesting behavior. The Hilbert scheme of n points on a smooth, irreducible projective surface is smooth and irreducible [Fo68]. In higher rank, the philosophy of Donaldson [Do90] and Gieseker and Li [GL94] expects the moduli spaces $M_X(r, c_1, c_2)$ to become better behaved as the second Chern class c_2 tends to infinity. For example, they become reduced and irreducible of the expected dimension if c_2 is sufficiently large (see [O’G96] and [Ni95, MS16] for sharper inequalities for surfaces in \mathbb{P}^3). On the other hand, if c_2 is small, then $M_X(r, c_1, c_2)$ may be reducible with components of different dimensions and may have nonreduced components [Mes97, MS11]. For example, [CH18] and [Fr89] construct moduli spaces with arbitrarily many irreducible components. It is reasonable to expect that for small c_2 , the moduli spaces of sheaves on surfaces satisfy a version of Murphy’s Law [Va06].

One can construct disconnected moduli spaces of sheaves on threefolds by using the Serre correspondence, which relates codimension 2 subschemes to rank 2 vector bundles (see §2 or [St19]). Subschemes with a given Hilbert polynomial are parameterized by a Hilbert scheme. If this Hilbert scheme is disconnected, the same can be true of the moduli space of sheaves associated through the Serre correspondence. We use a variant of this argument by relating the moduli space $M_X(r, c_1, c_2)$

Date: May 19, 2021.

2010 *Mathematics Subject Classification.* Primary: 14J60, 14J29. Secondary: 14D20.

During the preparation of this article the first author was partially supported by the NSF FRG grant DMS 1664296 and the second author was partially supported by NSF FRG grant DMS 1664303.

on the surface X to a disconnected Hilbert scheme of linear spaces on an ambient higher-dimensional variety.

Okonek and Van de Ven [OV86] and Kotschick [Ko89] found examples of disconnected moduli spaces on elliptic surfaces in their study of the topology of the underlying real fourfold (see also [FM88]). These disconnected moduli spaces are on certain elliptic surfaces with large Picard rank. In contrast, our construction uses general type surfaces with Picard rank one.

Let $n \geq 4$ be an integer. Let $2 < d_1 < d_2 \leq \dots \leq d_{n-2}$ be integers. If $n = 4$, further assume that $d_1 \geq 4$ and $d_2 \geq 6$. Set $e = \prod_{i=2}^{n-2} d_i$. Let $D_1 \subset \mathbb{P}^n$ be a hypersurface of degree d_1 that contains a linear space Φ of dimension $n - 3$. Assume that the singular locus of D_1 has codimension 5. For $2 \leq i \leq n - 2$, let $D_i \subset \mathbb{P}^n$ be very general hypersurfaces of degree d_i . Let X be the complete intersection $D_1 \cap \dots \cap D_{n-2}$, which is a smooth, projective surface of degree $d_1 e$. Let H denote the hyperplane class on X . Let $F_{n-3}(D_1)$ denote the Fano scheme parameterizing linear spaces on D_1 of dimension $n - 3$.

Theorem (3.4). *Every connected component of $F_{n-3}(D_1)$ corresponds to a distinct connected component of $M_X(2, H, (d_1 - 1)e)$ of the same dimension.*

In §4, we give two families of examples of $D_1 \subset \mathbb{P}^4$. In Example 4.2, $F_1(D_1)$ has $3d_1^2$ zero-dimensional connected components. In Example 4.5, $F_1(D_1)$ has d_1 one-dimensional connected components. Example 4.4 gives examples of $D_1 \subset \mathbb{P}^5$ where $F_2(D_1)$ has $15d_1^3$ zero-dimensional connected components. We conclude the following.

Corollary 1.1. *For any integer k , there exists a smooth surface X and Chern character \mathbf{v} on X such that $M_X(\mathbf{v})$ has at least k connected components.*

The original motivation for this note was a question about monotonicity of Betti numbers of $M_X(r, c_1, c_2)$. Coskun and Woolf [CW20] conjecture that as c_2 tends to infinity, the Betti numbers of $M_X(r, c_1, c_2)$ stabilize. Göttsche proved that the Betti numbers of the Hilbert scheme of n points $X^{[n]}$ stabilize as n tends to infinity [Go90]. In fact, the Betti numbers monotonically increase to the stable value as n increases. The same monotonicity occurs for certain moduli spaces of higher rank sheaves on K3 surfaces and rational surfaces. In talks given by the first author, several mathematicians raised the question whether one should expect the Betti numbers to always monotonically increase to the stable value. As examples of disconnected moduli spaces show, this is already false for b_0 .

Corollary 1.2. *The Betti numbers of the moduli spaces of sheaves $M_X(r, c_1, c_2)$ do not monotonically increase as c_2 increases.*

One could refine the question and ask whether the Betti numbers monotonically increase once c_2 reaches the O’Grady bound and the moduli spaces become reduced and irreducible of the expected dimension.

Organization of the paper. In §2, we recall basic facts about moduli spaces and complete intersections. In §3, we describe our main construction and prove Theorem 3.4. In §4, we give examples that illustrate interesting phenomena.

Acknowledgments. We would like to thank Lawrence Ein, Joe Harris, Daniel Huybrechts, Nicole Mestrano, Julius Ross, Carlos Simpson, Dennis Tseng, Matthew Woolf and Kota Yoshioka for valuable conversations. We thank the organizers and participants of ZAG, Rahul Pandharipande, and the participants of the Algebraic Geometry and Moduli seminar for stimulating discussions that led to this note.

2. PRELIMINARIES

In this section, we recall basic facts concerning moduli spaces of sheaves (see [HuL, MS16] for further details) and complete intersections.

Moduli spaces of sheaves. Let (X, H) be a smooth, polarized, complex projective variety of dimension n . If \mathcal{F} is a torsion-free coherent sheaf on X of rank $\text{rk}(\mathcal{F})$, then the Hilbert polynomial $P_{\mathcal{F}}(m)$ and the reduced Hilbert polynomial $p_{\mathcal{F}}(m)$ of \mathcal{F} are defined by

$$P_{\mathcal{F}}(m) = \chi(\mathcal{F}(mH)) = \text{rk}(\mathcal{F}) \frac{m^n}{n!} + \text{l.o.t.} \quad \text{and} \quad p_{\mathcal{F}}(m) = \frac{P_{\mathcal{F}}(m)}{\text{rk}(\mathcal{F})}.$$

The sheaf \mathcal{F} is called *(semi)stable* if for every proper subsheaf $\mathcal{F}' \subset \mathcal{F}$, we have $p_{\mathcal{F}'}(m) \underset{(-)}{\leq} p_{\mathcal{F}}(m)$ for $m \gg 0$. Gieseker and Maruyama constructed projective moduli spaces $M_X(\mathbf{v})$ parameterizing S-equivalence classes of semistable sheaves on X with Chern character \mathbf{v} [Gi77, Ma78]. The *H-slope* of \mathcal{F} is defined to be the number

$$\mu_H(\mathcal{F}) = \frac{c_1(\mathcal{F}) \cdot H^{n-1}}{\text{rk}(\mathcal{F})H^n}.$$

If $\mu_H(\mathcal{F}') < \mu_H(\mathcal{F})$ for all proper subsheaves $\mathcal{F}' \subset \mathcal{F}$, then \mathcal{F} is called μ_H -stable.

The Serre correspondence. Let X be a smooth projective surface. Let $Z \subset X$ be a zero-dimensional Gorenstein subscheme of length n . Then Z satisfies the *Cayley-Bacharach property* for a line bundle \mathcal{L} on X if for any subscheme $W \subset Z$ of length $n-1$, any section of \mathcal{L} vanishing on W vanishes on all of Z . Given a line bundle \mathcal{L} and a zero-dimensional scheme Z satisfying the Cayley-Bacharach property for $\omega_X \otimes \mathcal{L}$, the Serre correspondence constructs a vector bundle of rank 2 on X .

Theorem 2.1 (The Serre Correspondence [HuL, 5.1.1]). *Let X be a smooth projective surface, and $Z \subset X$ be a local complete intersection subscheme of dimension 0 and length n . Let \mathcal{L} be a line bundle on X . Then there exists an extension*

$$0 \rightarrow \mathcal{O}_X \rightarrow \mathcal{E} \rightarrow \mathcal{L} \otimes I_Z \rightarrow 0$$

with \mathcal{E} locally free if and only if Z satisfies the Cayley-Bacharach property for $\omega_X \otimes \mathcal{L}$.

The Cayley-Bacharach Theorem. For our construction, we will use the following Cayley-Bacharach theorem for projective space (see [EGH, Theorem CB7])¹.

Theorem 2.2 (Cayley-Bacharach). *Let $D_1, D_2, \dots, D_n \subset \mathbb{P}^n$ be divisors of degrees d_1, d_2, \dots, d_n respectively, meeting in a zero-dimensional scheme $Z = D_1 \cap D_2 \cap \dots \cap D_n$. Set $d = \sum_{i=1}^n d_i$.*

- (1) *If $D \subset \mathbb{P}^n$ is any divisor of degree $d - n - 1$ containing a subscheme Z' of Z of length one less than that of Z , then D contains all of Z .*
- (2) *Suppose Z', Z'' are residual subschemes of Z . Then for any $m \leq d - n - 1$, we have*

$$h^0(\mathbb{P}^n, I_{Z'}(m)) - h^0(\mathbb{P}^n, I_Z(m)) = h^1(\mathbb{P}^n, I_{Z''}(d - n - 1 - m)).$$

Complete intersections. Finally, we will use the following standard facts about complete intersections and their Fano schemes of linear spaces. Given a variety X , let $F_j(X)$ denote the Fano scheme parameterizing j -dimensional linear spaces contained in X .

- Proposition 2.3.**
- (1) *If $D \subset \mathbb{P}^4$ is a smooth hypersurface of degree $d \geq 4$, then $\dim(F_1(D)) \leq 1$. If $d \geq 6$ is and D is general, then $F_1(D)$ is empty.*
 - (2) *If $D \subset \mathbb{P}^5$ is a smooth hypersurface of degree $d \geq 3$, then $\dim(F_2(D)) \leq 0$. If D is general, then $F_2(D)$ is empty.*
 - (3) *Let $n \geq 5$. If $D \subset \mathbb{P}^n$ is a hypersurface of degree $d \geq 3$ which is smooth in codimension 4, then $\dim(F_{n-3}(D)) \leq 0$.*

¹The statement of Theorem CB7 in *op. cit.* should say divisors instead of curves.

Proof. Let $D \subset \mathbb{P}^4$ be a smooth hypersurface of degree d . Let Y be a component of $F_1(D)$ containing a line ℓ . If $d \geq 4$, then $\deg(N_{\ell/D}) = 3 - d < 0$. Hence, ℓ is not free and the lines parameterized by Y can only sweep a surface. If $\dim(Y) \geq 2$, this surface must be a plane. However, by the Lefschetz hyperplane theorem, a smooth hypersurface of degree $d > 1$ cannot contain a plane. We conclude that $\dim(F_1(D)) \leq 1$. A dimension count shows that a general hypersurface in \mathbb{P}^n of degree $d \geq 2n - 2$ does not contain lines. This proves (1).

Let $D \subset \mathbb{P}^5$ be a smooth hypersurface of degree $d \geq 3$ that contains a plane Λ . Choosing coordinates, we may assume that Λ is defined by $z_0 = z_1 = z_2 = 0$ and that the equation of the hypersurface is given by $z_0 F_0 + z_1 F_1 + z_2 F_2 = 0$, where F_i are polynomials of degree $d - 1$. For $1 \leq i \leq 3$, let \bar{F}_i denote the restriction of F_i to Λ . If the \bar{F}_i have a common zero on Λ , then D is singular at that point. The standard normal bundle sequence for $N_{\Lambda/D}$ is given by

$$0 \rightarrow N_{\Lambda/D} \rightarrow \mathcal{O}_{\Lambda}(1)^3 \xrightarrow{(\bar{F}_0, \bar{F}_1, \bar{F}_2)} \mathcal{O}_{\Lambda}(d) \rightarrow 0.$$

If $N_{\Lambda/D}$ has a section, then the \bar{F}_i must satisfy a relation of the form $\sum L_i \bar{F}_i = 0$ for some linear forms L_i . Since the polynomials \bar{F}_i have degree at least 2, they must have a common solution, contradicting the smoothness of D . Therefore $H^0(\Lambda, N_{\Lambda/D}) = 0$ and $\dim(F_2(D)) \leq 0$. Finally, a dimension count shows that for general D , $F_2(D)$ is empty. This proves (2), and (3) is an immediate consequence of (2) by intersecting with a general \mathbb{P}^5 . \square

Proposition 2.4. *Let $4 \leq d_1 \leq \dots \leq d_{n-3}$ be a sequence of integers such that $\sum_{i=1}^{n-3} d_i > n + 1$. Let $X \subset \mathbb{P}^n$ be a very general complete intersection of type d_1, \dots, d_{n-3} . Then any codimension 3 linear space intersects X in dimension 0.*

Proof. When $n = 4$ or 5 , the proposition asserts that X does not contain a line or a plane curve, respectively. A simple dimension count shows that this is the case as soon as $\sum_{i=1}^{n-3} d_i > n + 1$. When $n > 5$, it suffices to show that the codimension of the locus of forms (D_1, \dots, D_{n-3}) that contain a curve in \mathbb{P}^{n-3} is greater than the dimension of the Grassmannian $\mathbb{G}(n-3, n)$. This follows from the dimension counts in [Ts20, Lemmas 4.2 and 4.3]. \square

3. THE CONSTRUCTION

In this section, we will describe our construction and prove its main properties. We first establish some notation.

The surfaces. Let $n \geq 4$ be an integer. Let $2 < d_1 < d_2 \leq \dots \leq d_{n-2}$ be integers such that $d = \sum_{i=1}^{n-2} d_i \geq n + 1$. Set $e = \prod_{i=2}^{n-2} d_i$. Let $D_1 \subset \mathbb{P}^n$ be a hypersurface of degree d_1 that contains a linear space Φ of dimension $n - 3$. When $4 \leq n \leq 5$, we will assume that D_1 is smooth. If $n > 5$, then D_1 cannot be smooth, but we will assume that its singular locus has codimension 5. For $2 \leq i \leq n - 2$, let $D_i \subset \mathbb{P}^n$ be very general hypersurfaces of degree d_i . Let X be the complete intersection $D_1 \cap \dots \cap D_{n-2}$, which is a smooth, projective surface of degree $d_1 e$. Let H denote the hyperplane class on X . The following proposition summarizes the basic properties of X .

Proposition 3.1. *The surface X satisfies the following properties.*

- (1) *The canonical class of X is $K_X = (d - n - 1)H$.*
- (2) *We have $\text{Pic } X = \mathbb{Z}H$.*
- (3) *For any integer m and $i = 1, 2$, $h^i(\mathbb{P}^n, I_X(m)) = 0$.*
- (4) *For any integer m , $h^1(X, \mathcal{O}_X(m)) = 0$ and the map $H^0(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(m)) \rightarrow H^0(X, \mathcal{O}_X(m))$ is surjective.*

Proof. The canonical class of X is computed by adjunction. The intersection $Y = \cap_{i=1}^{n-3} D_i$ is a smooth, complete intersection threefold. By the Lefschetz hyperplane theorem, $\text{Pic}(Y)$ is cyclic

generated by the restriction of the hyperplane class. By assumption D_{n-2} is very general and K_X has a section, so Moishezon's Noether-Lefschetz theorem [Mo67] implies that $\text{Pic}(X) \cong \mathbb{Z}H$.

Set $W = \bigoplus_{i=1}^{n-2} \mathcal{O}_{\mathbb{P}^n}(-d_i)$. Since X is a complete intersection, the Koszul complex

$$0 \rightarrow \bigwedge^{n-2} W \rightarrow \bigwedge^{n-3} W \rightarrow \cdots \rightarrow \bigwedge^2 W \rightarrow W \rightarrow I_X \rightarrow 0$$

gives a resolution of I_X . Since $H^i(\mathbb{P}^n, (\bigwedge^j W)(m)) = 0$ for $0 < i < n$ and all m , we conclude that $h^i(\mathbb{P}^n, I_X(m)) = 0$ if $i = 1$ or 2 by an easy diagram chase. Finally, the long exact sequence associated to the standard sequence

$$0 \rightarrow I_X(m) \rightarrow \mathcal{O}_{\mathbb{P}^n}(m) \rightarrow \mathcal{O}_X(m) \rightarrow 0$$

shows that $h^1(X, \mathcal{O}_X(m)) = 0$ and the map $H^0(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(m)) \rightarrow H^0(X, \mathcal{O}_X(m))$ is surjective. \square

The bundles. By assumption, D_1 contains a linear space $\Phi \cong \mathbb{P}^{n-3}$. Let $\Lambda \cong \mathbb{P}^{n-2} \subset \mathbb{P}^n$ be a linear subspace of codimension 2 containing Φ . Since the singularities of D_1 occur in codimension 5, we must have $\Lambda \not\subset D_1$ and $\Lambda \cap D_1 = \Phi \cup Y$, where Y is a degree $d_1 - 1$ hypersurface in Λ . Furthermore, since D_2, \dots, D_{n-2} are very general hypersurfaces, $Z = Y \cap D_2 \cap \cdots \cap D_{n-2}$ is a zero-dimensional complete intersection scheme of length $(d_1 - 1)e$. Moreover, if Λ is general, then Z is reduced.

Proposition 3.2. *The scheme $Z \subset X$ satisfies the Cayley-Bacharach property for the line bundle $\mathcal{O}_X(d - n) \cong \omega_X(1)$.*

Proof. Observe that the scheme Z is a complete intersection of degrees $d_1 - 1, d_2, \dots, d_{n-2}, 1, 1$ in \mathbb{P}^n . Hence, by Theorem 2.2, the scheme Z satisfies the Cayley-Bacharach property for $\mathcal{O}_{\mathbb{P}^n}(d - n)$. By Proposition 3.1, every section of $\mathcal{O}_X(d - n)$ is a restriction of a section of $\mathcal{O}_{\mathbb{P}^n}(d - n)$. Hence, Z satisfies the Cayley-Bacharach property for $\mathcal{O}_X(d - n)$. \square

By the Serre correspondence, there is a locally free sheaf \mathcal{E} on X defined by a sequence

$$(*) \quad 0 \rightarrow \mathcal{O}_X \rightarrow \mathcal{E} \rightarrow I_Z(1) \rightarrow 0.$$

The following proposition summarizes the basic properties of \mathcal{E} .

Proposition 3.3. *Let \mathcal{E} be the locally free sheaf defined by (*). Then:*

- (1) *We have $\text{ext}^1(I_Z(1), \mathcal{O}_X) = 1$, so up to scalars there is a unique non-split extension \mathcal{E} .*
- (2) *The bundle \mathcal{E} is μ_H -stable and $\mathcal{E} \in M_X(2, H, (d_1 - 1)e)$.*
- (3) *Let $\delta_{i,j}$ denote the Kronecker delta function. Then $h^0(X, \mathcal{E}) = 3 + \delta_{2,d_1}$ and $h^1(X, \mathcal{E}) = 0$.*

Proof. Throughout the proof, let I_Z denote the ideal sheaf of Z in X . We will write $I_{Z/Y}$, such as I_{Z/\mathbb{P}^n} , when referring to the ideal sheaf of Z in an ambient space Y .

By Serre duality, we have

$$\text{ext}^1(I_Z(1), \mathcal{O}_X) = \text{ext}^1(\mathcal{O}_X, I_Z(K_X + H)) = h^1(X, I_Z(d - n)).$$

By the Cayley-Bacharach theorem, $h^1(\mathbb{P}^n, I_{Z/\mathbb{P}^n}(d - n)) = 1$. By Proposition 3.1,

$$H^1(\mathbb{P}^n, I_X(d - n)) = H^2(\mathbb{P}^n, I_X(d - n)) = 0.$$

The long exact sequence of cohomology associated to

$$0 \rightarrow I_X(d - n) \rightarrow I_{Z/\mathbb{P}^n}(d - n) \rightarrow I_Z(d - n) \rightarrow 0.$$

now implies that $h^1(X, I_Z(d - n)) = \text{ext}^1(I_Z(1), \mathcal{O}_X) = 1$, proving (1).

Since $\text{Pic}(X) \cong \mathbb{Z}H$, if \mathcal{E} is not μ_H -stable, then there exists a subbundle $\mathcal{O}_X(m) \subset \mathcal{E}$ with $m \geq 1$. The composition $\mathcal{O}_X(m) \rightarrow \mathcal{E} \rightarrow I_X(1)$ must be the zero map, so $\mathcal{O}_X(m)$ must factor through the map $\mathcal{O}_X \rightarrow \mathcal{E}$ in the sequence (*). This is impossible if $m \geq 1$, so \mathcal{E} must be μ_H -stable. A simple Chern class computation shows that $\mathcal{E} \in M_X(2, H, (d_1 - 1)e)$. This proves (2).

If $d_1 > 2$, then Z spans Λ . Hence, the hyperplanes containing Z are the hyperplanes containing Λ and $h^0(X, I_Z(1)) = 2$. Since $h^1(X, \mathcal{O}_X) = 0$, the long exact sequence associated to (*) shows that $h^0(X, \mathcal{E}) = 1 + h^0(X, I_Z(1)) = 3$. If $d_1 = 2$, then Z spans a linear space of dimension $n - 3$ and $h^0(X, I_Z(1)) = 3$. In this case, we get $h^0(X, \mathcal{E}) = 1 + h^0(X, I_Z(1)) = 4$.

In order to calculate $h^1(X, \mathcal{E})$, we first calculate $h^2(X, \mathcal{E}) = h^0(X, \mathcal{E}^*(d - n - 1))$. Since \mathcal{E} has rank 2, we have the isomorphism

$$\mathcal{E}^* \cong \mathcal{E} \otimes (\det \mathcal{E})^* = \mathcal{E}(-1).$$

Thus

$$h^2(X, \mathcal{E}) = h^0(X, \mathcal{E}(d - n - 2)).$$

We twist the sequence (*) by $\mathcal{O}_X(d - n - 2)$ to get

$$0 \rightarrow \mathcal{O}_X(d - n - 2) \rightarrow \mathcal{E}(d - n - 2) \rightarrow I_Z(d - n - 1) \rightarrow 0.$$

Since $h^1(X, \mathcal{O}_X(d - n - 2)) = 0$, we obtain

$$(1) \quad h^0(X, \mathcal{E}(d - n - 2)) = h^0(X, \mathcal{O}_X(d - n - 2)) + h^0(X, I_Z(d - n - 1)).$$

By Serre duality and the fact that the equations defining X have degree at least 2, we have

$$h^2(X, \mathcal{O}_X(d - n - 2)) = h^0(X, \mathcal{O}_X(1)) = n + 1.$$

Hence,

$$(2) \quad h^0(X, \mathcal{O}_X(d - n - 2)) = \chi(\mathcal{O}_X(d - n - 2)) - (n + 1).$$

Consider the ideal sheaf exact sequence

$$0 \rightarrow I_X(d - n - 1) \rightarrow I_{Z/\mathbb{P}^n}(d - n - 1) \rightarrow I_Z(d - n - 1) \rightarrow 0.$$

By Proposition 3.1, $h^1(\mathbb{P}^n, I_X(d - n - 1)) = 0$ and

$$(3) \quad h^0(X, I_Z(d - n - 1)) = h^0(\mathbb{P}^n, I_{Z/\mathbb{P}^n}(d - n - 1)) - h^0(\mathbb{P}^n, I_X(d - n - 1)).$$

The value of $h^0(\mathbb{P}^n, I_X(d - n - 1))$ is easily determined from the exact sequence

$$0 \rightarrow I_X(d - n - 1) \rightarrow \mathcal{O}_{\mathbb{P}^n}(d - n - 1) \rightarrow \mathcal{O}_X(d - n - 1) \rightarrow 0.$$

Since $\mathcal{O}_X(d - n - 1) = \omega_X$, we have $h^1(X, \omega_X) = 0$ and $h^2(X, \omega_X) = 1$. Furthermore, $h^3(\mathbb{P}^n, I_X(d - n - 1)) = 1$ and $h^i(\mathbb{P}^n, I_X(d - n - 1)) = 0$ for $i \neq 0, 3$. Hence,

$$(4) \quad h^0(\mathbb{P}^n, I_X(d - n - 1)) = \chi(I_X(d - n - 1)) + 1.$$

The scheme Z is a complete intersection of type $1, 1, (d_1 - 1), d_2, \dots, d_{n-2}$. The Koszul complex for $I_{Z/\mathbb{P}^n}(d - n - 1)$ implies that $h^1(\mathbb{P}^n, I_{Z/\mathbb{P}^n}(d - n - 1)) = n - 1 - \delta_{2, d_1}$. We conclude that

$$h^0(\mathbb{P}^n, I_{Z/\mathbb{P}^n}(d - n - 1)) = \chi(I_{Z/\mathbb{P}^n}(d - n - 1)) + n - 1 - \delta_{2, d_1}.$$

Combining this with Equations (1), (2), (3), and (4), we obtain

$$h^2(\mathcal{E}) = h^0(X, \mathcal{E}(d - n - 2)) = \chi(\mathcal{O}_X(d - n - 2)) + \chi(I_X(d - n - 1)) - 3 - \delta_{2, d_1}.$$

Since $\chi(\mathcal{E}(d - n - 2)) = \chi(\mathcal{O}_X(d - n - 2)) + \chi(I_X(d - n - 1))$ and $h^2(\mathcal{E}(d - n - 2)) = h^0(\mathcal{E}) = 3 + \delta_{2, d_1}$, we conclude that $h^1(X, \mathcal{E}) = h^1(\mathcal{E}(n - d - 2)) = 0$. This concludes the proof of the proposition. \square

We are now ready to state and prove our main theorem.

Theorem 3.4. *Let $n \geq 4$ and $d_1 > 2$. If $n = 4$, assume that $d_1 \geq 4$ and $d_2 \geq 6$. Then for every connected (respectively, irreducible) component of the Fano scheme $F_{n-3}(D_1)$, the moduli space $M_X(2, H, (d_1 - 1)e)$ has a connected (respectively, irreducible) component of the same dimension.*

Proof. Let $N \subset M_X(2, H, (d_1 - 1)e)$ be an irreducible component containing the bundle \mathcal{E} defined by (*). Endow N with its reduced induced structure. By Proposition 3.3, $h^1(X, \mathcal{E}) = 0$. Hence, by the upper semicontinuity of cohomology, $h^0(X, \mathcal{E}') \geq 3$ for every sheaf $\mathcal{E}' \in N$ and there is an exact sequence

$$0 \rightarrow \mathcal{O}_X \rightarrow \mathcal{E}' \rightarrow Q \rightarrow 0.$$

Computing Chern classes, we must have $\text{rk}(Q) = 1$, $c_1(Q) = H$, and $c_2(Q) = (1 - d_1)e$. We show that Q is torsion-free. Every element $\mathcal{E}' \in M_X(2, H, (d_1 - 1)e)$ is μ_H -stable with $c_1(\mathcal{E}') = H$. Hence, for any integer $m > 0$, we must have $h^0(X, \mathcal{E}'(-m)) = 0$. Since $h^1(X, \mathcal{O}_X(-m)) = 0$ by Proposition 3.1, we conclude that $h^0(X, Q(-m)) = 0$, so Q cannot have torsion in dimension zero.

If Q has torsion along a curve in X , let $T \subset Q$ be the torsion subsheaf and Q' be the quotient Q/T . Then Q' is a quotient of \mathcal{E}' , but $c_1(Q') = aH$ for some $a \leq 0$, which contradicts the stability of \mathcal{E}' . Thus Q is torsion-free, and consequently, $Q = I_{Z'}(1)$ for some scheme Z' of length $(d_1 - 1)e$.

The sequence

$$0 \rightarrow \mathcal{O}_X \rightarrow \mathcal{E}' \rightarrow I_{Z'}(1) \rightarrow 0$$

shows that $h^0(X, I_{Z'}(1)) \geq 2$. We next show that equality must hold. If

$$h^0(X, I_{Z'}(1)) > 2,$$

then Z' is contained in a linear space Γ of dimension at most $n - 3$. The length of Z' is $(d_1 - 1)e$, which is greater than the degree of $W = \cap_{i=2}^{n-2} D_i$. By Bézout's theorem, $W \cap \Gamma$ must contain a curve. Since D_i are very general, $d_i \geq 4$ and $\sum_{i=2}^{n-2} d_i > n + 1$, this is not possible by Proposition 2.4. We conclude that $h^0(X, I_{Z'}(1)) = 2$, $h^0(X, \mathcal{E}') = 3$ and the scheme Z' spans a linear space Λ' of dimension $n - 2$.

Since $\text{Pic}(X) \cong \mathbb{Z}H$, $\Lambda' \cap X$ must be a zero-dimensional complete intersection scheme of degree d_1e and cannot contain a curve. Since \mathcal{E}' is a nontrivial extension of $I_{Z'}(1)$ by \mathcal{O}_X , we have $h^1(X, I_{Z'}(d - n)) \geq 1$. The Cayley-Bacharach theorem implies that there exists a linear space $\Phi' \subset \Lambda'$ of dimension at most $n - 3$ containing the subscheme $Z'' \subset X \cap \Lambda'$ of length e residual to Z' . The dimension of Φ' must be $n - 3$. Otherwise, we could take the span of Φ' with additional points of $W = \cap_{i=2}^{n-2} D_i$ to obtain a linear space Δ of dimension $n - 3$ which would intersect W in more points than $e = \deg(W)$. Hence by Bezout's theorem, Δ would intersect W in a curve, contradicting Proposition 2.4. We conclude that $h^1(X, I_{Z'}(d - n)) = 1$ and $\dim(\Phi') = n - 3$.

The scheme Z'' of length e is contained in the intersection $\Phi' \cap W$. Since the two schemes have the same length, we conclude that $Z'' = \Phi' \cap W$. Since Z'' is a complete intersection scheme in Φ' of hypersurfaces of degree greater than d_1 , Z'' can only be contained in D_1 if $\Phi' \subset D_1$. Reversing the construction, we conclude that \mathcal{E}' is obtained by the same Serre construction as \mathcal{E} starting with $\Phi' \in F_{n-3}(D_1)$ instead of Φ . In particular, $H^1(X, \mathcal{E}') = 0$ and \mathcal{E}' is locally free.

Let \mathbb{E} be the pushforward of the universal sheaf over $X \times N$ to N . By the universal property of the Fano scheme, we get a morphism $\mathbb{P}\mathbb{E} \rightarrow F_{n-3}(D_1)$. In concrete terms, let (\mathcal{E}', s') be a sheaf together with a section up to scaling. Let Z' be the zero locus of s' and Λ' its span. The morphism sends (\mathcal{E}', s') to the span of the points residual to Z' in $\Lambda' \cap X$. Since \mathcal{E}' and \mathcal{E} are in the same irreducible component N , the $(n - 3)$ -plane Φ' must lie in the same irreducible component K of $F_{n-3}(D_1)$ as Φ . In particular, if $n \geq 5$, then Proposition 2.3 gives $\Phi = \Phi'$ and $\mathcal{E} = \mathcal{E}'$, and we conclude that N consists of a unique point. The same argument applies if $n = 4$ and $\dim(K) = 0$. If $n = 4$ and $\dim(K) = 1$, since there cannot be any nonconstant morphisms from \mathbb{P}^2 to a curve, we conclude that the linear space Φ' is canonically associated to \mathcal{E}' . Any irreducible component N' that intersects N must contain a bundle that arises via the Serre correspondence, hence must correspond to an irreducible component K' of $F_{n-3}(D_1)$ that intersects K . We thus conclude that the connected component containing \mathcal{E} corresponds to the connected component of $F_{n-3}(D_1)$ containing Φ and they have the same dimension. \square

Remark 3.5. If $n = 4$ or 5 and $d_1 = 2$, then $h^0(X, \mathcal{E}) = 4$ and $h^1(X, \mathcal{E}) = 0$. If \mathcal{E}' is in the same irreducible component N as \mathcal{E} , then $h^0(X, \mathcal{E}') \geq 4$. In this case, Z' spans at most a \mathbb{P}^{n-3} . If $n = 4$, since Z' has $e > 3$ points, it must span a line. If $n = 5$ and Z' was contained in a line, then the line would have to be contained in all the hypersurfaces defining X by Bézout's theorem. This would contradict the fact that $\text{Pic}(X) \cong \mathbb{Z}H$. We conclude that $h^0(X, \mathcal{E}') = 4$ for every bundle in N . The same argument shows that there is a connected component of $M_X(2, H, e)$ for every connected component of $F_{n-3}(D_1)$. In this case, the dimension count in the proof of Theorem 3.4 shows that the bundle corresponding to each connected component of $F_{n-3}(D_1)$ is unique.

Remark 3.6. Assume that $d_1 \geq 3$ when $n \geq 5$ and assume that $d_1 \geq 6$ when $n = 4$. Since the general D_1 does not contain a linear space of dimension $n - 3$, the proof of Theorem 3.4 shows that the bundles constructed on the special surfaces X do not deform to nearby surfaces.

4. EXAMPLES OF MODULI SPACES

In this section we give explicit examples of moduli spaces that exhibit interesting behaviors. We preserve the notation of Section 3.

Example 4.1 (Quintic threefold). Let D_1 be a general quintic threefold in \mathbb{P}^4 . Then D_1 contains 2875 lines. Let D_2 be a very general hypersurface of degree $d_2 \geq 6$. Let $X = D_1 \cap D_2$. By Theorem 3.4, $M_X(2, H, 4d_2)$ contains 2875 zero-dimensional connected components.

Example 4.2 (Moduli spaces with arbitrarily many isolated points I). Let $d_1 \geq 6$ and let D_1 be the hypersurface of degree d_1 defined by

$$f(z_0, \dots, z_4) = z_0^{d_1} - z_1^{d_1} + z_2^{d_1} - z_3^{d_1} + z_4 g(z_0, \dots, z_4) = 0$$

where g is a general homogeneous polynomial of degree $d_1 - 1$. Then D_1 is smooth and contains $3d_1^2$ lines contained in the Fermat surface $f = z_4 = 0$. Let ℓ be one of these lines. Up to permutations and roots of unity, we may assume that ℓ is defined by $z_0 - z_1 = z_2 - z_3 = z_4 = 0$. In the normal bundle exact sequence

$$0 \rightarrow N_{\ell/D_1} \rightarrow N_{\ell/\mathbb{P}^4} \cong \mathcal{O}_{\ell}(1)^{\oplus 3} \xrightarrow{M} \mathcal{O}_{\ell}(d_1) \rightarrow 0,$$

the map M is given by $[(d_1 - 1)s^{d_1-1}, (d_1 - 1)t^{d_1-1}, g(s, s, t, t, 0)]$ in local coordinates s, t on \mathbb{P}^1 [CR19, §3]. Hence, for a general choice g ,

$$N_{\ell/D_1} \cong \mathcal{O}_{\ell} \left(\left\lfloor \frac{3 - d_1}{2} \right\rfloor \right) \oplus \mathcal{O}_{\ell} \left(\left\lceil \frac{3 - d_1}{2} \right\rceil \right).$$

This implies $h^0(\ell, N_{\ell/D_1}) = 0$. We conclude that these $3d_1^2$ lines are isolated points of $F_1(D_1)$. Let $d_2 > d_1$ and let D_2 be a very general hypersurface in \mathbb{P}^4 of degree d_2 . Let $X = D_1 \cap D_2$. By Theorem 3.4, $M_X(2, H, (d_1 - 1)d_2)$ has at least $3d_1^2$ connected components of dimension 0.

Example 4.3 (Spinor bundles). Let D_1 be a smooth quadric fourfold in \mathbb{P}^5 . Let D_2 and D_3 be very general hypersurfaces of degrees $d_2, d_3 \geq 3$, respectively. Let $X = D_1 \cap D_2 \cap D_3$. Since $F_2(D_1)$ has two connected components, by Remark 3.5, $M_X(2, H, d_2d_3)$ has 2 connected components of dimension 0. One can be explicit about these two bundles: they are the restrictions of the two spinor bundles on the quadric D_1 . Identifying D_1 with the Grassmannian $G(2, 4)$ under the Plücker embedding, the spinor bundles are the universal quotient bundle Q and the dual S^* of the universal subbundle. These two bundles restrict to two bundles in $M_X(2, H, d_2d_3)$, which form two connected components. Since $h^0(G(2, 4), S^*) = h^0(G(2, 4), Q) = 4$ and the zero loci of the sections are planes, it is easy to see that the construction in §3 produces these two bundles.

Example 4.4 (Moduli spaces with arbitrarily many isolated points II). Let $d_1 \geq 3$ and let D_1 be the Fermat hypersurface in \mathbb{P}^5 defined by $\sum_{i=0}^5 x_i^{d_1} = 0$. Then D_1 contains $15d_1^3$ planes. Letting D_2, D_3

be very general hypersurfaces of degrees $d_2, d_3 > d_1$, Theorem 3.4 shows that $M_X(2, H, (d_1 - 1)d_2d_3)$ has at least $15d_1^3$ connected components.

Example 4.5 (Moduli spaces with arbitrarily many positive dimensional connected components). This example shows that there can be many positive dimensional connected components of $M_X(2, H, (d_1 - 1)e)$. Let $g(z_0, z_1)$ and $h(z_2, z_3, z_4)$ be general forms of degree $d_1 > 5$. Let $D_1 \subset \mathbb{P}^4$ be defined by the equation

$$f(z_0, \dots, z_4) = g(z_0, z_1) + h(z_2, z_3, z_4) = 0.$$

Let ℓ be the line defined by $z_2 = z_3 = z_4 = 0$ and let Λ be the plane defined by $z_0 = z_1 = 0$. Then ℓ intersects D_1 at the d_1 roots of $g(z_0, z_1)$. The tangent hyperplane at one of these points p_i intersects D_1 in the cone over the plane curve $z_0 = z_1 = h(z_2, z_3, z_4) = 0$ with vertex at p_i . Hence, D_1 contains d_1 disjoint one-parameter families of lines.

We claim that D_1 contains no other lines. Let $U \subset \mathbb{G}(1, 4)$ denote the open subset of lines not meeting ℓ . Consider the incidence correspondence

$$I = \{(g, h, m) \in H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(d_1)) \times H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(d_1)) \times U : (g(z_0, z_1) + h(z_2, z_3, z_4))|_m \equiv 0\}.$$

The PGL_5 action on U has two orbits: the lines that intersect Λ and the lines that do not. We can compute the fiber dimension of I over U by choosing the following representatives of the two orbits

$$m_1 : z_0 - z_1 = z_0 - z_3 = z_4 = 0 \quad \text{and} \quad m_2 : z_0 - z_2 = z_1 - z_3 = z_4 = 0.$$

Let $g = \sum_{i=0}^{d_1} a_i z_0^i z_1^{d_1-i}$ and let $h = \sum_{i,j} b_{i,j} z_2^i z_3^j z_4^{d_1-i-j}$. If $g+h$ vanishes on m_1 , then the coefficients of g and h must satisfy the $d_1 + 1$ linear conditions $b_{i,d_1-i} = 0$ for $1 \leq i \leq d_1$ and $b_{0,d_1} + \sum_{i=0}^{d_1} a_i = 0$. If $g+h$ vanishes on m_2 , then the coefficients of g and h must satisfy the $d_1 + 1$ linear conditions $a_i + b_{i,d_1-i} = 0$ for $0 \leq i \leq d_1$. Since $d_1 + 1 > 6 = \dim U$, we conclude that

$$\dim I < \dim H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(d_1)) \times H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(d_1)).$$

Hence, for a general choice of g and h , D_1 does not contain a line of U . Hence, any line in D_1 intersects ℓ and must pass through one of the points p_i . Since the tangent hyperplane at p_i intersects D_1 in a cone with vertex at p_i , the line must be one of the rulings of the cone.

Let $D_2 \subset \mathbb{P}^4$ be a very general hypersurface of degree $d_2 > d_1$ and let $X = D_1 \cap D_2$. By Theorem 3.4, $M_X(2, H, (d_1 - 1)d_2)$ has at least d_1 one-dimensional connected components.

REFERENCES

- [Ar07] E. Arrondo, A home-made Hartshorne-Serre correspondence, *Rev. Mat. Complut.* **20** no. 2 (2007), 423–443.
- [CH18] I. Coskun and J. Huizenga, The moduli spaces of sheaves on surfaces, pathologies, and Brill-Noether problems, in J. Christophersen and K. Ranestad (Eds.) *Geometry of Moduli* (2018), 75–105, Abel Symposia.
- [CR19] I. Coskun and E. Riedl, Normal bundles of rational curves on complete intersections, *Commun. Contemp. Math.* **21** no. 2 (2019).
- [CW20] I. Coskun and M. Woolf, The stable cohomology of moduli spaces of sheaves on surfaces, to appear in *J. Differential Geom.*
- [Do90] S. K. Donaldson, Polynomial invariants for smooth four-manifolds, *Topology*, **29** (1990), no. 3, 257–315.
- [Ei88] L. Ein, Generalized null correlation bundles, *Nagoya Math. J.*, **111** (1988), 13–24.
- [EGH] D. Eisenbud, M. Green, and J. Harris, Cayley-Bacharach theorems and conjectures, *Bull. Amer. Math. Soc.*, **33** no. 3 (1996), 295–324.
- [Fo68] J. Fogarty, Algebraic families on an algebraic surface. *Amer. J. Math.*, **90** (1968), 511–521.
- [Fr89] R. Friedman, Rank two vector bundles over regular elliptic surfaces, *Invent. Math.*, **96** (1989), 283–332.
- [FM88] R. Friedman and J. W. Morgan, On the diffeomorphism types of certain algebraic surfaces. I., *J. Differential Geom.*, **27** (1988), 297–369.
- [Gi77] D. Gieseker, On the moduli space of vector bundles on an algebraic surface, *Ann. Math.*, **106** (1977), 45–60.
- [GL94] D. Gieseker and J. Li, Irreducibility of moduli of rank 2 bundles on algebraic surfaces, *J. Diff. Geom.*, **40** (1994), 23–104.

- [Go90] L. Göttsche, The Betti numbers of the Hilbert scheme of points on a smooth projective surface, *Math. Ann.*, **286** no. 3 (1990), 193–207.
- [HuL] D. Huybrechts and M. Lehn, *The Geometry of Moduli Spaces of Sheaves*, Second edition, Cambridge Mathematical Library, Cambridge Univ. Press, Cambridge, 2010.
- [Ia72] A. Iarrobino, Reducibility of the families of 0-dimensional schemes on a variety, *Invent. Math.*, **15** (1972), 72–77.
- [Ko89] D. Kotschick, On manifolds diffeomorphic to $\mathbb{C}P^2 \# \overline{\mathbb{C}P^2}$, *Invent. Math.*, **95** no. 3 (1989), 591–600.
- [Ma78] M. Maruyama, Moduli of stable sheaves II, *J. Math. Kyoto*, **18** (1978), 557–614.
- [Mes97] N. Mestrano, Sur le espaces de modules de fibrés vectoriels de rang deux sur des hypersurfaces de \mathbb{P}^3 , *J. reine angew. Math.* **490** (1997), 65–79.
- [MS11] N. Mestrano and C. Simpson, Obstructed bundles of rank two on a quintic surface, *Int. J. Math.*, **22** (2011), 789–836.
- [MS16] N. Mestrano and C. Simpson, Moduli of sheaves, *Development of Moduli Theory–Kyoto 2013*, *Adv. Stud. Pure Math.* **69** (2016), 77–172, 2016.
- [Mo67] B. G. Moishezon, Algebraic homology classes on algebraic varieties, *Izv. Akad. Nauk. SSSR Ser. Mat.*, **31** no. 2 (1967), 225–268.
- [Ni95] P. Nijssse, The irreducibility of the moduli space of stable vector bundles of rank 2 on a quintic in \mathbb{P}^3 , Preprint arXiv:alg-geom/9503012 (1995).
- [O’G96] K. G. O’Grady, Moduli of vector bundles on projective surfaces: Some basic results, *Invent. Math.*, **123** (1996), 141–207.
- [OV86] C. Okonek and A. Van de Ven, Stable bundles and differentiable structures on certain elliptic surfaces, *Invent Math.*, **86** no. 2 (1986), 357–370.
- [St19] J. Starr, Connectedness of moduli space, URL (2019) mathoverflow.net/q/323673.
- [Ts20] D. Tseng, Collections of Hypersurfaces Containing a Curve, *Int. Math. Res. Not.*, Vol. 2020, No. 13, 3927–3977.
- [Va06] R. Vakil, Murphy’s Law in algebraic geometry: Badly-behaved deformation spaces, *Invent. Math.*, **164** (2006), 569–590.

DEPARTMENT OF MATHEMATICS, STATISTICS AND CS, UNIVERSITY OF ILLINOIS AT CHICAGO, CHICAGO, IL 60607
Email address: `icoskun@uic.edu`

DEPARTMENT OF MATHEMATICS, THE PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK, PA 16802
Email address: `huizenga@psu.edu`

DEPARTMENT OF MATHEMATICS, THE PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK, PA 16802
Email address: `kopper@psu.edu`