

MATH 553: MIDTERM PROBLEM SET

This problem set is due Friday February 28, 2020. You may work on the problem set in groups; however, the final write-up must be yours and reflect your own understanding.

Problem 1. Let $X \subset \mathbb{P}_k^n$ be a nonsingular hypersurface of degree d , where k is a field. Let $\mathcal{O}_X(e)$ denote the pullback of $\mathcal{O}_{\mathbb{P}^n}(e)$ under the inclusion i of X in \mathbb{P}_k^n .

- (1) For all $e \in \mathbb{Z}$ and all $i \geq 0$ compute $H^i(X, \mathcal{O}_X(e))$.
- (2) Show that $\omega_X \cong \mathcal{O}_X(d - n - 1)$ and compute $H^i(X, \omega_X)$.

Problem 2. Do Exercises III.5.1 and III.5.2 in Hartshorne. Compute the Hilbert polynomials of the following projective varieties using the definition in III.5.2:

- (1) A rational normal curve of degree d .
- (2) The Veronese image of \mathbb{P}^n under the d -uple Veronese embedding.
- (3) A hypersurface of degree d in \mathbb{P}^n .

Problem 3. Let k be a field. Prove the following theorem by following the given steps.

Theorem Any vector bundle V of rank r on \mathbb{P}_k^1 is isomorphic to a direct sum of line bundles $\bigoplus_{i=1}^r \mathcal{O}_{\mathbb{P}^1}(a_i)$ for a uniquely determined sequence of integers $a_1 \leq a_2 \leq \dots \leq a_r$.

- (1) Show the uniqueness.
- (2) Show that there exists a nonzero homomorphism $\mathcal{O}_{\mathbb{P}^1}(a) \rightarrow V$ for some $a \in \mathbb{Z}$.
- (3) Let

$$a_r := \max\{a \mid \text{Hom}(\mathcal{O}_{\mathbb{P}^1}(a), V) \neq 0\}.$$

Show that in the exact sequence induced by a nonzero homomorphism

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^1}(a_r) \rightarrow V \rightarrow W \rightarrow 0,$$

W is locally free. Hence, by induction on the rank we may assume that the theorem holds for W and $W = \bigoplus_{i=1}^{r-1} \mathcal{O}_{\mathbb{P}^1}(b_i)$.

- (4) Do exercise III.6.1 in Hartshorne.
- (5) Prove that $\text{Ext}^1(W, \mathcal{O}_{\mathbb{P}^1}(a_r)) = 0$ and conclude the theorem by induction. (Hint: Tensor the sequence defining W by $\mathcal{O}_{\mathbb{P}^1}(-a_r - 1)$ and compute cohomology.)

Problem 4. Let $n \geq 2$ and let $T\mathbb{P}_k^n$ denote the tangent bundle of \mathbb{P}_k^n .

- (1) Prove that if $\text{Hom}(\mathcal{O}_{\mathbb{P}^n}(a), T\mathbb{P}^n) \neq 0$, then $a \leq 1$.
- (2) Show that $T\mathbb{P}^n$ for $n \geq 2$ is not a direct sum of line bundles. Hence, the theorem in the previous problem does not generalize to \mathbb{P}^n for $n > 1$.
- (3) Show that $T\mathbb{P}^n(-i)$ is acyclic (has no cohomology) for $i = 2, 3, \dots, n$ and $i = n + 2$.

Problem 5. Let k be an algebraically closed field.

- (1) Let I_Z be the ideal sheaf of 4 points on \mathbb{P}_k^2 . Show that $H^1(\mathbb{P}^2, I_Z(2)) = 0$ if and only if the 4 points are not collinear.
- (2) Let I_Z be the ideal sheaf of 9 points which are the complete intersection of 2 cubic curves. Let W be any 8 of the points in Z . Show that $H^1(\mathbb{P}^2, I_W(3)) = 0$ and $H^1(\mathbb{P}^2, I_Z(3)) = k$.

Problem 6. Let X be a smooth, integral, projective surface over an algebraically closed field k . Let Z be a zero-dimensional scheme of length n on X and let I_Z denote its ideal sheaf. Show that $\dim_k(\text{Hom}_{\mathcal{O}_X}(I_Z, \mathcal{O}_Z)) = 2n$ by following the given steps:

- (1) Observe that $\text{Ext}^i(\mathcal{O}_X, \mathcal{O}_Z) = 0$ for $i > 0$.
 (2) Compute $\text{Hom}(\mathcal{O}_Z, \mathcal{O}_Z)$ and $\text{Ext}^2(\mathcal{O}_Z, \mathcal{O}_Z)$. Then use a locally free resolution of \mathcal{O}_Z to compute

$$\chi(\mathcal{O}_Z, \mathcal{O}_Z) = \sum_{i=0}^2 (-1)^i \dim(\text{Ext}^i(\mathcal{O}_Z, \mathcal{O}_Z))$$

directly and conclude that $\dim_k(\text{Ext}^1(\mathcal{O}_Z, \mathcal{O}_Z)) = 2n$.

- (3) Apply $\text{Hom}(-, \mathcal{O}_Z)$ to the standard exact sequence

$$0 \rightarrow I_Z \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_Z \rightarrow 0$$

to finish the computation.

Remark: Combined with the connectedness of the Hilbert scheme of points on a smooth, projective, integral surface X , this exercise shows that the Hilbert scheme of points on X is smooth.

Problem 7. A sheaf \mathcal{F} on a projective scheme X over an algebraically closed field k is called *exceptional* if $\text{Hom}(\mathcal{F}, \mathcal{F}) = k$ and $\text{Ext}^i(\mathcal{F}, \mathcal{F}) = 0$ for $i > 0$. A *strong exceptional collection* of sheaves is an ordered set of exceptional sheaves $\mathcal{F}_1, \dots, \mathcal{F}_r$ such that $\text{Ext}^i(\mathcal{F}_k, \mathcal{F}_l) = 0$ unless $k \leq l$ and $i = 0$.

- (1) Show that $\mathcal{O}_{\mathbb{P}^n}, \mathcal{O}_{\mathbb{P}^n}(1), \mathcal{O}_{\mathbb{P}^n}(2), \dots, \mathcal{O}_{\mathbb{P}^n}(n)$ is a strong exceptional collection on \mathbb{P}^n .
 (2) Show that $\mathcal{O}_{\mathbb{P}^2}, T\mathbb{P}^2(-1), \mathcal{O}_{\mathbb{P}^2}(1)$ is a strong exceptional collection on \mathbb{P}^2 .
 (3) Let X be the blowup of \mathbb{P}^2 at r distinct points p_1, \dots, p_r and let E_i denote the exceptional divisor over p_i . Let $\mathcal{O}_X(H)$ denote the pullback of $\mathcal{O}_{\mathbb{P}^2}(1)$ to X . Show that

$$\mathcal{O}_X(-2H), \mathcal{O}_X(-H), \mathcal{O}_X(-E_1), \mathcal{O}_X(-E_2), \dots, \mathcal{O}_X(-E_r), \mathcal{O}_X$$

is a strong exceptional collection on X . (Hint: Recall that $\mathcal{O}_{E_i}(E_i) \cong \mathcal{O}_{\mathbb{P}^1}(-1)$.)