

# PSti1l

A PostScript to PDF converter, written by Frank M. Siegert

PSti1l is NOT in the public domain. It is shareware.

This page will disappear in the registered PSti1l, to get a license key please send email to frank@this.net or refer to the registration.pdf file enclosed in the PSti1l archive.

Please register this software, otherwise further development is not possible.

This PDF file was generated by an unregistered PSti1l 1.10  
PSti1l is copyright Frank M. Siegert 1997, [frank@this.net]  
Please visit the PSti1l web page:

<http://www.this.net/~frank/psti1l.html>

PSti1l user: Floyd B. Hanson floyd@floyd2.[no getdomainname]  
PostScript, PDF, and others are Registered Trademarks of Adobe, Inc.  
and are used for identification purpose only. Other Trademarks belong  
to their respective owners and are used for the same purpose only.

**Directions:** Answer **All Questions** and show **All Work** in the **Exam Booklet** provided. Write your **Name**, **Social Security Number**, and **Discussion Section Hour/Day** on the Exam Book Cover Page. **Keep your eyes on your own work and keep your own work covered.** See *Table of Laplace Transforms* on Back.

1. (a)  $\mathcal{L} [7e^{-8t} \cos(3t) + t^5 + 11t^2 \delta(t - 3)]$

(b)  $\mathcal{L}^{-1} \left[ \frac{e^{-2s}}{s^2 + 12s + 32} \right]$  (25 points total)

(c)  $\mathcal{L}^{-1} \left[ \frac{s}{s^2 + 6s + 13} \right]$

2. (a) Find the general solution  $y(x)$ :

$$4x^2 y''(x) + 8xy'(x) + y(x) = 0, \quad x > 0.$$

(b) Find  $z(t)$ : (25 points total)

$$z''(t) + 3z'(t) + 2z(t) = 4\delta(t - \epsilon), \quad z(0) = 0, \quad z'(0) = 5, \quad \epsilon > 0.$$

What happens as  $\epsilon \rightarrow 0^+$ ?

3. (a) Find a power series approximation to the solution  $y(x)$  of the IVP:

$$y''(x) + 2xy'(x) - 4y(x) = 0, \quad y(0) = 2, \quad y'(0) = 1,$$

by assuming an expansion about  $x = 0$  of the form  $y(x) \simeq a_0 + a_1 x + a_2 x^2 + a_3 x^3 + a_4 x^4 + a_5 x^5$ , finding the coefficients  $\{a_0, a_1, a_2, a_3, a_4, a_5\}$ . You do not need any more terms!

(b) Classify all finite non-negative points of the ODE: (25 points total)

$$x^2 y''(x) + xy'(x) + (x^2 - 1/9)y(x) = 0, \quad x > 0$$

and then give only the general form of the power series solution in the regular case (say about  $x = a$ ) and the singular case, but do not evaluate any coefficients.

4. Solve the integral equation for  $y(t)$ : (25 points total)

$$y(t) + 8 \int_0^t y(v) \sin(t - v) dv = 9, \quad t \geq 0.$$

## Table of Laplace Transforms<sup>1</sup>

$f(t) = \mathcal{L}^{-1}(s)](t)$	$F(s) = \mathcal{L}[f(t)](s)$
$f(at)$	$\frac{1}{a}F\left(\frac{1}{a}\right)$
$e^{at}f(t)$	$F(s - a)$
$f^{(n)}(t)$	$s^n F(s) - s^{n-1}f(0) - \dots - f^{(n-1)}(0)$
$t^n f(t)$	$(-1)^n F^{(n)}(s)$
$\frac{1}{t}f(t), f(0) = 0$	$\int_s^\infty F(u)du$
$\int_0^t f(v)dv$	$F(s)/s$
$(f * g)(t)$	$F(s) \cdot G(s)$
$f(t - a)u(t - a), a \geq 0$	$e^{-as}F(s)$
$g(t)u(t - a), a \geq 0$	$e^{-as}\mathcal{L}[g(t + a)](s)$
$e^{at} \sin(bt)$	$\frac{b}{(s - a)^2 + b^2}$
$e^{at} \cos(bt)$	$\frac{s - a}{(s - a)^2 + b^2}$
$\sinh(bt)$	$\frac{b}{s^2 - b^2}$
$\cosh(bt)$	$\frac{s}{s^2 - b^2}$

---

<sup>1</sup>From Back Cover of Nagle and Saff Text