

INDEX THEORY AND NON-COMMUTATIVE GEOMETRY

I. HIGHER FAMILIES INDEX THEORY

MOULAY-TAHAR BENAMEUR AND JAMES L. HEITSCH

ABSTRACT. We prove an index theorem for foliated manifolds. We do so by constructing a push forward map in cohomology for a K -oriented map from an arbitrary manifold to the space of leaves of an oriented foliation, and by constructing a Chern-Connes character from the K -theory of the compactly supported smooth functions on the holonomy groupoid of the foliation to the Haefliger cohomology of the foliation. Combining these with the Connes-Skandalis topological index map and the classical Chern character gives a commutative diagram from which the index theorem follows immediately.

CONTENTS

1. Introduction	1
2. Notation and Review	3
3. Traces	4
4. The algebraic Chern character	8
5. Push forward maps	10
6. The index theorem	19
References	22

1. INTRODUCTION

This is the first in a series of papers on the different approaches to the higher index and Lefschetz theorems for foliations. In this paper, we give an alternative proof of the cohomological index theorem for families of elliptic operators defined along the leaves of a foliation of a compact manifold obtained in [HL99]. We do so by constructing a push forward map in cohomology for a K -oriented map from an arbitrary manifold to the space of leaves of the foliation, and by constructing a Chern-Connes character from the K -theory of the compactly supported smooth functions on the holonomy groupoid of the foliation to the Haefliger cohomology of the foliation. Combining these with the Connes-Skandalis topological index map and the classical Chern character gives a commutative diagram completely analogous to the commutative diagram of classical families index theory. As in the classical case, this immediately gives the cohomological index theorem obtained in [HL99].

In [CS84] and [Con81], Connes and Skandalis proved a families index theorem for foliations. Their theorem and its proof are algebraic in nature, showing that two push forward maps in K -theory are the same. Our constructions are more geometric and we get computable invariants in Haefliger cohomology. In addition, these invariants are more easily related to the geometry and topology of the foliation. The algebraic approach to invariants for non-commutative spaces is in some sense more fully developed than the geometric. It is our contention that the further development of the geometric approach will lend deep insight into these invariants. We expect that the application of the results here will give a much fuller understanding of the relationship of the index theory of leafwise operators with the geometry and topology of the foliation. Such insight is also likely to have applications to the Novikov conjecture, one of the central problems in topology.

This version of the paper incorporates its Corrigendum, [BH05].

In [A75], Atiyah identified the index of an elliptic differential operator D on a compact manifold with the trace in an associated von Neumann algebra of a primary K -theory invariant of the operator. This latter fundamental invariant lives in the K -theory of the C^* -algebra associated with the fundamental group of M and is a generalized index of the lifted operator $\tilde{D}$ of D to the universal cover $\tilde{M}$. The invariance of $\tilde{D}$ under the deck transformations insures that this deeper index class is well defined even though $\tilde{M}$ is non-compact in general. This construction led to many deep results in the topological and harmonic analysis of discrete groups, see for instance [BC84, BCH90, CM91, HK99, K88]. Atiyah's covering index theorem was generalized by Connes and Moscovici to higher traces in [CM91], giving a cyclic cohomology proof of the Novikov conjecture for hyperbolic groups.

In [Con79, Con81], Connes extended this construction to leafwise elliptic operators on compact foliated manifolds which are endowed with a holonomy invariant measure. To do so he replaced the lifting and deck transformations by a lifting to the holonomy covers of the leaves invariant under the natural action of the holonomy groupoid. Moreover, he defined an analytic index map from the K -theory of the tangent bundle to the foliation to the K -theory of the C^* algebra of the foliation, which plays the role of the K -theory of the space of leaves. In [CS84], Connes and Skandalis defined a push forward map in K -theory for any K -oriented map from a manifold to the space of leaves of a foliation of a compact manifold. This allowed them to define a topological index map from the K -theory of the tangent bundle of the foliation to the K -theory of its C^* algebra. Their main result is that the analytic and topological index maps are equal, an extension of the classical Atiyah-Singer families index theorem.

It is well known that the Atiyah-Singer families index theorem follows immediately from the commutative diagram

$$\begin{array}{ccc} K_c^0(N) & \xrightarrow{f_!} & K^0(M) \\ ch(\cdot) \wedge Td(f) \downarrow & & \downarrow ch \\ H_c^*(N; \mathbb{R}) & \xrightarrow{f_*} & H^*(M; \mathbb{R}). \end{array}$$

Here N and M are manifolds, with M compact. ch is the usual Chern character, and $f : N \rightarrow M$ is a K -oriented map with associated push forward map $f_!$. $Td(f)$ is the Todd class of f , and f_* is the push-forward map in cohomology defined using $f_* : H_*(N; \mathbb{R}) \rightarrow H_*(M; \mathbb{R})$ and Poincaré Duality.

We extend this result to foliations. Let F be an oriented foliation of M and $f : N \rightarrow M/F$ a K -oriented map to the space of leaves of F . Denote by $H_c^*(M/F)$ the Haefliger cohomology of F . We construct a push forward map in cohomology $f_* : H_c^*(N; \mathbb{R}) \rightarrow H_c^*(M/F)$. We also construct a Chern character $ch_a : K_0(C_c^\infty(\mathcal{G})) \rightarrow H_c^*(M/F)$, where $C_c^\infty(\mathcal{G})$ is the space of smooth functions with compact support on $\mathcal{G}$, the holonomy groupoid of F . Denote by $\hat{F}$ the foliation of $M \times \mathbb{R}^{2k}$ induced by F with the same leaf dimension. Its holonomy groupoid is $\mathcal{G}^{\mathbb{R}^{2k}} \simeq \mathcal{G} \times \mathbb{R}^{2k}$ and its leaf space is $M \times \mathbb{R}^{2k} / \hat{F} \simeq M/F \times \mathbb{R}^{2k}$. Composing $ch_a : K_0(C_0^\infty(\mathcal{G}^{\mathbb{R}^{2k}})) \rightarrow H_c^*(M \times \mathbb{R}^{2k} / \hat{F})$ with integration over $\mathbb{R}^{2k}$, $\int_{\mathbb{R}^{2k}} : H_c^*(M \times \mathbb{R}^{2k} / \hat{F}) \rightarrow H_c^*(M/F)$, gives the map $ch_a^{\mathbb{R}^{2k}} : K_0(C_c^\infty(\mathcal{G}^{\mathbb{R}^{2k}})) \rightarrow H_c^*(M/F)$. Given an imbedding of N in $\mathbb{R}^{2k}$, there is a Connes-Skandalis push forward map $f_!^{\mathbb{R}^{2k}} : K_c^0(N) \rightarrow K_0(C_c^\infty(\mathcal{G}^{\mathbb{R}^{2k}}))$. We are forced to work with this latter space because in general there is no Bott isomorphism between $K_0(C_c^\infty(\mathcal{G}^{\mathbb{R}^{2k}}))$ and $K_0(C_c^\infty(\mathcal{G}))$. Our main result is a complete extension of the above result to foliations.

Theorem 5.11. The following diagram commutes,

$$\begin{array}{ccc} K_c^0(N) & \xrightarrow{f_!^{\mathbb{R}^{2k}}} & K_0(C_c^\infty(\mathcal{G}^{\mathbb{R}^{2k}})) \\ ch(\cdot) \wedge Td(f) \downarrow & & \downarrow ch_a^{\mathbb{R}^{2k}} \\ H_c^*(N; \mathbb{R}) & \xrightarrow{f_*} & H_c^*(M/F). \end{array}$$

This result extends immediately to other interesting algebras, e.g. the space of superexponentially decaying operators in Connes' C^* -algebra, [HL02].

Denote by TF the tangent bundle of F , and by $\pi : TF \rightarrow M \rightarrow M/F$ the natural map. Then for k sufficiently large, there is a Connes-Skandalis push forward map $\pi_! : K_c^0(TF) \rightarrow K_0(C_c^\infty(\mathcal{G}^{\mathbb{R}^{2k}}))$. By standard methods, Theorem 5.11 immediately yields a cohomological index theorem, namely

Theorem 6.2. For any $u \in K_c^0(TF)$, $\text{ch}_a^{\mathbb{R}^{2k}} \circ \pi_!(u)$, the algebraic Chern character of the topological index of u , is given by

$$\text{ch}_a^{\mathbb{R}^{2k}} \circ \pi_!(u) = (-1)^p \int_F \pi_{F!}(\text{ch}(u)) \text{Td}(TF \otimes \mathbb{C}) \in H_c^*(M/F),$$

where $\int_F : H^*(M; \mathbb{R}) \rightarrow H_c^*(M/F)$ is integration over the leaves of F , $\pi_{F!} : H_c^*(TF; \mathbb{R}) \rightarrow H^*(M; \mathbb{R})$ is integration along the fibers, and Td is the Todd class.

Finally, in [Con86], A. Connes showed that the pairing of any holonomy invariant current extends to the K -theory of the C^* -algebra of the foliation. Therefore, our theorem 6.2 proves, using the Connes-Skandalis index theorem in K -theory [CS84] a cohomological index formula in Haefliger cohomology. In particular, our results partially answer a conjecture stated in [He95], namely that the index formula in Haefliger cohomology, for all elliptic pseudodifferential operators on foliations, compare [He95, HL99], is a consequence of the Connes-Skandalis index theorem.

Note that a similar index formula was obtained independently, for Dirac operators on foliations with Hausdorff graph, by Gorokhovsky-Lott, [GL03, GL05].

Acknowledgments. We are indebted to J.-M. Bismut, A. Connes, M. Crainic, T. Fack, J. Lott, I. Moerdijk, V. Nistor, H. Oyono-Oyono and G. Skandalis for many useful discussions.

Part of this work was done while the first author was visiting the University of Illinois at Chicago, and while the second author was visiting the Erwin Schrödinger Institute in Vienna, and the University of Lyon I. Both authors are most grateful for the warm hospitality and generous support of their hosts.

2. NOTATION AND REVIEW

Throughout the paper, M denotes a smooth compact Riemannian manifold of dimension n , and F denotes an oriented foliation of M of dimension p and codimension q . So $n = p + q$. The tangent and cotangent bundles of F will be denoted TF and T^*F respectively. If $E \rightarrow N$ is a vector bundle over a manifold N , we denote the space of smooth sections by $C^\infty(E)$ or by $C^\infty(N; E)$ if we want to emphasize the base space of the bundle. The compactly supported sections are denoted by $C_c^\infty(E)$ or $C_c^\infty(N; E)$. The space of differential k -forms on N is denoted $\mathcal{A}^k(N)$, and we set $\mathcal{A}(N) = \bigoplus_{k \geq 0} \mathcal{A}^k(N)$. The space of compactly supported k -forms is denoted $\mathcal{A}_c^k(N)$, and $\mathcal{A}_c(N) = \bigoplus_{k \geq 0} \mathcal{A}_c^k(N)$. The tangent and cotangent bundles of N are denoted TN and T^*N respectively.

The holonomy groupoid $\mathcal{G}$ of F consists of equivalence classes of paths $\gamma : [0, 1] \rightarrow M$ such that the image of γ is contained in a leaf of F . Two such paths γ_1 and γ_2 are equivalent if $\gamma_1(0) = \gamma_2(0)$, $\gamma_1(1) = \gamma_2(1)$, and the holonomy germ along them is the same. Two classes may be composed if one ends where the second begins and the composition is just the juxtaposition of the two paths. This makes $\mathcal{G}$ a groupoid. The space $\mathcal{G}^{(0)}$ of units of $\mathcal{G}$ consists of the equivalence classes of the constant paths, and we identify $\mathcal{G}^{(0)}$ with M .

$\mathcal{G}$ is a (in general non-Hausdorff) dimension $2p + q$ manifold. The basic open sets defining its manifold structure are given as follows. Let $\mathcal{U}$ be a finite good cover of M by foliation charts, [HL90]. Given U and V in this cover and a leafwise path γ starting in U and ending in V , define (U, γ, V) to be the set of equivalence classes of leafwise paths starting in U and ending in V which are homotopic to γ through a homotopy of leafwise paths whose end points remain in U and V respectively. It is easy to see, using the holonomy defined by γ from a transversal in U to a transversal in V , that if $U, V \simeq \mathbb{R}^p \times \mathbb{R}^q$, then we have the local diffeomorphism $(U, \gamma, V) \simeq \mathbb{R}^p \times \mathbb{R}^p \times \mathbb{R}^q$.

If $\mathcal{G}$ is non-Hausdorff, it is not true that compact sets are always closed, nor that the closure of a compact set is compact. Because of this, we define the notion of having compact support as follows. Given a bundle E over $\mathcal{G}$ and any set (U, γ, V) as above, consider $E|_{(U, \gamma, V)}$, the restriction of E to (U, γ, V) .

The space $C_c^\infty(E| (U, \gamma, V))$ has a natural inclusion into the space of sections of E over $\mathcal{G}$ by extending any element of $C_c^\infty(E| (U, \gamma, V))$ to all of $\mathcal{G}$ by defining it to be zero outside (U, γ, V) . We define the space $C_c^\infty(E) = C_c^\infty(\mathcal{G}; E)$ of smooth sections of E over $\mathcal{G}$ with compact support to be all finite sums $\sum s_i$ where each s_i is the image of an element $\hat{s}_i$ in some $C_c^\infty(E| (U, \gamma, V))$. The space of smooth functions with compact support, namely $C_c^\infty(\mathcal{G}; \mathbb{R})$, will be denoted $C_c^\infty(\mathcal{G})$.

The source and range maps of the groupoid $\mathcal{G}$ are the two natural maps $s, r : \mathcal{G} \rightarrow M$ given by $s([\gamma]) = \gamma(0)$, $r([\gamma]) = \gamma(1)$. $\mathcal{G}$ has two natural transverse foliations F_s and F_r whose leaves are respectively $\tilde{L}_x = s^{-1}(x)$, $\tilde{L}^x = r^{-1}(x)$ for $x \in M$. Note that $r : \tilde{L}_x \rightarrow L$ is the holonomy covering of L .

The Haefliger cohomology of F , [H80], is given as follows. For each $U_i \in \mathcal{U}$, let $T_i \subset U_i$ be a transversal and set $T = \bigcup T_i$. We may assume that the closures of the T_i are disjoint. Let $\mathcal{H}$ be the holonomy pseudogroup induced by F on T . Denote by $\mathcal{A}_c^k(M/F)$ the quotient of $\mathcal{A}_c^k(T)$ by the vector subspace generated by elements of the form $\alpha - h^*\alpha$ where $h \in \mathcal{H}$ and $\alpha \in \mathcal{A}_c^k(T)$ has support contained in the range of h . Give $\mathcal{A}_c^k(M/F)$ the quotient topology of the usual C^∞ topology on $\mathcal{A}_c^k(T)$, so this is not a Hausdorff space in general. The exterior derivative $d_T : \mathcal{A}_c^k(T) \rightarrow \mathcal{A}_c^{k+1}(T)$ induces a continuous differential $d_H : \mathcal{A}_c^k(M/F) \rightarrow \mathcal{A}_c^{k+1}(M/F)$. Note that $\mathcal{A}_c^k(M/F)$ and d_H are independent of the choice of cover $\mathcal{U}$. The complex $\{\mathcal{A}_c(M/F), d_H\}$ and its cohomology $H_c^*(M/F)$ are, respectively, the Haefliger forms and Haefliger cohomology of F .

As the bundle TF is oriented, there is a continuous open surjective linear map, called integration over the leaves,

$$\int_F : \mathcal{A}^{p+k}(M) \longrightarrow \mathcal{A}_c^k(M/F)$$

which commutes with the exterior derivatives d_M and d_H . Given $\omega \in \mathcal{A}(M)$, write $\omega = \sum_i \omega_i$ where $\omega_i \in \mathcal{A}_c(U_i)$. Integrate ω_i along the fibers of the submersion $\pi_i : U_i \rightarrow T_i$ to obtain $\int_{U_i} \omega_i \in \mathcal{A}_c(T_i)$. Define

$\int_F \omega \in \mathcal{A}_c(M/F)$ to be the class of $\sum_i \int_{U_i} \omega_i$. It is independent of the choice of the ω_i and of the cover $\mathcal{U}$.

As $\int_F$ commutes with d_M and d_H , it induces the map $\int_F : H^{p+k}(M) \rightarrow H_c^k(M/F)$.

3. TRACES

In this section, we give some algebraic background and construct the trace which will be used in the sequel.

There is a canonical lift of the normal bundle ν of F to a bundle $\nu_{\mathcal{G}} \subset T\mathcal{G}$ so that $T\mathcal{G} = TF_s \oplus TF_r \oplus \nu_{\mathcal{G}}$ and $r_*\nu_{\mathcal{G}} = \nu$, $s_*\nu_{\mathcal{G}} = \nu$. It is given as follows. Let $[\gamma] \in \mathcal{G}$ with $s([\gamma]) = x$, $r([\gamma]) = y$. Denote by $\exp : \nu \rightarrow M$ the exponential map. Given $X \in \nu_x$ and $t \in \mathbb{R}$ sufficiently small, there is a unique curve $\gamma_t : [0, 1] \rightarrow M$ so that

$$\text{i) } \gamma_t(0) = \exp tX \quad \text{ii) } \gamma_t \subset L_{\exp tX} \quad \text{iii) } \gamma_t(s) \in \exp(\nu_{\gamma(s)}).$$

In particular $\gamma_0 = \gamma$. Thus the family $[\gamma_t]$ in $\mathcal{G}$ defines a tangent vector $\hat{X} \in T\mathcal{G}_{[\gamma]}$. It is easy to check that $s_*(\hat{X}) = X$ and $r_*(\hat{X})$ is the parallel translate of X along γ to ν_y .

The metric g_0 on M induces a canonical metric g_0 on $\mathcal{G}$ as follows. $T\mathcal{G} = TF_s \oplus TF_r \oplus \nu_{\mathcal{G}}$ and these bundles are mutually orthogonal. On $TF_r \oplus \nu_{\mathcal{G}}$ we define g_0 to be s^*g_0 and on TF_s we define g_0 to be $r^*(g_0|TF)$.

The foliation F_s has normal bundle $\nu_s = TF_r \oplus \nu_{\mathcal{G}}$, so $\nu_s \simeq s^*(TM)$ and $\nu_s^* \simeq s^*(T^*M)$. Denote the space of smooth sections of the bundle $\Lambda^k T^*F_s \otimes \Lambda^h \nu_s^*$ by $\mathcal{A}^{k,h}(\mathcal{G}, F_s)$. The exterior derivative d splits with respect to the decomposition

$$\mathcal{A}^m(\mathcal{G}) \simeq \bigoplus_{k+h=m} \mathcal{A}^{k,h}(\mathcal{G}, F_s)$$

into three bihomogeneous components

$$d = d_F + d_\nu + d_{-1,2},$$

where d_F is the leafwise differential for F_s with bidegree $(1, 0)$ with respect to the splitting but which is independent of the choice of ν [BN03], d_ν is the $(0, 1)$ homogeneous component, and $d_{-1,2}$ is an extra differential which has bidegree $(-1, 2)$, see [T97] for more details.

Let E_1 be a complex vector bundle over M with Hermitian metric and connection, and set $E = r^*E_1$ with the pulled back metric and connection, denoted ∇ . We may regard ∇ as an operator of degree one on $C^\infty(E \otimes \Lambda T^*\mathcal{G})$ where on decomposable sections $\phi \otimes \omega$, $\nabla(\phi \otimes \omega) = (\nabla\phi) \wedge \omega + \phi \otimes d\omega$. Then ∇ defines a *quasi-connection* ∇^ν acting on $C^\infty(E \otimes \Lambda\nu_s^*)$ by the composition

$$C^\infty(E \otimes \Lambda\nu_s^*) \xrightarrow{i} C^\infty(E \otimes \Lambda T^*\mathcal{G}) \xrightarrow{\nabla} C^\infty(E \otimes \Lambda T^*\mathcal{G}) \xrightarrow{p_\nu} C^\infty(E \otimes \Lambda\nu_s^*),$$

where i is the inclusion and p_ν is induced by the projection $p_\nu : T^*\mathcal{G} \rightarrow \nu_s^*$. Given $\omega \in C^\infty(\Lambda\nu_s^*)$, denote by $M_\omega : C^\infty(E \otimes \Lambda\nu_s^*) \rightarrow C^\infty(E \otimes \Lambda\nu_s^*)$ the operator given by exterior multiplication on the left by ω . Then for any $\omega \in C^\infty(\Lambda^j\nu_s^*)$ and any $\phi \in C^\infty(E \otimes \Lambda\nu_s^*)$, ∇^ν satisfies the relation

$$\nabla^\nu(\omega\phi) - (-1)^j\omega\nabla^\nu\phi = (d_\nu\omega)\phi.$$

In other words, the graded commutator $[\nabla^\nu, M_\omega]$ of these operators on $C^\infty(E \otimes \Lambda\nu_s^*)$ is the operator $M_{d_\nu\omega}$ of exterior multiplication (on the left) by $d_\nu\omega$.

If $T \in \text{End}(C^\infty(E \otimes \Lambda\nu_s^*))$ we say that T is homogeneous of degree k if it maps $E \otimes \Lambda^j\nu_s^*$ to $E \otimes \Lambda^{j+k}\nu_s^*$. $C^\infty(E \otimes \Lambda\nu_s^*)$ is an $\mathcal{A}(M)$ -module where for $\phi \in C^\infty(E \otimes \Lambda\nu_s^*)$, and $\omega \in \mathcal{A}(M)$, we set

$$M_\omega\phi = p_\nu(s^*(\omega))\phi,$$

where again the map $p_\nu : \mathcal{A}(\mathcal{G}) \rightarrow C^\infty(\Lambda\nu_s^*)$ is induced by the projection $p_\nu : T^*\mathcal{G} \rightarrow \nu_s^*$. Denote by Ω^k the space of $\mathcal{A}(M)$ -equivariant endomorphisms of $C^\infty(E \otimes \Lambda\nu_s^*)$ with degree k and set

$$\Omega = \bigoplus_k \Omega^k.$$

Then $T \in \Omega^k$ if and only if for all $\omega \in \mathcal{A}^j(M)$,

$$[T, M_\omega] \equiv T \circ M_\omega - (-1)^{kj} M_\omega \circ T = 0.$$

A pseudo-differential $\mathcal{G}$ -operator with uniform support acting on E is a *smooth* family $(P_x)_{x \in M}$ of $\mathcal{G}$ -invariant pseudo-differential operators, where for each x , P_x is an operator acting on $E|_{\tilde{L}_x}$. The $\mathcal{G}$ -invariance property means that for any $\gamma \in \tilde{L}_x^y = \tilde{L}_x \cap \tilde{L}^y$, we have

$$(\gamma \cdot P)_y \equiv U_\gamma \circ P_x \circ U_\gamma^{-1} = P_y,$$

where $U_\gamma : C_c^\infty(\tilde{L}_x, E) \rightarrow C_c^\infty(\tilde{L}_y, E)$ is the induced action of the groupoid $\mathcal{G}$ defined by

$$[U_\gamma\xi](\gamma') = \xi(\gamma'\gamma), \quad \gamma' \in \tilde{L}_y.$$

The smoothness assumption is rigorously defined in [NWX96]. If we denote by K_x the Schwartz kernel of P_x , then the $\mathcal{G}$ -invariance assumption implies that the family $(K_x)_{x \in M}$ induces a distributional section K of the bundle $\text{Hom}(E)$ over the holonomy groupoid, whose fiber at a point γ is a homomorphism from $E_{s(\gamma)}$ to $E_{r(\gamma)}$, which is smooth outside $\mathcal{G}^{(0)} = M$. Since M is compact, the uniform support condition becomes the assumption that the support of K is compact in $\mathcal{G}$. We denote by $\Psi^m(\mathcal{G}; E)$ the space of all such uniformly supported pseudo-differential $\mathcal{G}$ -operators of order m , i.e. such that each P_x has order m . $\Psi^m(\mathcal{G}; E)$ is a $C^\infty(M)$ module, where $f \in C^\infty(M)$ acts by multiplication by $s^*(f)$. The space of all uniformly supported pseudo-differential $\mathcal{G}$ -operators is denoted $\Psi^\infty(\mathcal{G}; E)$ and the space of uniformly supported regularizing $\mathcal{G}$ -operators is denoted $\Psi^{-\infty}(\mathcal{G}; E)$. So,

$$\Psi^\infty(\mathcal{G}; E) = \bigcup_{m \in \mathbb{Z}} \Psi^m(\mathcal{G}; E) \text{ and } \Psi^{-\infty}(\mathcal{G}; E) = \bigcap_{m \in \mathbb{Z}} \Psi^m(\mathcal{G}; E).$$

The Schwartz Kernel Theorem identifies $\Psi^{-\infty}(\mathcal{G}; E)$ with $C_c^\infty(\mathcal{G}, \text{Hom}(E))$, see [NWX96]. We may consider the algebra

$$\Psi^{-\infty}(\mathcal{G}; E) \hat{\otimes}_{C^\infty(M)} \mathcal{A}(M)$$

as a subspace of Ω by using the $\mathcal{A}(M)$ module structure of $C^\infty(E \otimes \Lambda\nu_s^*)$.

Denote by $\partial_\nu : \text{End}(C^\infty(E \otimes \Lambda \nu_s^*)) \rightarrow \text{End}(C^\infty(E \otimes \Lambda \nu_s^*))$ the linear operator which on homogeneous elements of degree k is given by

$$\partial_\nu(T) = [\nabla^\nu, T] = \nabla^\nu \circ T - (-1)^k T \circ \nabla^\nu.$$

Lemma 3.1. ∂_ν preserves the subspace Ω , and $(\partial_\nu)^2$ is given by the commutator with $\theta = (\nabla^\nu)^2 \in \Omega^2$. In addition, ∂_ν preserves the subspace $\Psi^{-\infty}(\mathcal{G}; E) \widehat{\otimes}_{C^\infty(M)} \mathcal{A}(M)$.

Proof. Let T be a homogeneous endomorphism of $C^\infty(E \otimes \Lambda \nu_s^*)$ of degree k and let $\omega \in \mathcal{A}^j(M)$. Then

$$\begin{aligned} [\partial_\nu(T), M_\omega] &= \nabla^\nu \circ T \circ M_\omega - (-1)^k T \circ \nabla^\nu \circ M_\omega - \\ &\quad (-1)^{(k+1)j} M_\omega \circ \nabla^\nu \circ T + (-1)^{(k+1)j} (-1)^k M_\omega \circ T \circ \nabla^\nu \\ &= (-1)^{kj} (\nabla^\nu \circ M_\omega) \circ T - (-1)^k T \circ ((-1)^j M_\omega \circ \nabla^\nu + M_{d_\nu \omega}) - \\ &\quad (-1)^{kj+j} M_\omega \circ \nabla^\nu \circ T + (-1)^{kj+j+k} M_\omega \circ T \circ \nabla^\nu \\ &= (-1)^{kj+j} M_\omega \circ \nabla^\nu \circ T + (-1)^{kj} M_{d_\nu \omega} \circ T - (-1)^{k+j+kj} M_\omega \circ T \circ \nabla^\nu - \\ &\quad (-1)^{k(j+1)+k} M_{d_\nu \omega} \circ T - (-1)^{kj+j} M_\omega \circ \nabla^\nu \circ T + (-1)^{kj+k+j} M_\omega \circ T \circ \nabla^\nu \\ &= 0. \end{aligned}$$

In the same way we have

$$\begin{aligned} (\partial_\nu \circ \partial_\nu)(T) &= \nabla^\nu \circ \partial_\nu(T) - (-1)^{k+1} \partial_\nu(T) \circ \nabla^\nu \\ &= \nabla^\nu \circ (\nabla^\nu \circ T - (-1)^k T \circ \nabla^\nu) - (-1)^{k+1} (\nabla^\nu \circ T - (-1)^k T \circ \nabla^\nu) \circ \nabla^\nu \\ &= [\theta, T]. \end{aligned}$$

Finally, in order to prove that ∂_ν preserves $\Psi^{-\infty}(\mathcal{G}; E) \widehat{\otimes}_{C^\infty(M)} \mathcal{A}(M)$, we use the $\mathcal{A}(M)$ -module structure to reduce to the case where the element T under consideration actually belongs to $\Psi^{-\infty}(\mathcal{G}; E) = C_c^\infty(\mathcal{G}; \text{Hom}(E))$. By linearity, we may assume that T comes from a compactly supported smooth Schwartz kernel $K \in C_c^\infty((U, \gamma, V); \text{Hom}(E))$ for some basic open set (U, γ, V) . But then it is obvious that $\partial_\nu(K)$ is also smooth. Thus the proof is complete. $\square$

For $T \in \Psi^{-\infty}(\mathcal{G}; E) \widehat{\otimes}_{C^\infty(M)} \mathcal{A}^k(M) = C_c^\infty(\mathcal{G}; \text{Hom}(E)) \widehat{\otimes}_{C^\infty(M)} \mathcal{A}^k(M)$, define the trace of T to be the Haefliger k -form $\text{Tr}(T)$ given by

$$\text{Tr}(T) = \int_F \text{tr}(K(\bar{x}, \bar{x})) dx,$$

where K is the smooth Schwartz kernel of T , $\bar{x}$ is the class of the constant path at x , $\text{tr}(K(\bar{x}, \bar{x}))$ is the usual trace of $K(\bar{x}, \bar{x}) \in \text{End}(E_{\bar{x}}) \otimes \Lambda^k T^* M_x$ and so belongs to $\Lambda^k T^* M_x$, and dx is the leafwise volume form associated with the fixed orientation of the foliation F .

Lemma 3.2. The map $\text{Tr} : \Psi^{-\infty}(\mathcal{G}; E) \widehat{\otimes}_{C^\infty(M)} \mathcal{A}(M) \rightarrow \mathcal{A}_c(M/F)$ is a graded trace which satisfies $\text{Tr} \circ \partial_\nu = d_H \circ \text{Tr}$.

Proof. This is a corollary of Lemma 2.5, p. 443 of [HL02] which says that if K is a smooth section of $\text{Hom}(E) \widehat{\otimes}_{C^\infty(M)} \mathcal{A}(M)$ over $\mathcal{G}$ which is uniformly exponentially decaying (considered as a family of sections over the $\tilde{L}_x \times \tilde{L}_x$) and H is a continuous section of $\text{Hom}(E) \widehat{\otimes}_{C^\infty(M)} \mathcal{A}(M)$ over $\mathcal{G}$ with uniformly bounded coefficients, then $\text{Tr}([H, K]) = 0$. As the kernels in question here are all uniformly supported, they always have uniformly bounded coefficients. Likewise, for any such kernel, there is an $R > 0$ so that on any $\tilde{L}_x \times \tilde{L}_x$, the kernel is zero off the R neighborhood of the diagonal and so is uniformly exponentially decaying.

Now suppose that $T \in \Psi^{-\infty}(\mathcal{G}; E) \widehat{\otimes}_{C^\infty(M)} \mathcal{A}(M)$. As ∇^ν is local operator and everything is linear, we may assume that T is given by a section $K \in C_c^\infty((U, \bar{w}, U); \text{Hom}(E) \otimes \Lambda T^* U)$, where $w \in U$. The foliation chart $U \simeq \mathbb{R}^p \times \mathbb{R}^q$, and $(U, \bar{w}, U) \simeq \mathbb{R}^p \times \mathbb{R}^q \times \mathbb{R}^p$, with coordinates (x, y, z) . The map $s : (U, \bar{w}, U) \rightarrow U$ is

given by $s(x, y, z) = (x, y)$ and the foliation F_s restricted to $(U, \bar{w}, U)$ has leaves of the form $(x, y) \times \mathbb{R}^p$. In this setting, $\nabla^\nu = d_x + d_y + A$ where $A \in C^\infty((U, \bar{w}, U); \text{Hom}(E)) \hat{\otimes}_{C^\infty(U)} \mathcal{A}^1(U)$. Thus

$$\text{Tr}(\partial_\nu T) = \int_F \text{tr}([d_x + d_y + A, K])dx = \int_F \text{tr}(d_x K)dx + \int_F \text{tr}(d_y K)dx + \int_F \text{tr}([A, K])dx.$$

The term $\int_F \text{tr}(d_x K)dx$ is obviously zero. A direct computation shows that $\int_F \text{tr}([A, K])dx = 0$. Finally, note that in these coordinates, d_H is d_y if we take the transversal in U to be $(0, 0) \times \mathbb{R}^q$. Thus $\int_F \text{tr}(d_y K)dx = \int_F d_y \text{tr}(K)dx = d_H \int_F \text{tr}(K)dx = d_H \text{Tr}(T)$. $\square$

Since ∂_ν^2 is not necessarily zero, we now employ Connes' X -trick to construct a new graded differential algebra $(\tilde{\mathcal{A}}_{-\infty}, \delta)$ out of the graded quasi-differential algebra $(\mathcal{A}_c^\infty(\mathcal{G}|F_s; E), \partial_\nu)$, see [Con94], p. 229. Write $\mathcal{A}_{-\infty}$ for $\mathcal{A}_c^\infty(\mathcal{G}|F_s; E)$ and note that the curvature operator $\theta = \nabla^\nu \circ \nabla^\nu$ is a multiplier of $\mathcal{A}_{-\infty}$. As a vector space $\tilde{\mathcal{A}}_{-\infty} = M_2(\mathcal{A}_{-\infty})$. An element $\tilde{T} \in \tilde{\mathcal{A}}_{-\infty}$ is homogeneous of degree $\partial \tilde{T} = k$ if

$$k = \partial \tilde{T}_{11} = \partial \tilde{T}_{12} + 1 = \partial \tilde{T}_{21} + 1 = \partial \tilde{T}_{22} + 2.$$

The differential δ on $\tilde{\mathcal{A}}_{-\infty}$ is defined on elements of the form $\begin{pmatrix} \tilde{T}_{11} & 0 \\ 0 & 0 \end{pmatrix}$ by

$$\delta \tilde{T} = \begin{pmatrix} \partial_\nu \tilde{T}_{11} & (-1)^{\partial \tilde{T}_{11}} \tilde{T}_{11} \\ \tilde{T}_{11} & 0 \end{pmatrix},$$

for homogeneous $\tilde{T}_{11} \in \mathcal{A}_{-\infty}$. On homogeneous elements of $\tilde{\mathcal{A}}_{-\infty}$ it is given by

$$\delta \tilde{T} = \begin{pmatrix} \partial_\nu \tilde{T}_{11} & \partial_\nu \tilde{T}_{12} \\ -\partial_\nu \tilde{T}_{21} & -\partial_\nu \tilde{T}_{22} \end{pmatrix} + \begin{pmatrix} 0 & -\theta \\ 1 & 0 \end{pmatrix} \tilde{T} + (-1)^{\partial \tilde{T}} \tilde{T} \begin{pmatrix} 0 & 1 \\ -\theta & 0 \end{pmatrix}.$$

A straightforward computation gives $\delta^2 = 0$. Set

$$\Theta = \begin{pmatrix} 1 & 0 \\ 0 & \theta \end{pmatrix}$$

and define a new product on $\tilde{\mathcal{A}}_{-\infty}$ by

$$\tilde{T} * \tilde{T}' = \tilde{T} \Theta \tilde{T}'.$$

This makes $(\tilde{\mathcal{A}}_{-\infty}, \delta)$ a graded differential algebra.

The graded algebra $\mathcal{A}_{-\infty}$ embeds as a subalgebra of $\tilde{\mathcal{A}}_{-\infty}$ by using the map

$$T \mapsto \begin{pmatrix} T & 0 \\ 0 & 0 \end{pmatrix}.$$

We shall therefore also denote by T the image in $\tilde{\mathcal{A}}_{-\infty}$ of any $T \in \mathcal{A}_{-\infty}$.

For homogeneous $\tilde{T} \in \tilde{\mathcal{A}}_{-\infty}$ define

$$\Phi(\tilde{T}) = \text{Tr}(\tilde{T}_{11}) - (-1)^{\partial \tilde{T}} \text{Tr}(\tilde{T}_{22} \theta),$$

and extend to arbitrary elements by linearity.

Theorem 3.3. *The map $\Phi : \tilde{\mathcal{A}}_{-\infty} \rightarrow \mathcal{A}_c^*(M/F)$ is a graded trace, and $\Phi \circ \delta = d_H \circ \Phi$.*

Proof. Let $\tilde{T}, \tilde{S}$ be two homogeneous elements of $\tilde{\mathcal{A}}_{-\infty}$ with degree respectively k and ℓ . Then

$$\Phi([\tilde{T}, \tilde{S}]) = \text{Tr}([\tilde{T}_{11}, \tilde{S}_{11}]) + \text{Tr}([\tilde{T}_{12} \theta, \tilde{S}_{21}]) + (-1)^{k+\ell+1} \text{Tr}([\tilde{T}_{21}, \tilde{S}_{12} \theta]) + (-1)^{k+\ell+1} \text{Tr}([\tilde{T}_{22} \theta, \tilde{S}_{22} \theta]) = 0,$$

by Lemma 3.2.

For a homogeneous element $\tilde{T} \in \tilde{\mathcal{A}}_{-\infty}^{k-1}$,

$$(\Phi \circ \delta)(\tilde{T}) = (\text{Tr} \circ \partial_\nu)(\tilde{T}_{11}) + (-1)^k (\text{Tr} \circ \partial_\nu)(\tilde{T}_{22} \theta) - (\text{Tr} \circ (\partial_\nu)^2)(\tilde{T}_{21}),$$

since $\partial_\nu(\theta) = 0$. Using Lemma 3.2 again gives

$$(\Phi \circ \delta)(\tilde{T}) = (d_H \circ \text{Tr}) \left(\tilde{T}_{11} + (-1)^k \tilde{T}_{22} \theta \right) - (d_H^2 \circ \text{Tr})(\tilde{T}_{21}).$$

Since $d_H^2 = 0$, we get

$$(\Phi \circ \delta)(\tilde{T}) = d_H \left(\text{Tr} \tilde{T}_{11} + (-1)^k \text{Tr} \tilde{T}_{22} \theta \right),$$

and the proof is complete. $\square$

4. THE ALGEBRAIC CHERN CHARACTER

In this section, we use the map Φ to construct an algebraic Chern-Connes character

$$\text{ch}_a : K_0(C_c^\infty(\mathcal{G})) \longrightarrow H_c^*(M/F).$$

As $K_0(C_c^\infty(\mathcal{G})) \simeq K_0(C_c^\infty(\mathcal{G}; \text{Hom}(E))) \simeq K_0(\Psi^{-\infty}(\mathcal{G}; E))$, we may work with $K_0(\Psi^{-\infty}(\mathcal{G}; E))$ in place of $K_0(C_c^\infty(\mathcal{G}))$. Denote by $M_N(\Psi^{-\infty}(\mathcal{G}; E) \oplus \mathbb{C})$ the space of $N \times N$ matrices with coefficients in $\Psi^{-\infty}(\mathcal{G}; E) \oplus \mathbb{C}$.

Theorem 4.1. *Let $B = [\tilde{e}_1] - [\tilde{e}_2]$ be an element of $K_0(\Psi^{-\infty}(\mathcal{G}; E))$, where $\tilde{e}_1 = (e_1, \lambda_1)$ and $\tilde{e}_2 = (e_2, \lambda_2)$ are idempotents in $M_N(\Psi^{-\infty}(\mathcal{G}; E) \oplus \mathbb{C})$, which we consider as elements of $M_N(\tilde{\mathcal{A}}_{-\infty} \oplus M_2(\mathbb{C}))$. Then the Haefliger form*

$$(\Phi \circ \text{tr}) \left(e_1 \exp \left(\frac{-(\delta e_1)^2}{2i\pi} \right) \right) - (\Phi \circ \text{tr}) \left(e_2 \exp \left(\frac{-(\delta e_2)^2}{2i\pi} \right) \right)$$

is closed and its Haefliger cohomology class depends only on B . Here $\text{tr} : M_N(\tilde{\mathcal{A}}_{-\infty}) \rightarrow \tilde{\mathcal{A}}_{-\infty}$ is the usual trace.

Definition 4.2. *Let $B = [\tilde{e}_1] - [\tilde{e}_2] \in K_0(\Psi^{-\infty}(\mathcal{G}; E))$ where $\tilde{e}_1 = (e_1, \lambda_1)$ and $\tilde{e}_2 = (e_2, \lambda_2)$ are idempotents. The algebraic Chern character $\text{ch}_a(B)$ of B is the Haefliger cohomology class*

$$\text{ch}_a(B) = [(\Phi \circ \text{tr}) \left(e_1 \exp \left(\frac{-(\delta e_1)^2}{2i\pi} \right) \right)] - [(\Phi \circ \text{tr}) \left(e_2 \exp \left(\frac{-(\delta e_2)^2}{2i\pi} \right) \right)].$$

Proof. The derivation δ extends to a derivation on $\tilde{\mathcal{A}}_{-\infty} \oplus \mathbb{C}$ (and so also to $M_N(\tilde{\mathcal{A}}_{-\infty} \oplus \mathbb{C})$) by setting $\delta(e, \lambda) = (\delta e, 0)$. In addition, Φ extends to a graded trace $\Phi : \tilde{\mathcal{A}}_{-\infty} \oplus \mathbb{C} \rightarrow \mathcal{A}_c(M/F)$ by setting $\Phi((e, \lambda)) = \Phi(e)$, and Theorem 3.3 remains valid for this extended trace.

Let $e \in M_N(\tilde{\mathcal{A}}_{-\infty} \oplus M_2(\mathbb{C}))$ be an idempotent, and set $u = 2e - 1$. Then

$$u^2 = 1 \text{ and } u \cdot \delta e + \delta e \cdot u = 0.$$

By Theorem 3.3, $d_H(\Phi \circ \text{tr}) \left(e(\delta e)^{2k} \right) = (\Phi \circ \text{tr}) \left((\delta e)^{2k+1} \right)$ and we have

$$(\Phi \circ \text{tr}) \left((\delta e)^{2k+1} \right) = (\Phi \circ \text{tr}) \left(u^2 (\delta e)^{2k+1} \right) = -(\Phi \circ \text{tr}) \left(u (\delta e)^{2k+1} u \right).$$

The last equality is a consequence of the relation $u \cdot \delta e = -\delta e \cdot u$. On the other hand, since $\Phi \circ \text{tr}$ is a trace, we have

$$-(\Phi \circ \text{tr}) \left(u (\delta e)^{2k+1} u \right) = -(\Phi \circ \text{tr}) \left((\delta e)^{2k+1} u^2 \right) = -(\Phi \circ \text{tr}) \left((\delta e)^{2k+1} \right),$$

so

$$d_H \left((\Phi \circ \text{tr}) (e(\delta e)^{2k}) \right) = 0.$$

Now suppose e_t is a smooth path of idempotents in $M_N(\Psi^{-\infty}(\mathcal{G}; E) \oplus \mathbb{C})$ and set $a_t = \frac{de_t}{dt} (2e_t - 1)$. Then

$$\begin{aligned} [a_t, e_t] &= a_t e_t - e_t a_t \\ &= \frac{de_t}{dt} (2e_t - 1) e_t - e_t \frac{de_t}{dt} (2e_t - 1) \\ &= \frac{de_t}{dt} e_t + e_t \frac{de_t}{dt} - 2e_t \frac{de_t}{dt} e_t. \end{aligned}$$

Since $e_t^2 = e_t$ for any t , we have

$$\frac{de_t}{dt}e_t + e_t\frac{de_t}{dt} = \frac{de_t}{dt} \text{ and } e_t\frac{de_t}{dt}e_t = 0,$$

so

$$\frac{de_t}{dt} = [a_t, e_t].$$

This implies that the zero th order component of $\frac{d}{dt}(\Phi \circ \text{tr})\left(e_t \exp\left[\frac{-(\delta e_t)^2}{2i\pi}\right]\right)$ is zero.

For any $k \geq 1$ we have by multilinearity

$$\begin{aligned} \frac{d}{dt}(\Phi \circ \text{tr})\left(e_t(\delta e_t)^{2k}\right) &= (\Phi \circ \text{tr})\left(\frac{de_t}{dt}(\delta e_t)^{2k}\right) + \sum_{j=1}^{2k} (\Phi \circ \text{tr})\left(e_t(\delta e_t)^{j-1}\left(\delta \frac{de_t}{dt}\right)(\delta e_t)^{2k-j}\right) \\ &= (\Phi \circ \text{tr})\left([a_t, e_t](\delta e_t)^{2k}\right) + \sum_{j=1}^{2k} (\Phi \circ \text{tr})\left(e_t(\delta e_t)^{j-1}\delta[a_t, e_t](\delta e_t)^{2k-j}\right) \\ &= (\Phi \circ \text{tr})\left([a_t, e_t](\delta e_t)^{2k}\right) + \sum_{j=1}^{2k} (\Phi \circ \text{tr})\left([e_t(\delta e_t)^{j-1}a_t(\delta e_t)^{2k-j+1} - e_t(\delta e_t)^ja_t(\delta e_t)^{2k-j}\right) + \\ &\quad \sum_{j=1}^{2k} (\Phi \circ \text{tr})\left(e_t(\delta e_t)^{j-1}[(\delta a_t)e_t - e_t(\delta a_t)](\delta e_t)^{2k-j}\right). \end{aligned}$$

The first sum collapses to

$$(\Phi \circ \text{tr})\left(e_t a_t (\delta e_t)^{2k}\right) - (\Phi \circ \text{tr})\left(e_t (\delta e_t)^{2k} a_t\right) = -(\Phi \circ \text{tr})\left([a_t, e_t](\delta e_t)^{2k}\right),$$

so the first two terms cancel.

To analyze the second sum, note that since e_t is an idempotent and δ is a derivation, by induction we have

$$e_t(\delta e_t)^{2h}e_t = e_t(\delta e_t)^{2h} = (\delta e_t)^{2h}e_t, \quad e_t(\delta e_t)^{2h+1}e_t = 0,$$

$$\text{and } e_t(\delta e_t)^{2h+1} + (\delta e_t)^{2h+1}e_t = (\delta e_t)^{2h+1} \quad \forall h \geq 0.$$

The second sum is then

$$\sum_{j=1}^{2k} (\Phi \circ \text{tr})\left(e_t(\delta e_t)^{j-1}(\delta a_t)e_t(\delta e_t)^{2k-j} - e_t(\delta e_t)^{j-1}e_t(\delta a_t)(\delta e_t)^{2k-j}\right).$$

Using the fact that $\Phi \circ \text{tr}$ is a graded trace, this equals

$$\begin{aligned} &\sum_{j=1}^{2k} (\Phi \circ \text{tr})\left((-1)^j e_t(\delta e_t)^{2k-j} e_t(\delta e_t)^{j-1}(\delta a_t)\right) + \sum_{j=1}^{2k} (\Phi \circ \text{tr})\left((-1)^{(j+1)}(\delta e_t)^{2k-j} e_t(\delta e_t)^{j-1}e_t(\delta a_t)\right) \\ &= \sum_{i=1}^k (\Phi \circ \text{tr})\left(e_t(\delta e_t)^{2k-2i} e_t(\delta e_t)^{2i-1}(\delta a_t)\right) + \sum_{i=1}^k (\Phi \circ \text{tr})\left((\delta e_t)^{2k-2i+1} e_t(\delta e_t)^{2i-2} e_t(\delta a_t)\right) \\ &= \sum_{i=1}^k (\Phi \circ \text{tr})\left(e_t(\delta e_t)^{2k-1}(\delta a_t)\right) + \sum_{i=1}^k (\Phi \circ \text{tr})\left(-e_t(\delta e_t)^{2k-1}(\delta a_t) + (\delta e_t)^{2k-1}(\delta a_t)\right) \\ &= \sum_{i=1}^k (\Phi \circ \text{tr})\left((\delta e_t)^{2k-1}(\delta a_t)\right) = k d_H(\Phi \circ \text{tr})\left(e_t(\delta e_t)^{2k-2}(\delta a_t)\right). \end{aligned}$$

Thus $\text{ch}_a(B)$ does not depend on the choice of representative of B .

Now suppose that ${}^t\nabla$ is a smooth family of connections on E_1 with associated differentials ∂_t , and δ_t . On $M \times I$, where $I = [0, 1]$, we have the foliation $\widehat{F}$ whose leaves are of the form $L \times \{t\}, t \in I$. Then

$\mathcal{G}_{\widehat{F}} = \mathcal{G} \times I$, and $M \times I/\widehat{F} = M/F \times I$. Denote by $\pi : \mathcal{G} \times I \rightarrow \mathcal{G}$ the projection. On the bundle π^*E , we have the connection ∇ determined by ${}^t\nabla$. In particular, for $\phi \in C^\infty(E)$, and $X \in T(\mathcal{G} \times I)$, $\nabla_X(\phi \circ \pi)(x, t) \equiv {}^t\nabla_{\pi_*(X)}(\phi)(x)$, where we identify $\pi^*E_{(x,t)}$ with E_x . Thus on $\mathcal{G} \times \{t\}$, $\nabla = {}^t\nabla + d_t$, where $d_t = dt \otimes \partial/\partial t$, so the associated partial connection ∇^ν on $\mathcal{A}(\mathcal{G}_{\widehat{F}}, \pi^*E)$ is $\nabla^\nu = {}^t\nabla^\nu + d_t$. Therefore, the curvature θ of ∇^ν satisfies $i_t^*(\theta) = \theta_t$ the curvature of ${}^t\nabla^\nu$, where $i_t : \mathcal{G} \rightarrow \mathcal{G} \times I$ is $i_t(x) = (x, t)$. Finally, the differential ∂_ν associated to ∇ is $\partial_\nu = \partial_t + d_t$.

Denote by $\int_I : A_c(M \times I/\widehat{F}) \rightarrow A_c(M/F)$ integration along the fiber of $M \times I/\widehat{F} = M/F \times I \rightarrow M/F$, and let $r_j : M/F \rightarrow M \times I/\widehat{F}$ be given by $r_j(x) = (x, j)$. The usual proof shows that

$$r_1^* - r_0^* = \int_I \circ d_{\widehat{H}} \pm d_H \circ \int_I.$$

For any idempotent $(e, \lambda) \in M_N(\Psi^{-\infty}(\mathcal{G}; E) \oplus \mathbb{C})$, the associated idempotent in $M_N(\Psi^{-\infty}(\mathcal{G} \times I; E) \oplus \mathbb{C})$ is denoted by (π^*e, λ) . As

$$d_{\widehat{H}}((\Phi \circ \text{tr})(\pi^*e(\delta\pi^*e)^{2k})) = 0,$$

we have

$$r_1^*((\Phi \circ \text{tr})(\pi^*e(\delta\pi^*e)^{2k})) - r_0^*((\Phi \circ \text{tr})(\pi^*e(\delta\pi^*e)^{2k})) = \pm d_H \left(\int_I (\Phi \circ \text{tr})(\pi^*e(\delta\pi^*e)^{2k}) \right).$$

Now (up to signs whose ambiguity is the same in both expressions below)

$$(\delta\pi^*e)^2 = \begin{pmatrix} (\partial_\nu \pi^*e)^2 + (\pi^*e)\theta(\pi^*e) & (\partial_\nu \pi^*e)\pi^*e \\ \pi^*e(\partial_\nu \pi^*e) & (\pi^*e)^2 \end{pmatrix},$$

so

$$i_j^*(\delta\pi^*e)^2 = \begin{pmatrix} (\partial_j e)^2 + e\theta_j e & (\partial_j e)e \\ e(\partial_j e) & e^2 \end{pmatrix} = (\delta_j e)^2$$

as $d_j(\pi^*(e)) = 0$. Since $i_j^*(\theta) = \theta_j$, we have $i_j^*((\delta\pi^*e)^{2k}) = (\delta_j e)^{2k}$. But $r_j^* \circ \Phi \circ \text{tr} = \Phi \circ \text{tr} \circ i_j^*$, and $i_j^*(\pi^*e) = e$, so

$$r_j^*((\Phi \circ \text{tr})(\pi^*e(\delta\pi^*e)^{2k})) = ((\Phi \circ \text{tr})(i_j^*(\pi^*e(\delta\pi^*e)^{2k}))) = (\Phi \circ \text{tr})(e(\delta_j e)^{2k}),$$

and we have the theorem. $\square$

5. PUSH FORWARD MAPS

In this section we construct a push forward map in cohomology for oriented maps from a manifold N to the space of leaves M/F . We will also give a geometric construction of the Connes-Skandalis push forward map [CS84] in K-theory.

Recall the following from [CS84].

Definition 5.1. A map $f : N \rightarrow M/F$ is a cocycle $(V_\alpha, f_{\alpha\beta})$ on N with values in $\mathcal{G}$.

Here $\mathcal{V} = \{V_\alpha\}$ is a locally finite open cover of N and the maps $f_{\alpha\beta} : V_\alpha \cap V_\beta \rightarrow \mathcal{G}$ satisfy $f_{\alpha\beta}(x)f_{\beta\gamma}(x) = f_{\alpha\gamma}(x)$ for all $x \in V_\alpha \cap V_\beta \cap V_\gamma$.

Note that as $f_{\alpha\alpha}f_{\alpha\alpha} = f_{\alpha\alpha}$, $f_{\alpha\alpha}$ maps V_α into the space of units $\mathcal{G}^{(0)} = M$. The holonomy maps along the $f_{\alpha\beta}$ provide the gluing maps to glue the local bundles $f_{\alpha\alpha}^*\nu \rightarrow V_\alpha$ together to get the pull-back vector bundle $f^*\nu \rightarrow N$ of the normal bundle ν . The map f is oriented if the vector bundle $TN \oplus f^*\nu$ is oriented. It is a submersion provided that each $f_{\alpha\alpha} : V_\alpha \rightarrow M$ is transverse to F . We assume that f is an oriented submersion until further notice. As above, $\mathcal{U} = \{U_i\}$ is a finite good cover of M by foliation charts for F . We may assume that each $\overline{f_{\alpha\alpha}(V_\alpha)} \subset U_i$ for some i , for if not, we may replace $\mathcal{V}$ with a locally finite refinement for which this is true. For each α , choose $i(\alpha)$ so that $\overline{f_{\alpha\alpha}(V_\alpha)} \subset U_{i(\alpha)}$.

Since the $f_{\alpha\alpha}$ are submersions, they induce foliations of the V_α and because of the cocycle condition, these local foliations fit together to give a global foliation F_N of N . We may assume that the V_α are foliation charts for F_N . Let S_α be a transversal of V_α . The map $f_{\alpha\alpha}$ induces a map $f_\alpha : S_\alpha \rightarrow T_{i(\alpha)}$ which is a diffeomorphism onto its image, so we get a map $f_{\alpha*} : \mathcal{A}_c(S_\alpha) \rightarrow \mathcal{A}_c(T_{i(\alpha)})$. Let $\omega = (\omega_\alpha) \in \mathcal{A}_c(N/F_N)$, be a

Haeffliger form, where $\omega_\alpha \in \mathcal{A}_c(S_\alpha)$. As ω has compact support and the cover $\mathcal{V}$ is locally finite, there are at most a finite number of α for which $\omega_\alpha \neq 0$. It then follows easily that $\sum_\alpha f_{\alpha*}$ induces a well defined map denoted $f_* : \mathcal{A}_c(N/F_N) \rightarrow \mathcal{A}_c(M/F)$. Since f is oriented, F_N is an oriented foliation on N . Recall the map $\int_{F_N} : \mathcal{A}_c(N) \rightarrow \mathcal{A}_c(N/F_N)$.

Definition 5.2. For an oriented submersion $f : N \rightarrow M/F$, $\int_f : \mathcal{A}_c(N) \rightarrow \mathcal{A}_c(M/F)$ is defined to be

$$\int_f = f_* \circ \int_{F_N}.$$

Denote the cohomology class of any closed differential form α by $[\alpha]$, and similarly for Haeffliger forms.

Definition 5.3. For an oriented submersion $f : N \rightarrow M/F$, the push forward map $f_! : H_c^*(N; \mathbb{R}) \rightarrow H_c^*(M/F)$ is defined by $f_!([\omega]) = [\int_f \omega]$.

That this is well defined follows immediately from

Proposition 5.4.

$$d_H \circ \int_f = \int_f \circ d_N.$$

Proof. Denote the Haeffliger differentials on M and N by d_H^M and d_H^N respectively. The result then follows from the fact that $d_H^M \circ f_* = f_* \circ d_H^N$ and $d_H^N \circ \int_{F_N} = \int_{F_N} \circ d_N$. $\square$

This construction may be generalized as follows. Suppose that F_1 is another foliation of N and that $\widehat{f} : N/F_1 \rightarrow M/F$ is a submersion. This means that there is a submersion $f : N \rightarrow M/F$ so that the leaves of F_1 are submanifolds of the leaves F_N . We can then construct a map $\widehat{f}_! : H_c^*(N/F_1) \rightarrow H_c^*(M/F)$ using the techniques above. Assume that the cover $\mathcal{V}$ is a cover by foliation charts for F_1 as well as for F_N . Denote the transversal of V_α for F_1 by $\widehat{S}_\alpha$. We may assume that $\widehat{S}_\alpha$ is a fiber bundle over S_α with projection π . Integration over the fibers of these bundles defines a map $\int_\pi : \mathcal{A}_c(N/F_1) \rightarrow \mathcal{A}_c(N/F_N)$. We then define $\widehat{f}_! : H_c^*(N/F_1) \rightarrow H_c^*(M/F)$ to be $\widehat{f}_!([\omega]) = [f_* (\int_\pi \omega)]$. Our particular case above is given by taking F_1 to be the foliation of N by its points, so that $H_c^*(N/F_1) = H_c^*(N; \mathbb{R})$.

Recall that a map $f : N \rightarrow M$ between two manifolds is oriented if the bundle $TN \oplus f^*TM$ is oriented. Given such a map, there is a push forward map $f_! : H_c^*(N; \mathbb{R}) \rightarrow H_c^*(M; \mathbb{R})$ which is defined using Poincaré duality. If $g : M \rightarrow W$ is another oriented map, then $(g \circ f)_! = g_! \circ f_!$. If f is a submersion, $f_!$ is also given by integrating over the fibers of f . (This is just the construction above, taking F to be the foliation of M by its points).

Now suppose that $f : N \rightarrow M/F$ is an arbitrary oriented map. We shall show below that there is a manifold W and oriented maps $i : N \rightarrow W$ and $\widehat{f} : W \rightarrow M/F$ where $\widehat{f}$ is a submersion, and $f = \widehat{f} \circ i$.

Definition 5.5. Suppose $f : N \rightarrow M/F$ is written as $f = \widehat{f} \circ i$ as above. Define $f_! : H_c^*(N; \mathbb{R}) \rightarrow H_c^*(M/F)$ to be the composition $f_! = \widehat{f}_! \circ i_!$.

Proposition 5.6. $f_!$ is well defined. In particular, if W_1, W_2 are smooth manifolds, and $i_j : N \rightarrow W_j$ and $f_j : W_j \rightarrow M/F$ are oriented smooth maps where $f = f_j \circ i_j$ and each f_j is a submersion, then $f_! \circ i_{1!} = f_{2!} \circ i_{2!}$.

Proof. [See [CS84], p.1169] First note that if we have maps $X \xrightarrow{g} W \xrightarrow{f} M/F$ where both g and f are oriented submersions, then $f \circ g$ is an oriented submersion, and $(f \circ g)_! = f_! \circ g_!$. This is just Fubini's Theorem, coupled with the fact that $g_!$ can be defined by integration over its fibers.

In [CS84], the following commutative diagram is constructed

$$\begin{array}{ccccc}
 & & W_1 & & \\
 & \nearrow i_1 & \uparrow \pi_1 & \searrow f_1 & \\
 N & \xrightarrow{i} & W & \xrightarrow{f} & M/F \\
 & \searrow i_2 & \downarrow \pi_2 & \nearrow f_2 & \\
 & & W_2 & &
 \end{array}$$

where W is essentially the fibered product of W_1 and W_2 over M/F , and f , π_1 and π_2 are submersions. It is clear that since i_1 and i_2 are oriented, so is the fibered product map i . In the same way the maps π_1 and π_2 are clearly oriented. Then we have

$$f_{1!} \circ i_{1!} = f_{1!} \circ (\pi_1 \circ i)_! = f_{1!} \circ \pi_{1!} \circ i_! = (f_1 \circ \pi_1)_! \circ i_! = f_! \circ i_!.$$

Similarly, $f_{2!} \circ i_{2!} = f_! \circ i_!$. □

To construct a manifold W and oriented maps $i : N \rightarrow W$ and $\hat{f} : W \rightarrow M/F$ where $\hat{f}$ is a submersion and $f = \hat{f} \circ i$, we follow [Con81], Section 11. For the benefit of the reader, let us sketch briefly the construction of W . Recall the foliated microbundle $N \xrightarrow{i_1} W_1 \xrightarrow{p} N$ associated to a Γ_q structure $\mathcal{F}$ on N introduced by Haefliger, [H58, H70]. Suppose the foliation F is given by the submersions $k_i : U_i \rightarrow \mathbb{R}^q$ with changes of coordinates $k_{ji} : k_i(U_i \cap U_j) \rightarrow k_j(U_i \cap U_j)$. For each i , let $k_i^{-1} : \mathbb{R}^q \rightarrow U_i$ be a transversal (so $k_i k_i^{-1} = Id$). Note that in general, $k_i^{-1} k_i(x) \neq x$, but these points are always on the same plaque of U_i . As above, assume that $\overline{f_{\alpha\alpha}(V_\alpha)} \subset U_{i(\alpha)}$. The Γ_q structure $\mathcal{F}$ on N associated to f and F is then given by $\mathcal{F} = (V_\alpha, h_\alpha, h_{\alpha\beta}^x)$, where $h_\alpha = k_{i(\alpha)} f_{\alpha\alpha} : V_\alpha \rightarrow \mathbb{R}^q$. For each $x \in V_\alpha \cap V_\beta$, the local diffeomorphism $h_{\alpha\beta}^x$ from a neighborhood of $h_\beta(x)$ to a neighborhood of $h_\alpha(x)$ is $h_{\alpha\beta}^x = k_{i(\alpha)} H_{\alpha\beta}^x k_{i(\beta)}^{-1}$, where $H_{\alpha\beta}^x$ is the holonomy determined by the path $f_{\alpha\beta}(x)$ from $f_{\beta\beta}(x)$ to $f_{\alpha\alpha}(x)$ acting from its domain in $k_{i(\beta)}^{-1}(\mathbb{R}^q)$ to its range in $k_{i(\alpha)}^{-1}(\mathbb{R}^q)$. Note that $\{h_{\alpha\beta}^x\}$ is then a cocycle. Set $\tilde{V}_\alpha = V_\alpha \times \mathbb{R}^q$ and consider the disjoint union $\coprod \tilde{V}_\alpha$. Each of the sets $\tilde{V}_\alpha$ has the codimension q foliation whose leaves are given by $V_\alpha \times \{r\}$ where $r \in \mathbb{R}^q$. Define $i_\alpha : V_\alpha \rightarrow \tilde{V}_\alpha$ by $i_\alpha(x) = (x, h_\alpha(x))$ and $p_\alpha : \tilde{V}_\alpha \rightarrow V_\alpha$ by $p_\alpha(x, r) = x$. The space $\tilde{W}$ is then obtained from $\coprod \tilde{V}_\alpha$ by identifying $(x, s) \in \tilde{V}_\beta$ with $(x, h_{\alpha\beta}^x(s)) \in \tilde{V}_\alpha$. The maps i_α and p_α determine well defined maps, $i_1 : N \rightarrow \tilde{W}$, and $p : \tilde{W} \rightarrow N$. The map p is a submersion, and $p \circ i_1 = Id$. The local foliations also fit together to give a global foliation $F_{\tilde{W}}$ of $\tilde{W}$, and it is easy to see that $\mathcal{F}$ is induced by i_1 and $F_{\tilde{W}}$. Note that $\tilde{W}$ is not a manifold in general as it will not be locally Euclidean due to the fact that the $h_{\alpha\beta}^x$ are only locally defined. This problem is eliminated by taking W_1 to be a sufficiently small neighborhood of the image of the i_1 in $\tilde{W}$.

Next we construct a submersion $f_1 : W_1 \rightarrow M/F$ so that $f = f_1 \circ i_1$. Each foliation chart U_i has coordinates $U_i \simeq \mathbb{R}^p \times \mathbb{R}^q$ with the plaques given by the $\mathbb{R}^p \times \{r\}$. Set $\hat{V}_\alpha = p^{-1}V_\alpha$ which we may consider as a subset of $\tilde{V}_\alpha = V_\alpha \times \mathbb{R}^q$. Define $f_{1\alpha,\alpha} : \hat{V}_\alpha \rightarrow U_{i(\alpha)}$ by

$$f_{1\alpha,\alpha}(x, r) = f_{\alpha,\alpha}(x) + (0, r).$$

We may assume that W_1 is a sufficiently small enough neighborhood of the image of i_1 in $\tilde{W}$ so that the $f_{1\alpha,\alpha}$ are well defined. It is obvious that each $f_{1\alpha,\alpha}$ is transverse to F . Now suppose that $\tilde{V}_\alpha \cap \tilde{V}_\beta \neq \emptyset$. A point z in this intersection has two different coordinates, namely $(x, r) \in V_\alpha \times \mathbb{R}^q$ and $(x, h_{\beta,\alpha}^x(r)) \in V_\beta \times \mathbb{R}^q$. Define $f_{1\beta,\alpha} : \hat{V}_\alpha \cap \hat{V}_\beta \rightarrow \mathcal{G}$ by $f_{1\beta,\alpha}(z)$ is the class of the leafwise path starting at $f_{1\alpha,\alpha}(x, r)$, ending at the point $f_{1\beta,\beta}(x, h_{\beta,\alpha}^x(r))$ and parallel to the leafwise path determined by $f_{\beta,\alpha}(x)$. We leave it to the reader to check that this is well defined and gives a submersion $f_1 : W_1 \rightarrow M/F$. It is obvious that $f = f_1 \circ i_1$.

To finish the construction of W , i and $\widehat{f}$, we take care of the problem of making both $\widehat{f}$ and i oriented. Let $\xi \rightarrow N$ be the normal bundle of an imbedding of N in some Euclidean space. Let $W = p^*\xi$ and denote by $\pi : W \rightarrow W_1$ the projection, and by $s : W_1 \rightarrow W$ the zero section. Set

$$i = s \circ i_1 : N \rightarrow W \quad \text{and} \quad \widehat{f} = f_1 \circ \pi : W \rightarrow M/F.$$

Then $f = \widehat{f} \circ i$ since $\pi \circ s = Id$, and $\widehat{f}$ is a submersion.

It is easy to see that $i_1^*TW_1 \simeq TN \oplus f^*\nu$ and $s^*TW = s^*Tp^*\xi = TW_1 \oplus p^*\xi$, so $i^*TW \simeq TN \oplus f^*\nu \oplus \xi$. Thus $TN \oplus i^*TW \simeq TN \oplus f^*\nu \oplus TN \oplus \xi$ is oriented since f is oriented and $TN \oplus \xi$ is a trivial bundle.

Finally, using the fact that $i_1(N)$ is a deformation retract of W_1 , which implies that $(i_1 \circ p)^*E \simeq E$ for any bundle E over W_1 , we have

$$\begin{aligned} TW \oplus \widehat{f}^*\nu &\simeq \pi^*(TW_1 \oplus p^*\xi) \oplus \pi^*f_1^*\nu \\ &\simeq \pi^*(p^*(TN \oplus f^*\nu) \oplus p^*\xi \oplus (i_1 \circ p)^*f_1^*\nu) \\ &\simeq \pi^*(p^*(TN \oplus f^*\nu) \oplus p^*\xi \oplus p^*i_1^*f_1^*\nu) \\ &\simeq \pi^*p^*(TN \oplus f^*\nu \oplus \xi \oplus f^*\nu) \\ &\simeq \pi^*p^*(TN \oplus \xi \oplus f^*\nu \oplus f^*\nu), \end{aligned}$$

which is oriented.

We now give a geometric construction of the push forward map $f_! : K_c^0(N) \rightarrow K_0(C^*(M, F))$ associated to the K-oriented map $f : N \rightarrow M/F$ constructed by Connes and Skandalis in [CS84]. $C^*(M, F)$ is Connes' foliation C^* algebra for F . We will use the fact that $f_!$ takes values in the image of $K_0(C_c^\infty(\mathcal{G}))$ under the inclusion map in K -theory. The map f is K-oriented provided that the bundle $T^*N \oplus f^*(\nu)$ is endowed with a Spin^c structure. If f is K-oriented, it is oriented. The fact that f is K-oriented implies that i is K-oriented, so we have the push forward map $i_! : K_c^0(N) \rightarrow K_c^0(W)$.

Denote by $\widehat{F}$ the foliation of $M \times \mathbb{R}^{2k}$ whose leaves are of the form $L \times \{x\}$ where L is a leaf of F and $x \in \mathbb{R}^{2k}$. If we denote the holonomy groupoid of $\widehat{F}$ by $\mathcal{G}^{\mathbb{R}^{2k}}$, then $\mathcal{G}^{\mathbb{R}^{2k}} = \mathcal{G} \times \mathbb{R}^{2k}$, and the space of leaves of $\widehat{F}$ is $M \times \mathbb{R}^{2k}/\widehat{F} \simeq M/F \times \mathbb{R}^{2k}$. Let the map $\widehat{f}$ be represented by the cocycle $\{\widehat{f}_{\alpha\beta}, V_\alpha\}$ where $\mathcal{V} = \{V_\alpha\}$ is a locally finite open cover of W . Choose an imbedding $g : W \rightarrow \mathbb{R}^{2k}$ and let $h : W \rightarrow M \times \mathbb{R}^{2k}/\widehat{F}$ be given by the cocycle $\{h_{\alpha\beta}, V_\alpha\}$ where

$$h_{\alpha\beta}(x) = (\widehat{f}_{\alpha\beta}(x), g(x)).$$

Now denote by $\mathcal{U} = \{U_i\}$ a locally finite good cover of $M \times \mathbb{R}^{2k}$ by foliation charts for $\widehat{F}$. As above, we may assume that each $\overline{h_{\alpha\alpha}(V_\alpha)} \subset U_i$ for some i , and for each α we may choose $i(\alpha)$ so that $\overline{h_{\alpha\alpha}(V_\alpha)} \subset U_{i(\alpha)}$. Note that for each α , $h_{\alpha\alpha}$ is an imbedding of V_α in a transversal of the foliation $\widehat{F}$. This implies that each $h_{\alpha\beta}$ is an imbedding in a transversal of the combined foliation $\widehat{F}_s \oplus \widehat{F}_r$ of $\mathcal{G}^{\mathbb{R}^{2k}}$.

Recall the open cover of $\mathcal{G}^{\mathbb{R}^{2k}}$ associated to the cover $\mathcal{U}$ of $M \times \mathbb{R}^{2k}$. For each $U_i \in \mathcal{U}$ choose a smooth connected transversal $T_i \subset U_i$. We may assume that these transversals are pairwise disjoint. Let U_i and U_j be in $\mathcal{U}$, and let γ_{ij} be a smooth map $\gamma_{ij} : T_i \rightarrow \mathcal{G}^{\mathbb{R}^{2k}}$ so that for each $y \in \text{domain}(\gamma_{ij}) \subset T_i$, $s(\gamma_{ij}(y)) \in T_i$ and $r(\gamma_{ij}(y)) \in T_j$. The open subset of $\mathcal{G}^{\mathbb{R}^{2k}}$

$$(U_i, \gamma_{ij}, U_j)$$

consists of the classes of all leafwise paths σ which start in U_i , end in U_j , and if the path σ is in the leaf of $\gamma_{ij}(y)$ then σ is parallel to $\gamma_{ij}(y)$. We may also describe (U_i, γ_{ij}, U_j) as the classes of all leafwise paths of the form $\gamma_j \circ \gamma_{ij}(y) \circ \gamma_i$, where γ_i , respectively γ_j , is entirely contained in the plaque of U_i , respectively U_j , containing $s(\gamma_{ij}(y))$, respectively $r(\gamma_{ij}(y))$. Note that the image $\gamma_{ij}(T_i)$, denoted T_{ij} , is a submanifold of (U_i, γ_{ij}, U_j) which is a transversal of the combined foliation $\widehat{F}_s \oplus \widehat{F}_r$.

The image of any map $h_{\alpha\beta} : V_\alpha \cap V_\beta \rightarrow \mathcal{G}^{\mathbb{R}^{2k}}$ is contained in a chart of the form $(U_{i(\alpha)}, \gamma_{\alpha\beta}, U_{i(\beta)})$ as above. We now alter the $h_{\alpha\beta}$ so that the new $h_{\alpha\beta}$ satisfy $h_{\alpha\beta}(V_\alpha \cap V_\beta) \subset \gamma_{\alpha\beta}(T_{i(\alpha)})$. For each $x \in V_\alpha \cap V_\beta$, $h_{\alpha\beta}(x)$ may be written as the class of some $\gamma_\beta(x) \circ \gamma_{\alpha\beta}(y) \circ \gamma_\alpha(x)$, where $y \in T_{i(\alpha)}$ is in the leaf containing

$h_{\alpha\beta}(x)$, and the paths $\gamma_\alpha(x), \gamma_\beta(x)$ are entirely contained in the plaques of $U_{i(\alpha)}$ and $U_{i(\beta)}$ respectively determined by the end points of $\gamma_{\alpha\beta}(y)$. We define the new $h(x)$ to be $\gamma_{\alpha\beta}(y)$. We leave it to the reader to check that this does define a map $h : W \rightarrow M \times \mathbb{R}^{2k}/\widehat{F}$ which is equivalent to the original h , and with each new $h_{\alpha\alpha}$ an immersion of V_α in a transversal of the foliation $\widehat{F}$, so each new $h_{\alpha\beta}$ is still an immersion in a transversal of the combined foliation $\widehat{F}_s \oplus \widehat{F}_r$.

Remark 5.7. *Given any finite subset $\widehat{\mathcal{V}} \subset \mathcal{V}$, we may alter the map h by sliding images along leaves as above and refining the cover $\mathcal{U}$ to obtain an equivalent map h which satisfies, $U_{i(\alpha)} \cap U_{i(\beta)} = \emptyset$ for $V_\alpha, V_\beta \in \widehat{\mathcal{V}}$, $\alpha \neq \beta$. This will be useful later.*

We now extend the map h so that it is étale. Let ν^g be the normal bundle to the image of TF_W in $\mathbb{R}^{2k}$. In particular, $\nu_x^g = [g_*(TF_{W,x})]^\perp$. Denote by $\pi_W : \nu^g \rightarrow W$ the projection and by $j : W \rightarrow \nu^g$ the zero section. On ν^g , we have the open cover by the sets $\widehat{V}_\alpha = \pi_W^{-1}(V_\alpha)$. The map $\widehat{h} : \nu^g \rightarrow M \times \mathbb{R}^{2k}/\widehat{F}$ is given by the cocycle $\{\widehat{h}_{\alpha\beta}, \widehat{V}_\alpha\}$, where

$$\widehat{h}_{\alpha\beta}(x, \xi) = (\widehat{f}_{\alpha\beta}(x), g(x) + \xi).$$

We may choose an open neighborhood X of the zero section $j(W) \subset \nu^g$ so that each $\widehat{h}_{\alpha\beta}$ satisfies

$$\widehat{h}_{\alpha\beta}(\widehat{V}_\alpha \cap \widehat{V}_\beta \cap X) \subset (U_{i(\alpha)}, \gamma_{\alpha\beta}, U_{i(\beta)}),$$

and $\widehat{h}_{\alpha\beta}$ is a diffeomorphism of $\widehat{V}_\alpha \cap \widehat{V}_\beta \cap X$ onto an open subset of the transversal $T_{i(\alpha)i(\beta)}$ of $(U_{i(\alpha)}, \gamma_{\alpha\beta}, U_{i(\beta)})$. Set $h^X = \{\widehat{h}_{\alpha\beta}, \widehat{V}_\alpha \cap X\}$. Then $h^X : X \rightarrow M \times \mathbb{R}^{2k}/\widehat{F}$ is an étale map, and $h = h^X \circ j$.

The map $j : W \rightarrow X$ is K-oriented. To see this, denote by F_W the foliation of W induced by $\widehat{f}$ and F , and denote its normal bundle by ν_{F_W} . Then $TW \oplus j^*TX \simeq TW \oplus TW \oplus \nu^g \simeq (p \circ \pi)^*(TN \oplus \xi) \oplus \nu_{F_W} \oplus \nu_{F_W} \oplus TF_W \oplus \nu^g$. As $TN \oplus \xi$ and $TF_W \oplus \nu^g$ are trivial and $\nu_{F_W} \oplus \nu_{F_W}$ has a complex structure, so is Spin^c , j is K-oriented. Thus we have the push forward map $j_! : K_c^0(W) \rightarrow K_c^0(X)$.

We need two more constructions in order to complete the definition of $\widehat{f}_!$. The first is the non-commutative space $\mathcal{R}$ associated to the open cover $\mathcal{V}_X = \{\widehat{V}_\alpha \cap X\}$ of X . See [Con94] Section II.2.α. Consider the space $\mathcal{X}$ which is the disjoint union of all the $\widehat{V}_\alpha \cap X$. There is an obvious map $q : \mathcal{X} \rightarrow X$, and we write $z \sim z'$ for $z, z' \in \mathcal{X}$ if $q(z) = q(z')$. This is an equivalence relation and we define $\mathcal{R} \subset \mathcal{X} \times \mathcal{X}$ to be the graph of this relation with the induced topology. Note that $\mathcal{R}$ is the disjoint union of the open sets $\mathcal{R}_{\alpha\beta} = \{(z, z') \in (\widehat{V}_\alpha \cap X) \times (\widehat{V}_\beta \cap X) \mid z \sim z'\} \simeq \widehat{V}_\alpha \cap \widehat{V}_\beta \cap X$. The space of compactly supported smooth functions $C_c^\infty(\mathcal{R})$ on $\mathcal{R}$ is a $*$ algebra with the operations being given by

$$(f * g)(z, z'') = \sum_{z \sim z' \sim z''} f(z, z')g(z', z'')$$

$$f^*(z, z') = \overline{f(z', z)}.$$

The C^* algebra associated to this $*$ algebra is Morita equivalent to the C^* algebra associated to $C_c^\infty(X)$, see Proposition 1 of [Con94]. This equivalence is effected by the $*$ -algebra homomorphism $\Upsilon : C_c^\infty(X) \rightarrow C_c^\infty(\mathcal{R})$, given by

$$\Upsilon(f) = (f \circ q) \cdot a,$$

where $a \in C_c^\infty(\mathcal{R})$ is constructed as follows. Choose a partition of unity $\{\phi_\alpha\}$ on X subordinate to the cover $\mathcal{V}_X$. The value of a at the point $(z, z') \in \mathcal{R}_{\alpha\beta}$ is

$$a(z, z') = \sqrt{\phi_\alpha(q(z))\phi_\beta(q(z'))}.$$

Note that a is an idempotent, i.e. $a = a^2 = a^*$, and it has rank one when thought of as a matrix of functions $[a_{\alpha\beta}]$ where $a_{\alpha\beta} \in C_c^\infty(\mathcal{R}_{\alpha\beta})$. Υ induces the isomorphism

$$\Upsilon_* : K_c^*(X) = K_0(C_c^\infty(X)) \rightarrow K_0(C_c^\infty(\mathcal{R})).$$

The second construction is the sub-groupoid $\mathcal{G}_T \subset \mathcal{G}^{\mathbb{R}^{2k}}$, which consists of the classes all leafwise paths which start in the transversal $T_i \subset U_i$ and end in the transversal $T_j \subset U_j$, where U_i, U_j run over all elements

of $\mathcal{U}$. $\mathcal{G}_T$ is the union of all the transversals T_{ij} of the sets (U_i, γ_{ij}, U_j) . The algebra structure on $C_c^\infty(\mathcal{G}_T)$ is given by

$$(f * g)(\gamma) = \sum_{\gamma_1 \gamma_0 = \gamma} f(\gamma_1)g(\gamma_0)$$

$$f^*(\gamma) = \overline{f(\gamma^{-1})}.$$

As above the C^* -algebras associated to $C_c^\infty(\mathcal{G}_T)$ and $C_c^\infty(\mathcal{G}^{\mathbb{R}^{2k}})$ are Morita equivalent, where the equivalence is effected as follows. For each transversal T_i , let $\varrho : D_i \rightarrow T_i$ be an imbedded normal disc bundle. We may assume that the D_i are pairwise disjoint. Let $\ell = \dim \widehat{F}$, and choose a smooth function $\psi : \mathbb{D}^\ell \rightarrow \mathbb{R}$ with compact support so that

$$\int_{\mathbb{D}^\ell} \psi^2 = 1.$$

On each $D_i \simeq T_i \times \mathbb{D}^\ell$ with coordinates (x, y) , define $\psi_i(x, y) = \psi(y)$. Then ψ_i has fiber compact support and for all $x \in T_i$,

$$\int_{\varrho^{-1}(x)} \psi_i^2 = 1.$$

Define $\Psi : C_c^\infty(\mathcal{G}_T) \rightarrow C_c^\infty(\mathcal{G}^{\mathbb{R}^{2k}})$ by

$$\Psi(f)(\gamma) = \psi_i(s(\gamma))f(\gamma_{ij}(y))\psi_j(r(\gamma)),$$

where $\gamma \in (U_i, \gamma_{ij}, U_j)$, $s(\gamma) \in D_i$, $r(\gamma) \in D_j$, and $s(\gamma)$ is in the plaque determined by $y \in T_i$. For any γ which does not start in some D_i or does not end in some D_j , $\Psi(f)(\gamma) = 0$. It is easy to check that Ψ is a well defined $*$ -algebra homomorphism, so it induces the map

$$\Psi_* : K_0(C_c^\infty(\mathcal{G}_T)) \rightarrow K_0(C_c^\infty(\mathcal{G}^{\mathbb{R}^{2k}})).$$

Now it is quite clear that $h^X : X \rightarrow M \times \mathbb{R}^{2k}/\widehat{F}$ induces a well defined map $\widehat{h}^X : \mathcal{R} \rightarrow \mathcal{G}_T$ and also a well defined $*$ -algebra map $\widehat{h}_*^X : C_c^\infty(\mathcal{R}) \rightarrow C_c^\infty(\mathcal{G}_T)$, and so the map $\widehat{h}_!^X : K_0(C_c^\infty(\mathcal{R})) \rightarrow K_0(C_c^\infty(\mathcal{G}_T))$. Recall that $C^*(M, F)$ and $C^*(M \times \mathbb{R}^{2k}, \widehat{F})$ are Connes' foliation C^* algebras for F and $\widehat{F}$ respectively. As $\mathcal{G}^{\mathbb{R}^{2k}} = \mathcal{G} \times \mathbb{R}^{2k}$, we have $K_0(C^*(M \times \mathbb{R}^{2k}, \widehat{F})) \simeq K_0(C^*(M, F) \widehat{\otimes} C_0(\mathbb{R}^{2k}))$ which by Bott periodicity is isomorphic to $K_0(C^*(M, F))$. Denote the composition of these two isomorphisms by $B : K_0(C^*(M \times \mathbb{R}^{2k}, \widehat{F})) \rightarrow K_0(C^*(M, F))$.

Definition 5.8. *The push forward map $f_! : K_c^0(N) \rightarrow K_0(C^*(M, F))$ is the composition of the following maps*

$$K_c^0(N) \xrightarrow{f_!^{\mathbb{R}^{2k}}} K_0(C_c^\infty(\mathcal{G}^{\mathbb{R}^{2k}})) \rightarrow K_0(C^*(M \times \mathbb{R}^{2k}, \widehat{F})) \xrightarrow{B} K_0(C^*(M, F)),$$

where $K_0(C_c^\infty(\mathcal{G}^{\mathbb{R}^{2k}})) \rightarrow K_0(C^*(M \times \mathbb{R}^{2k}, \widehat{F}))$ is induced by the inclusion $C_c^\infty(\mathcal{G}^{\mathbb{R}^{2k}}) \hookrightarrow C^*(M \times \mathbb{R}^{2k}, \widehat{F})$, and $f_!^{\mathbb{R}^{2k}}$ is the composite map

$$K_c^0(N) \xrightarrow{i_!} K_c^0(W) \xrightarrow{j_!} K_c^0(X) \xrightarrow{\Upsilon_*} K_0(C_c^\infty(\mathcal{R})) \xrightarrow{\widehat{h}_!^X} K_0(C_c^\infty(\mathcal{G}_T)) \xrightarrow{\Psi_*} K_0(C_c^\infty(\mathcal{G}^{\mathbb{R}^{2k}})).$$

The following result is proved in [CS84].

Theorem 5.9. *The map $f_! : K_c^*(N) \rightarrow K_*(C^*(M, F))$ is well defined.*

Proof. This is a consequence of the homotopy invariance of K -theory for C^* -algebras by using the homotopy described in [ASI][page 498]. For more details, see [CS84]. $\square$

Definition 5.10. *Let $f : N \rightarrow M/F$ be a K -oriented map and $g : W \hookrightarrow \mathbb{R}^{2k}$ an imbedding of W as above. Set*

$$\text{ch}_a^{\mathbb{R}^{2k}} = \int_{\mathbb{R}^{2k}} \circ \text{ch}_a : K_0(C_c^\infty(\mathcal{G}^{\mathbb{R}^{2k}})) \rightarrow H_c^*(M/F),$$

where $\text{ch}_a : K_0(C_c^\infty(\mathcal{G}^{\mathbb{R}^{2k}})) \rightarrow H_c^*(M \times \mathbb{R}^{2k}/\widehat{F})$ is the Chern character associated to the foliation $\widehat{F}$ on $M \times \mathbb{R}^{2k}$, and $\int_{\mathbb{R}^{2k}} : H_c^*(M \times \mathbb{R}^{2k}/\widehat{F}) \simeq H_c^*(M/F) \otimes H_c^*(\mathbb{R}^{2k}; \mathbb{R}) \rightarrow H_c^*(M/F)$ is integration over $\mathbb{R}^{2k}$.

The fact that $\text{ch}_a^{\mathbb{R}^{2k}} \circ f_!^{\mathbb{R}^{2k}} : K_c^0(N) \rightarrow H_c^*(M/F)$ is well defined is an immediate consequence of our main theorem.

Theorem 5.11. *Let F be an oriented foliation of a compact manifold M , and $f : N \rightarrow M/F$ a K -oriented map. Then the following diagram commutes.*

$$\begin{array}{ccc} K_c^0(N) & \xrightarrow{f_!^{\mathbb{R}^{2k}}} & K_0(C_c^\infty(\mathcal{G}^{\mathbb{R}^{2k}})) \\ \text{ch}(\cdot) \wedge \text{Td}(f) \downarrow & & \downarrow \text{ch}_a^{\mathbb{R}^{2k}} \\ H_c^*(N; \mathbb{R}) & \xrightarrow{f_!} & H_c^*(M/F). \end{array}$$

Proof. We need only show that the following diagram commutes.

$$\begin{array}{ccccccc} K_c^0(N) & \xrightarrow{i_!} & K_c^0(W) & \xrightarrow{j_!} & K_c^0(X) & \xrightarrow{h_!^X} & K_0(C_c^\infty(\mathcal{G}^{\mathbb{R}^{2k}})) \\ \downarrow \text{ch}(\cdot) \wedge \text{Td}(f) & & \downarrow \text{ch}(\cdot) \wedge \text{Td}(j) & & \downarrow \text{ch} & & \downarrow \text{ch}_a \\ H_c^*(N; \mathbb{R}) & \xrightarrow{i_!} & H_c^*(W; \mathbb{R}) & \xrightarrow{j_!} & H_c^*(X; \mathbb{R}) & \xrightarrow{h_!^X} & H_c^*(M \times \mathbb{R}^{2k}/\widehat{F}), \end{array}$$

where $h_!^X : K_c^0(X) \rightarrow K_0(C_c^\infty(\mathcal{G}^{\mathbb{R}^{2k}}))$ is the composition

$$K_c^0(X) \xrightarrow{\Upsilon_*} K_0(C_c^\infty(\mathcal{R})) \xrightarrow{\widehat{h_!^X}} K_0(C_c^\infty(\mathcal{G}_T)) \xrightarrow{\Psi_*} K_0(C_c^\infty(\mathcal{G}^{\mathbb{R}^{2k}})),$$

$\text{Td}(f) = \text{Td}(TN)/\text{Td}(f^*\nu)$, and $\text{Td}(j) = \text{Td}(TW)/\text{Td}(j^*TX)$. To see this, observe that

$$f_! = \int_{\mathbb{R}^{2k}} \circ h_!^X \circ j_! \circ i_! : H_c^*(N; \mathbb{R}) \rightarrow H_c^*(M/F).$$

The classical results of [ASIII] and [CS84] imply that the second rectangle commutes up to sign. The sign comes from the definition of the Chern character, see again [ASIII]. By the same token, the left rectangle would commute if we removed the term $\wedge \text{Td}(j)$ and replaced $\text{Td}(f)$ by $\text{Td}(i) = \text{Td}(TN)/\text{Td}(i^*TW)$. Thus to see that it does commute, we need to show that $\text{Td}(f) = \text{Td}(i) \wedge i^*(\text{Td}(j))$. Let $\rho : M \times \mathbb{R}^{2k} \rightarrow M$ be the projection and denote the normal bundle of $\widehat{F}$ by $\nu_{\widehat{F}}$. Then

$$f = \rho \circ h^X \circ j \circ i,$$

so $f^*\nu = i^*j^*h^{X*}\rho^*\nu$ and $f^*(\nu \oplus \mathbb{R}^{2k}) = i^*j^*h^{X*}\rho^*(\nu \oplus \mathbb{R}^{2k}) = i^*j^*h^{X*}(\nu_{\widehat{F}}) = i^*j^*(TX)$, since h^X is étale. Thus $\text{Td}(f^*\nu) = \text{Td}(f^*(\nu \oplus \mathbb{R}^{2k})) = \text{Td}(i^*j^*TX)$, so

$$\text{Td}(f) = \text{Td}(TN)/\text{Td}(i^*j^*TX).$$

As $\text{Td}(i) = \text{Td}(TN)/\text{Td}(i^*TW)$ and $\text{Td}(j) = \text{Td}(TW)/\text{Td}(j^*TX)$,

$$\text{Td}(i) \wedge i^*(\text{Td}(j)) = \text{Td}(TN)/\text{Td}(i^*j^*TX),$$

and the left rectangle commutes.

To finish the proof we show that the third rectangle commutes by a direct computation. Let $\widehat{e} = (e, \lambda)$ be an idempotent in $M_N(C_c^\infty(X) \oplus \mathbb{C})$. We now apply Remark 5.7 to the open sets $\mathcal{V}_X$ covering the support of e to obtain a representative of the map h which satisfies the conclusions of that remark. In particular, this implies that the image of e on the transversal $T_{i(\alpha)}$ under the composition $h^X \circ \text{ch}$ is the Haefliger form for $\widehat{F}$

$$h_{\alpha*}^X(\phi_\alpha \text{tr}(e \exp\left(\frac{-(de)^2}{2i\pi}\right))),$$

where d is the deRham differential on X . The degree $2j$ part of this form is

$$\left(\frac{-1}{2i\pi}\right)^j \frac{1}{j!} h_{\alpha*}^X(\phi_\alpha \operatorname{tr}(e(de)^{2j})).$$

The map $h_\alpha^X : \widehat{V}_\alpha \cap X \rightarrow T_{i(\alpha)}$ is a diffeomorphism onto its image, so we may assume that we have coordinates (x_0, s_0) on $D_{i(\alpha)} \subset U_{i(\alpha)} \subset M \times \mathbb{R}^{2k}$ of the form $(\widehat{V}_\alpha \cap X) \times \mathbb{D}^\ell$, and the transversal coordinate x_0 is given by $(h_\alpha^X)^{-1}$. Then the degree $2j$ part of $h^X \operatorname{ch} e$ in these coordinates at the point $(x_0, 0) \in T_{i(\alpha)}$ is just

$$(h^X \operatorname{ch} e)_{2j} = \left(\frac{-1}{2i\pi}\right)^j \frac{1}{j!} \phi_\alpha \operatorname{tr}(e(de)^{2j})(x_0),$$

where d is now the deRham differential on $T_{i(\alpha)}$.

We now compute the degree $2j$ part of $\operatorname{ch}_a h^X e$ on $T_{i(\alpha)}$ in these coordinates and show that it is the same as $(h^X \operatorname{ch} e)_{2j}$. As above, we have coordinates $(\widehat{V}_\alpha \cap X) \times \mathbb{D}^\ell \times \mathbb{D}^\ell$ on the open subset of the chart $(U_{i(\alpha)}, \gamma_{\alpha\beta}, U_{i(\beta)})$ of $\mathcal{G}^{\mathbb{R}^{2k}}$ determined by $D_{i(\alpha)}$ and $D_{i(\beta)}$. The local coordinate functions are denoted (x_0, s_0, r_0) , where s_0 is the fiber coordinate on $D_{i(\alpha)}$ and r_0 the fiber coordinate on $D_{i(\beta)}$. Note that $h^X e$ can be non-zero on the chart $(U_{i(\alpha)}, \gamma_{\alpha\beta}, U_{i(\beta)})$ only if $\widehat{V}_\alpha \cap \widehat{V}_\beta \cap X \neq \emptyset$, and it has support contained in the open subset $(\widehat{V}_\alpha \cap X) \times \mathbb{D}^\ell \times \mathbb{D}^\ell$. The element $h^X e$ in these coordinates is given by

$$h^X e = \psi_{i(\beta)}(r_0) \psi_{i(\alpha)}(s_0) (e\sqrt{\phi_\beta \phi_\alpha})(x_0).$$

If we ignore the constant $\left(\frac{-1}{2i\pi}\right)^j \frac{1}{j!}$, the degree $2j$ part of Haeffliger form $\operatorname{ch}_a h^X e$ on the transversal $T_{i(\alpha)}$ is

$$(\operatorname{ch}_a h^X e)_{2j} = \int_{\mathbb{D}^\ell} \operatorname{tr}_2 \left(h^X e (\delta(h^X e))^{2j} \right) ds_0,$$

where tr_2 takes into account the 2×2 matrix $\delta(h^X e)$ obtained from the definition of δ . An easy induction argument using the fact that $h^X e (\partial_\nu(h^X e))^{2\ell-1} h^X e = 0$ for all $\ell > 0$, shows that

$$h^X e (\delta(h^X e))^{2j} = \begin{pmatrix} h^X e ((\partial_\nu(h^X e))^2 + (h^X e)\theta(h^X e))^j & 0 \\ 0 & 0 \end{pmatrix}.$$

Thus

$$(\operatorname{ch}_a h^X e)_{2j} = \int_{\mathbb{D}^\ell} \operatorname{tr} \left(h^X e ((\partial_\nu(h^X e))^2 + (h^X e)\theta(h^X e))^j \right) ds_0.$$

We will show below that we can choose the connection we use to define δ so that $\theta = 0$, so

$$(\operatorname{ch}_a h^X e)_{2j} = \int_{\mathbb{D}^\ell} \operatorname{tr} \left(h^X e (\partial_\nu(h^X e))^{2j} \right) ds_0.$$

At the point $(x_0, 0) \in T_{i(\alpha)}$, this is given by

$$\int_{\mathbb{D}^\ell} \operatorname{tr} \left(\int_{\tilde{L}_{s(\gamma_{2j})}} \cdots \int_{\tilde{L}_{s(\gamma_1)}} (h^X e)(\gamma_{2j}) \partial_\nu(h^X e)(\gamma_{2j-1}) \cdots \partial_\nu(h^X e)(\gamma_0) d\gamma_1 \cdots d\gamma_{2j} \right) ds_0,$$

where $s(\gamma_0) = (x_0, s_0) \in (\widehat{V}_\alpha \cap X) \times \mathbb{D}^\ell \subset U_{i(\alpha)}$, and $\gamma_{2j} \cdots \gamma_0$ is equivalent to the constant path at the point (x_0, s_0) . The element $\gamma_0 \in (U_{i(\alpha)}, \gamma_{\alpha\beta}, U_{i(\beta_1)})$, and for $a = 1, \dots, 2j$, $\gamma_a \in (U_{i(\beta_a)}, \gamma_{\beta_a \beta_{a+1}}, U_{i(\beta_{a+1})})$, where we have the convention $U_{i(\beta_{2j+1})} = U_{i(\alpha)}$. $(U_{i(\beta_a)}, \gamma_{\beta_a \beta_{a+1}}, U_{i(\beta_{a+1})})$ has coordinates (x_a, s_a, r_a) from $(\widehat{V}_{\beta_a} \cap X) \times \mathbb{D}^\ell \times \mathbb{D}^\ell$. In these coordinates $h^X e$ is given by

$$h^X e|_{(U_{i(\beta_a)}, \gamma_{\beta_a \beta_{a+1}}, U_{i(\beta_{a+1})})} = \psi_{i(\beta_{a+1})}(r_a) \psi_{i(\beta_a)}(s_a) (e\sqrt{\phi_{\beta_{a+1}} \phi_{\beta_a}})(q(x_a)).$$

A straightforward computation shows that the above integral is the same as

$$\int_{\mathbb{D}^\ell} \operatorname{tr} \left(\sum_{\beta_1 \cdots \beta_{2j}} \int_{\mathbb{D}^\ell} \cdots \int_{\mathbb{D}^\ell} \psi_{i(\alpha)}(r_{2j}) \psi_{i(\beta_{2j})}(s_{2j}) (e\sqrt{\phi_\alpha \phi_{\beta_{2j}}})(q(x_{2j})) \right).$$

$$\partial_\nu \left\{ \psi_{i(\beta_{2j})}(r_{2j-1}) \psi_{i(\beta_{2j-1})}(s_{2j-1}) (e\sqrt{\phi_{\beta_{2j}}\phi_{\beta_{2j-1}}})(q(x_{2j-1})) \right\} \wedge \cdots \wedge \\ \partial_\nu \left\{ \psi_{i(\beta_1)}(r_0) \psi_{i(\alpha)}(s_0) (e\sqrt{\phi_{\beta_1}\phi_\alpha})(q(x_0)) \right\} ds_1 \cdots ds_{2j} \Big) ds_0.$$

The sum is over all β_a with $x_0 \in q(\widehat{V}_{\beta_a} \cap X)$. Note that $q(x_a) = x_0$ and $r_a = s_{a+1}$ for $a = 0, \dots, 2j$, where we set $s_{2j+1} = s_0$. Thus the integral is

$$\int_{\mathbb{D}^\ell} \text{tr} \left(\sum_{\beta_1 \cdots \beta_{2j}} \int_{\mathbb{D}^\ell} \cdots \int_{\mathbb{D}^\ell} \psi_{i(\alpha)}(s_0) \psi_{i(\beta_{2j})}(s_{2j}) (e\sqrt{\phi_\alpha \phi_{\beta_{2j}}})(x_0) \cdot \right. \\ \left. \partial_\nu \left\{ \psi_{i(\beta_{2j})}(s_{2j}) \psi_{i(\beta_{2j-1})}(s_{2j-1}) (e\sqrt{\phi_{\beta_{2j}}\phi_{\beta_{2j-1}}})(x_0) \right\} \wedge \cdots \wedge \right. \\ \left. \partial_\nu \left\{ \psi_{i(\beta_1)}(s_1) \psi_{i(\alpha)}(s_0) (e\sqrt{\phi_{\beta_1}\phi_\alpha})(x_0) \right\} ds_1 \cdots ds_{2j} \right) ds_0.$$

Now, the computation is local and in order to get the local expression for the transverse differential ∂_ν on $(U_{i(\beta_a)}, \gamma_{\beta_a\beta_{a+1}}, U_{i(\beta_{a+1})})$ we apply the material from the proof of Lemma 3.2 where E_1 is the N dimensional trivial bundle over $M \times \mathbb{R}^{2k}$, since $h^X e \in M_N(C_c^\infty(\mathcal{G}^{\mathbb{R}^{2k}}))$. We choose the connection on E_1 to be the canonical flat connection. Then ∇^ν on $(U_{i(\beta_a)}, \gamma_{\beta_a\beta_{a+1}}, U_{i(\beta_{a+1})})$ is given by

$$\nabla^\nu = d_{x_a} + d_{s_a},$$

and $\theta = (\nabla^\nu)^2 = 0$ as promised above. Locally

$$\partial_\nu(h^X e) = (d_{x_a} + d_{s_a})(h^X e).$$

Using these facts, the fact that we may identify each d_{x_a} with d on $T_{i(\alpha)}$, as well as the fact that $\int_{\mathbb{D}^\ell} \psi_i d_{s_i} \psi_i = 0$, the integral becomes

$$\int_{\mathbb{D}^\ell} \text{tr} \left(\sum_{\beta_1 \cdots \beta_{2j}} \int_{\mathbb{D}^\ell} \cdots \int_{\mathbb{D}^\ell} \psi_{i(\alpha)}(s_0) \psi_{i(\beta_{2j})}(s_{2j}) (e\sqrt{\phi_\alpha \phi_{\beta_{2j}}})(x_0) \cdot \right. \\ \left\{ \psi_{i(\beta_{2j})}(s_{2j}) \psi_{i(\beta_{2j-1})}(s_{2j-1}) d(e\sqrt{\phi_{\beta_{2j}}\phi_{\beta_{2j-1}}})(x_0) \right\} \wedge \cdots \wedge \\ \left. \left\{ \psi_{i(\beta_1)}(s_1) \psi_{i(\alpha)}(s_0) d(e\sqrt{\phi_{\beta_1}\phi_\alpha})(x_0) \right\} ds_1 \cdots ds_{2j} \right) ds_0.$$

Since $\int_{\mathbb{D}^\ell} \psi_i^2 = 1$, this equals

$$\text{tr} \left(\sum_{\beta_1 \cdots \beta_{2j}} e\sqrt{\phi_\alpha \phi_{\beta_{2j}}} d \left(e\sqrt{\phi_{\beta_{2j}}\phi_{\beta_{2j-1}}} \right) \wedge \cdots \wedge d \left(e\sqrt{\phi_{\beta_1}\phi_\alpha} \right) \right) (x_0),$$

which equals

$$\text{tr} \left(\sum_{\beta_1 \cdots \beta_{2j}} \phi_\alpha \phi_{\beta_{2j}} \cdots \phi_{\beta_1} e d e \wedge \cdots \wedge d e \right) (x_0) + \text{other terms}.$$

As $\sum_\beta \phi_\beta = 1$ the first term above is just

$$\phi_\alpha \text{tr}(e(de)^{2j})(x_0),$$

so we need only show that the “other terms” are zero in Haefliger cohomology. These terms come in several types.

(1) Those which have a repeated $d\phi_{\beta_i}$, such as

$$\text{tr} \left(\sum_{\beta_1 \cdots \beta_{2j}} e\sqrt{\phi_\alpha} \frac{d\phi_{\beta_{2j}}}{2\sqrt{\phi_{\beta_{2j}}}} \wedge e \frac{d\phi_{\beta_{2j}}}{2\sqrt{\phi_{\beta_{2j}}}} \sqrt{\phi_{\beta_{2j-1}}} \wedge \cdots \wedge d(e\sqrt{\phi_{\beta_1}\phi_\alpha}) \right),$$

which are obviously zero as forms.

- (2) Those which have a $d\phi_{\beta_i}$ which is not repeated, such as

$$\mathrm{tr} \left(\sum_{\beta_1 \cdots \beta_{2j}} e\sqrt{\phi_\alpha} \frac{1}{2} d\phi_{\beta_{2j}} \wedge e\frac{1}{2} d\phi_{\beta_{2j-1}} \wedge \cdots \wedge d(e\sqrt{\phi_{\beta_1} \phi_\alpha}) \right).$$

These are zero as forms since they equal

$$\mathrm{tr} \left(\sum_{\beta_1 \cdots \beta_{2j-1}} e\sqrt{\phi_\alpha} \frac{1}{2} d \left(\sum_{\beta_{2j}} \phi_{\beta_{2j}} \right) \wedge e\frac{1}{2} d\phi_{\beta_{2j-1}} \wedge \cdots \wedge d(e\sqrt{\phi_{\beta_1} \phi_\alpha}) \right),$$

and $\sum_{\beta_{2j}} \phi_{\beta_{2j}} = 1$ near x_0 .

- (3) Those with no $d\phi_{\beta_i}$, in which case we have the form

$$\mathrm{tr} \left(\sum_{\beta_1 \cdots \beta_{2j}} e\sqrt{\phi_{\beta_{2j}}} de\sqrt{\phi_{\beta_{2j}} \phi_{\beta_{2j-1}}} \wedge \cdots \wedge de\sqrt{\phi_{\beta_2} \phi_{\beta_1}} \wedge e\sqrt{\phi_{\beta_1}} \wedge \frac{1}{2} d\phi_\alpha \right).$$

The Haefliger class determined by all these local forms is zero since it is the local image under h^X of the global form on X

$$\mathrm{tr} \left(\sum_{\beta_1 \cdots \beta_{2j}} e\sqrt{\phi_{\beta_{2j}}} de\sqrt{\phi_{\beta_{2j}} \phi_{\beta_{2j-1}}} \wedge \cdots \wedge de\sqrt{\phi_{\beta_2} \phi_{\beta_1}} \wedge e\sqrt{\phi_{\beta_1}} \wedge \frac{1}{2} d \left(\sum_{\alpha} \phi_\alpha \right) \right),$$

which is zero since $\sum_{\alpha} \phi_\alpha = 1$.

□

Remark 5.12. As pointed out to us by G. Skandalis, the results of [Con86] hold for any holonomy invariant current. Therefore, the pairing of the Chern-Connes character ch_a with any holonomy invariant current extends to the K -theory of the foliation C^* -algebra. Hence, the proof above shows using the Connes-Skandalis index theorem in the K -theory of the foliation C^* -algebra that $\mathrm{ch}_a^{\mathbb{R}^{2k}} \circ f_!^{\mathbb{R}^{2k}}$ is given by

$$\mathrm{ch}_a^{\mathbb{R}^{2k}} \circ f_!^{\mathbb{R}^{2k}} = \mathrm{ch}_a \circ f_!^{an},$$

where $f_!^{an}$ is the analytic shriek map described in [CS84].

6. THE INDEX THEOREM

In this section, we compute the Chern character of the topological index of any class $u \in K_c^0(TF)$. This permits us to translate the Connes-Skandalis index theorem [CS84] to Haefliger cohomology and yields an alternative proof of the longitudinal index theorem in Haefliger cohomology for all longitudinal pseudodifferential operators, compare [HL99, GL03, GL05].

We first describe the Chern character of the topological index, i.e. the homomorphism $\mathrm{ch}_a(\mathrm{Ind}_t) : K_c^0(TF) \rightarrow H_c^*(M/F)$. Choose a smooth imbedding $g : M \hookrightarrow \mathbb{R}^{2k}$. Denote by $\pi_F : TF \rightarrow M$ the projection, and by $\pi_N : N \rightarrow M$ the normal bundle to TF in $\mathbb{R}^{2k}$. Let $j : M \hookrightarrow N$ be the zero section of N and $f : TF \rightarrow N$ the composite map

$$f = j \circ \pi_F.$$

The map f is K -oriented by a complex structure since $T(TF) \simeq \pi_F^*(TF \oplus TM)$ and $f^*(TN) \simeq \pi_F^*(N \oplus TM)$, so

$$T(TF) \oplus f^*(TN) \simeq \pi_F^*(TF \oplus N \oplus TM \oplus TM) \simeq \pi_F^*(g^*(T\mathbb{R}^{2k}) \oplus TM \oplus TM).$$

Therefore the Atiyah-Singer shriek map $f_! : K_c^0(TF) \rightarrow K_c^0(N)$ is well defined, see for instance [ASI]. Moreover the Todd class of f is given by

$$\mathrm{Td}(f) = \mathrm{Td}(T(TF)) / \mathrm{Td}(f^*(TN)) = \mathrm{Td}(\pi_F^*(TF \oplus TM)) / \mathrm{Td}(\pi_F^*(N \oplus TM)) =$$

$$\pi_F^*(\mathrm{Td}(TF) / \mathrm{Td}(N)) = \pi_F^*(\mathrm{Td}(TF) \cdot \mathrm{Td}(TF)) = \pi_F^*(\mathrm{Td}(TF \oplus TF)) = \pi_F^*(\mathrm{Td}(TF \otimes \mathbb{C})),$$

since $1 / \mathrm{Td}(N) = \mathrm{Td}(TF)$ as $N \oplus F \simeq M \times \mathbb{R}^{2k}$.

Recall the foliation $\widehat{F}$ of $M \times \mathbb{R}^{2k}$ and its holonomy groupoid $\mathcal{G}^{\mathbb{R}^{2k}} = \mathcal{G} \times \mathbb{R}^{2k}$. The bundle N may be identified with an open transversal of $\widehat{F}$ by the map $i : N \rightarrow M \times \mathbb{R}^{2k}$ where

$$i(x, \xi) = (x, g(x) + \xi).$$

On a suitable neighborhood of the zero section, this gives an open transversal and we may assume that the neighborhood is in fact all of N . Thus we have the map

$$i_! : K_c^0(N) \longrightarrow K_0(C_c^\infty(\mathcal{G}^{\mathbb{R}^{2k}})).$$

The inclusion $\rho^{\mathbb{R}^{2k}} : C_c^\infty(\mathcal{G}^{\mathbb{R}^{2k}}) \hookrightarrow C^*(M \times \mathbb{R}^{2k}, \widehat{F})$ gives $\rho_*^{\mathbb{R}^{2k}} : K_0(C_c^\infty(\mathcal{G}^{\mathbb{R}^{2k}})) \rightarrow K(C^*(M \times \mathbb{R}^{2k}, \widehat{F}))$. Finally, we have the Bott isomorphism $B : K_0(C^*(M \times \mathbb{R}^{2k}, \widehat{F})) \simeq K_0(C^*(M, F) \widehat{\otimes} C_0(\mathbb{R}^{2k})) \rightarrow K_0(C^*(M, F))$.

Definition 6.1. [CS84] *The topological index morphism is the composite map*

$$\text{Ind}_t = B \circ \rho_*^{\mathbb{R}^{2k}} \circ i_! \circ f_! : K_c^0(TF) \longrightarrow K_0(C^*(M, F)).$$

Ind_t is well defined, see [CS84]. Moreover, if $\pi : TF \rightarrow M/F$ is the natural map, then π is a K -oriented submersion and it is clear that $i_! \circ f_! = \pi_!^{\mathbb{R}^{2k}}$. We therefore denote by $\text{ch}_a(\text{Ind}_t)$ the map

$$\text{ch}_a(\text{Ind}_t) = \text{ch}_a^{\mathbb{R}^{2k}} \circ \pi_!^{\mathbb{R}^{2k}} : K_c^0(TF) \longrightarrow H_c^*(M/F).$$

Theorem 6.2. *For any $u \in K_c^0(TF)$, $\text{ch}_a(\text{Ind}_t)(u)$, the algebraic Chern character of the topological index of u , is given by*

$$\text{ch}_a(\text{Ind}_t)(u) = (-1)^p \int_F \pi_{F!}(\text{ch}(u)) \text{Td}(TF \otimes \mathbb{C}) \in H_c^*(M/F),$$

where $\pi_{F!} : H_c^*(TF; \mathbb{R}) \rightarrow H^*(M; \mathbb{R})$ is integration along the fibers, and $\text{ch} : K_c^0(TF) \rightarrow H_c^*(TF)$ is the usual Chern character.

Proof. As $i : N \rightarrow M \times \mathbb{R}^{2k}$ is a transversal to $\widehat{F}$, we may regard i as a map from N to $M \times \mathbb{R}^{2k}/\widehat{F}$ and we have the push forward map

$$i_! : H_c^*(N; \mathbb{R}) \rightarrow H_c^*(M \times \mathbb{R}^{2k}/\widehat{F}).$$

The essence of the theorem is that the following diagram commutes up to sign.

$$\begin{array}{ccccccc} K_c^0(TF) & \xrightarrow{f_!} & K_c^0(N) & \xrightarrow{i_!} & K_0(C_c^\infty(\mathcal{G}^{\mathbb{R}^{2k}})) & & \\ \downarrow \text{ch}(\cdot) \wedge \text{Td}(f) & & \downarrow \text{ch} & & \downarrow \text{ch}_a^{\mathbb{R}^{2k}} & & \\ H_c^*(TF; \mathbb{R}) & \xrightarrow{f_!} & H_c^*(N; \mathbb{R}) & \xrightarrow{i_!} & H_c^*(M \times \mathbb{R}^{2k}/\widehat{F}) & \xrightarrow{\int_{\mathbb{R}^{2k}}} & H_c^*(M/F) \\ \pi_{F!} \searrow & & \nearrow j_! & & \searrow \simeq & & \nearrow \int_{\mathbb{R}^{2k}} \\ & H^*(M; \mathbb{R}) & & & H_c^*(M/F) \otimes H_c^*(\mathbb{R}^{2k}; \mathbb{R}) & & \end{array}$$

The classical results of [ASIII] and [CS84] imply that the left rectangle commutes up to sign. The sign comes from the definition of the Chern character and can be fixed by inspecting trivial examples, see again [ASIII]. The triangles commute by definition. The fact that the rest of the diagram commutes is just Theorem 5.11. Now observe that

$$\int_{\mathbb{R}^{2k}} \circ \simeq \circ i_! \circ j_! = \int_F.$$

The theorem follows immediately since $\text{Td}(f) = \pi_F^* \text{Td}(TF \otimes \mathbb{C})$. $\square$

We now discuss the relationship of the results here with the Connes-Skandalis index theorem, and answer a question asked in [He95]. As in Section 3, let P be a uniformly supported elliptic pseudo-differential $\mathcal{G}$ -operator acting on the bundle $E = r^*(E_1)$. Restricting P to the holonomy cover of a leaf L of F gives a $\mathcal{G}(L)$ -invariant pseudo-differential operator, where $\mathcal{G}(L)$ is the holonomy group of the leaf L . Therefore, the principal symbol of P can be defined using the induced operator on $E_1|_L$. We get in this way a section $\sigma(P)$

of the bundle $\text{Hom}(\pi_F^* E_1)$ over TF . An element $P \in \Psi^\infty(\mathcal{G}; E)$ is elliptic if $\sigma(P)(\xi) : (\pi_F^* E_1)_\xi \rightarrow (\pi_F^* E_1)_\xi$ is an isomorphism for all $\xi \neq 0$, $\xi \in TF$. Denote by $\text{Ind}_a : K_c^0(TF) \rightarrow K_0(C_c^\infty(\mathcal{G}))$ Connes-Skandalis analytic index map [CS84]. This map associates to an element u the analytic index $\text{Ind}_a(P)$ of any elliptic pseudo-differential $\mathcal{G}$ -operator P with symbol $\sigma(P) = u$. The construction of P from u is given in [Con79]. Denote by

$$\rho_* : K_0(C_c^\infty(\mathcal{G})) \rightarrow K_0(C^*(M, F)),$$

the homomorphism induced by the inclusion map $\rho : C_c^\infty(\mathcal{G}) \hookrightarrow C^*(M, F)$. Note that ρ_* need not be injective. Recall the Connes-Skandalis leafwise index theorem.

Theorem 6.3. [CS84] $\rho_* \circ \text{Ind}_a = \text{Ind}_t : K_c^0(TF) \rightarrow K_0(C^*(M, F))$.

Let $\mathcal{A}$ be any $*$ -subalgebra of $C^*(M \times \mathbb{R}^{2k}, \widehat{F})$ which contains the smooth convolution algebra $C_c^\infty(\mathcal{G}^{\mathbb{R}^{2k}})$, and denote by $\rho^{\mathcal{A}} : \mathcal{A} \rightarrow C^*(M \times \mathbb{R}^{2k}, \widehat{F})$ the inclusion. An example of such algebra is the algebra of operators in $C^*(M \times \mathbb{R}^{2k}, \widehat{F})$ with smooth kernels having superexponential decay, see [HL02]. In that case, the Chern character $\text{ch}_a^{\mathbb{R}^{2k}}$ extends to the K -theory of $\mathcal{A}$.

Theorem 6.4. Suppose that the Chern character $\text{ch}_a^{\mathbb{R}^{2k}}$ extends to $K(\mathcal{A})$ and that $\text{Ker}(\rho_*^{\mathcal{A}}) \subset \text{Ker}(\text{ch}_a^{\mathbb{R}^{2k}})$. Then for all $u \in K_c^0(TF)$,

$$\text{ch}_a(\text{Ind}_t)(u) = \text{ch}_a(\text{Ind}_a(u)).$$

Proof. The Connes-Skandalis index theorem gives

$$B^{-1} \circ \rho_* \circ \text{Ind}_a = \rho_*^{\mathbb{R}^{2k}} \circ \pi_!^{\mathbb{R}^{2k}}.$$

Denote by $\varphi : K_0(C_c^\infty(\mathcal{G})) \rightarrow K_0(C_c^\infty(\mathcal{G}^{\mathbb{R}^{2k}}))$ the Bott homomorphism. Then

$$B^{-1} \circ \rho_* = \rho_*^{\mathbb{R}^{2k}} \circ \varphi.$$

Denote by $i_*^{\mathbb{R}^{2k}} : K_0(C_c^\infty(\mathcal{G}^{\mathbb{R}^{2k}})) \rightarrow K_0(\mathcal{A})$ the map induced by the inclusion, and set

$$\text{Ind}_a^{\mathcal{A}}(u) = (i_*^{\mathbb{R}^{2k}} \circ \varphi)(\text{Ind}_a(u)) \quad \text{and} \quad \text{Ind}_t^{\mathcal{A}}(u) = (i_*^{\mathbb{R}^{2k}} \circ \pi_!^{\mathbb{R}^{2k}})(u).$$

These satisfy

$$\rho_*^{\mathcal{A}}(\text{Ind}_t^{\mathcal{A}}(u)) = \rho_*^{\mathcal{A}}(\text{Ind}_a^{\mathcal{A}}(u)),$$

so

$$\text{Ind}_a^{\mathcal{A}}(u) - \text{Ind}_t^{\mathcal{A}}(u) \in \text{Ker}(\rho_*^{\mathcal{A}}),$$

and

$$\text{ch}_a^{\mathbb{R}^{2k}}(\text{Ind}_t^{\mathcal{A}}(u)) = \text{ch}_a^{\mathbb{R}^{2k}}(\text{Ind}_a^{\mathcal{A}}(u)) \in H_c^*(M \times \mathbb{R}^{2k}/\widehat{F}).$$

Integrating over $\mathbb{R}^{2k}$, we get by definition of the Chern character of the topological index

$$\text{ch}_a(\text{Ind}_t)(u) = \int_{\mathbb{R}^{2k}} \text{ch}_a^{\mathbb{R}^{2k}}(\text{Ind}_t^{\mathcal{A}}(u)) = \int_{\mathbb{R}^{2k}} \text{ch}_a^{\mathbb{R}^{2k}}(\text{Ind}_a^{\mathcal{A}}(u)) = \text{ch}_a(\text{Ind}_a(u)).$$

□

From the deep extension theorem of [Con86], it follows that $\text{Ker}(\rho_*^{\mathbb{R}^{2k}}) \subset \text{Ker}(\text{ch}_a^{\mathbb{R}^{2k}})$ i.e. Theorem 6.4 holds for the case $\mathcal{A} = C_c^\infty(\mathcal{G}^{\mathbb{R}^{2k}})$. Therefore, we deduce from our computation of the Chern character of the topological index, the cohomological formula for all elliptic pseudodifferential operators on foliations.

Corollary 6.5. For any $u \in K_c^0(TF)$, $\text{ch}_a(\text{Ind}_t)(u) = \text{ch}_a(\text{Ind}_a(u))$ and

$$\text{ch}_a(\text{Ind}_a(u)) = (-1)^p \int_F \pi_{F!}(\text{ch}(u)) \text{Td}(TF \otimes \mathbb{C}) \in H_c^*(M/F).$$

Thus we have the following.

Theorem 6.6. *Suppose that P is a leafwise elliptic pseudodifferential operator defined along the leaves of a foliation. Then*

$$\mathrm{ch}_a(\mathrm{Ind}_t)([\sigma(P)]) = \mathrm{ch}_a(\mathrm{Ind}_a(P))$$

in Haefliger cohomology.

A particularly interesting case is that of spin^c Dirac operators. Compare with [GL03, GL05]:

Corollary 6.7. *Suppose TF admits a Spin^c structure and denote by L the line bundle corresponding to that structure. Let $\alpha : K^0(M) \rightarrow K_c^0(TF)$ be the corresponding Thom isomorphism. Then for any $u \in K^0(M)$,*

$$\mathrm{ch}_a(\mathrm{Ind}_a(\alpha(u))) = \int_F \hat{A}(TF) e^{c_1(L)/2} \mathrm{ch}(u) \in H_c^*(M/F),$$

where $c_1(L)$ is the Chern class of L , $\hat{A}(TF)$ is the $\hat{A}$ -genus of TF , [LM89], and $\mathrm{ch} : K^0(M) \rightarrow H^(M; \mathbb{R})$ is the usual Chern character.*

Proof. The Thom isomorphism α is exactly the Gysin shriek map corresponding to the K -oriented inclusion of the zero section of TF [ASI]. Therefore, by functoriality of the Gysin shriek maps [CS84], the composite map $f_! \circ \alpha$ coincides with the Gysin shriek $j_!$ map corresponding to the inclusion of the zero section $j : M \hookrightarrow N$ of the K -oriented bundle N . Hence

$$\mathrm{ch}_a(\mathrm{Ind}_t)(\alpha(u)) = (-1)^p \int_F \mathrm{Td}(j) \mathrm{ch}(u).$$

Now an inspection of the cohomology class $\mathrm{Td}(j)$ shows that

$$\mathrm{Td}(j) = 1/\mathrm{Td}(N) = \mathrm{Td}(TF),$$

where here Td means the Todd class of the Spin^c bundle, see [LM89, BD81]. But the Spin^c Todd class of the bundle TF is precisely $(-1)^p \hat{A}(TF) e^{c_1(L)/2}$ [BD81]. $\square$

As an immediate corollary, we have an index formula for twisted spin Dirac operators [LM89].

Corollary 6.8. *Assume that the foliation F is spin and let D be a leafwise Dirac operator associated to the spin structure on TF and complex vector bundle E over M . Then*

$$\mathrm{ch}_a(\mathrm{Ind}_a([\sigma(D)])) = \int_F \hat{A}(TF) \mathrm{ch}(E) \in H_c^*(M/F).$$

Proof. The integrand of Theorem 6.2 is known to be equal to $\hat{A}(TF) \mathrm{ch}(E)$ for such Dirac operators by classical characteristic computations, [ASIII]. $\square$

REFERENCES

- [A75] M. F. Atiyah, *Elliptic operators, discrete groups and von Neumann algebras*, Asterisque **32/33** (1976) 43-72.
- [ASI] M. F. Atiyah and I. Singer, *The index of elliptic operators I*, Ann. of Math. **87** 1968, 484-530.
- [ASIII] M. F. Atiyah and I. Singer, *The index of elliptic operators III*, Ann. of Math. **87** 1968, 564-604.
- [BC84] P. Baum and A. Connes, *K-theory for Lie groups and foliations*, Enseign. Math. **46** (2000), no. 1-2, 3-42.
- [BCH90] P. Baum, A. Connes and N. Higson, *Classifying space for proper actions and K-theory of group C*-algebras*. C*-algebras: 1943-1993 (San Antonio, TX, 1993), 240-291, Contemp. Math., **167**, Amer. Math. Soc., Providence, RI, 1994.
- [BD81] P. Baum and R. Douglas, *K-homology and index theory*, Operator algebras and applications, Part I, Proc. Sympos. Pure Math **38**, Amer. Math. Soc., (1982) 117-173.
- [BH05] M-T. Benaméur and J. L. Heitsch, *Index theory and Non-Commutative Geometry I. Higher Families Index Theory*, Corrigendum, K-Theory **36** (2005) 397-402.
- [BN03] M-T. Benaméur and V. Nistor, *Residues and an index theorem for foliations*, to appear in Math. Scand.
- [Con79] A. Connes, *Sur la théorie de l'intégration non commutative*. Lect. Notes in Math. **725**, 1979.
- [Con81] A. Connes. *A survey of foliations and operator algebras*. Operator algebras and applications, Part I, Proc. Sympos. Pure Math **38**, Amer. Math. Soc., (1982) 521-628.
- [Con86] A. Connes. *Cyclic cohomology and the transverse fundamental class of a foliation*, Geometric methods in operator algebras (Kyoto, 1983), Res. Notes in Mathematics **123**, Pitman, London, (1986) 52-144.
- [Con94] A. Connes, *Noncommutative Geometry*, Academic Press, New York, 1994.

- [CM91] A. Connes and H. Moscovici, *Cyclic cohomology and the Novikov conjecture for hyperbolic groups*, *Topology* **29** (1990) 345–388.
- [CS84] A. Connes, and G. Skandalis. *The longitudinal index theorem for foliations*, *Publ. RIMS Kyoto* **20** (1984) 1139–1183.
- [GL03] A. Gorokhovsky, and J. Lott. *Local index theory over étale groupoids*, *J. reine angew Math* **560** (2003) 151–198.
- [GL05] A. Gorokhovsky, and J. Lott. *Local index theory over foliation groupoids*, *preprint* (2005).
- [H58] A. Haefliger. *Structures feuilletées et cohomologie à valeur dans un faisceau de groupoïdes*, *Comm. Math. Helv.* **32** (1958) 248–329.
- [H70] A. Haefliger. *Feuilletages sur les variétés ouvertes*, *Topology*, **9** (1970) 183–194.
- [H80] A. Haefliger. *Some remarks on foliations with minimal leaves*, *J. Diff. Geo.* **15** (1980) 269–284.
- [He95] J. L. Heitsch, *Bismut superconnections and the Chern character for Dirac operators on foliated manifolds*. *K-Theory* **9** (1995), no. 6, 507–528.
- [HL90] J. L. Heitsch and C. Lazarov. *A Lefschetz theorem for foliated manifolds*, *Topology*, **29** (1990) 127–162.
- [HL99] J. L. Heitsch and C. Lazarov. *A general families index theorem*, *K-Theory*, **18** (1999) 181–202.
- [HL02] J. L. Heitsch and C. Lazarov. *Riemann-Roch-Grothendieck and torsion for foliations* *J. Geo. Anal.* **12** (2002) 437–468.
- [HK99] N. Higson and G. Kasparov, *E-theory and KK-theory for groups which act properly and isometrically on Hilbert space*, *Invent. Math.* **144** (2001), no. 1, 23–74.
- [K88] G. Kasparov, *Equivariant KK-theory and the Novikov conjecture*, *Inv. Math.* **91** (1988) 147–201.
- [LM89] H. B. Lawson and M.-L. Michelson, *Spin Geometry*, *Princeton Math. Series* **38**, Princeton, 1989.
- [NWX96] V. Nistor, A. Weinstein and P. Xu, *Pseudodifferential operators on differential groupoids* *Pacific J. Math.* **189** (1999), no. 1, 117–152.
- [T97] P. Tondeur, *Geometry of foliations*, *Monographs in Mathematics*, 90. Birkhauser Verlag, Basel, 1997.

UNIVERSITÉ DE METZ, ISGMP

E-mail address: `benameur@math.univ-metz.fr`

MATHEMATICS, STATISTICS, AND COMPUTER SCIENCE, UNIVERSITY OF ILLINOIS AT CHICAGO

E-mail address: `heitsch@math.uic.edu`