PIERI HOMOTOPIES FOR PROBLEMS IN ENUMERATIVE GEOMETRY APPLIED TO POLE PLACEMENT IN LINEAR SYSTEMS CONTROL*

BIRKETT HUBER† AND JAN VERSCHELDE‡

Abstract. Huber, Sottile, and Sturmfels proposed Pieri homotopies to enumerate all p-planes in $\mathbb{C}^{m+p}$ that meet n given $(m+1-k_i)$-planes in general position, with $k_1+k_2+\cdots+k_n=mp$ as condition to have a finite number of solution p-planes. Pieri homotopies turn the deformation arguments of classical Schubert calculus into effective numerical methods by expressing the deformations algebraically and applying numerical path-following techniques. We describe the Pieri homotopy algorithm in terms of a poset of simpler problems. This approach is more intuitive and more suitable for computer implementation than the original chain-oriented description and provides also a self-contained proof of correctness. We extend the Pieri homotopies to the quantum Schubert calculus problem of enumerating all polynomial maps of degree q into the Grassmannian of p-planes in $\mathbb{C}^{m+p}$ that meet $mp+q(m+p)$ given m-planes in general position sampled at $mp+q(m+p)$ interpolation points. Our approach mirrors existing counting methods for this problem and yields a numerical implementation for the dynamic pole placement problem in the control of linear systems.

Key words. cheater's homotopy, combinatorial root count, continuation methods, control theory, dynamic pole placement problem, enumerative geometry, Grassmannian, linear system, localization pattern, numerical Schubert calculus, Pieri homotopy, polynomial system, poset, quantum Schubert calculus.

AMS subject classifications. 14N10, 14M15, 65H10, 68Q40, 93B27, 93B55.

1. Introduction.

A general method to solve geometric problems proceeds by moving the input data to special position, solving the special configuration and then deforming the solutions of the special problem into those of the original configuration. As long as the number of solutions remains finite, this number is constant during deformation. This principle of "conservation of number" was developed by Mario Pieri [27], Hermann Schubert [34], and Hieronymus Zeuthen [50].

A classical example in enumerative geometry as explained in [18] consists of finding the two lines in projective 3-space that meet four given lines. More generally, we want to enumerate all p-planes in $\mathbb{C}^{m+p}$ that meet mp given m-planes. In the late nineteenth century, Hermann Schubert [35] established recursive and explicit formulas for the *number* of p planes meeting a set of mp m-planes in general position. This problem was also treated by Mario Pieri in [27]. In the early eighties, Roger Brockett and Christopher Byrnes [5] showed that these p-planes correspond to feedback laws which control a machine whose evolution is governed by a linear system of first-order differential equations. This problem of control theory ([6], [10], [17], [48]) is known as the (static) pole placement problem.

Based on his geometric proof of Pieri's formula [27], Frank Sottile introduced intrinsic deformations in [38] to solve general intersection conditions given by n planes in general position of dimension $m+1-k_i$, for $i=1,2,\ldots,n$. The condition that $k_1+k_2+\cdots+k_n=mp$ guarantees a finite number of p-planes meeting those n given

*Date: 5 October 1999. This work was done while the authors were post-doctors at the Mathematical Sciences Research Institute, 1000 Centennial Drive, Berkeley, CA 94720-5070, USA.

† Wolfram Research, Inc. 100 Trade Center Drive Champaign, IL 61820-7237 E-mail address: birkh@wolfram.com.

‡Supported in part by NSF under Grant DMS - 9804846 at the Department of Mathematics, Michigan State University, East Lansing, MI 48824-1027, USA. E-mail addresses: jan@math.msu.edu and jan.verschelde@na-net.ornl.gov. URL: http://www.mth.msu.edu/~jan.

planes. In [16], Birkett Huber, Frank Sottile, and Bernd Sturmfels reinterpreted these deformations as computational procedures. The resulting algorithm was called the Pieri homotopy algorithm.

Our understanding arose from a first implementation of the Pieri homotopy algorithm and yielded the notion of what we call a "localization pattern". We feel that this is the key concept in our derivation of the algorithm. It is the vehicle through which the combinatorial techniques for enumerating solutions are translated back into synthetic geometry and realized. Since our aim is to understand practical complexity issues, we excuse ourselves from the explicit use of Schubert varieties (as in [15]) in our description. Nevertheless we obtain a self-contained proof of correctness, independent of [16]. The backbone of our approach is the poset of localization patterns used to count the roots combinatorially [43]. We use this poset to set up the homotopies and to control the flow of data between them. Our current description of the algorithm in [16] is both simpler than the original and also more suitable for computer implementation.

A novel feature of this paper is the development of Pieri homotopies for the dynamic pole placement problem. The connection between this problem and the Schubert calculus was established by Joachim Rosenthal in [31] and further developed in [28], [29], [46], and [47]. We arrived at this generalization thanks to a hint of Frank Sottile (explicitly pronounced in [40]) by reverse engineering the root counting procedure described in [30]. Given $mp+q(m+p)$ general m-planes in $\mathbb{C}^{m+p}$ sampled at $mp+q(m+p)$ interpolation points, we can compute all polynomial maps of degree q producing p-planes that meet those given m-planes at the prescribed interpolation points.

Other root counts for these enumerative problems are given in [4]. In [42], quantum Gröbner and SAGBI homotopies are derived. The adjective "quantum" refers to the important connections between this problem of enumerative geometry and physics [7], although as mentioned in [4], one could also speak of "modular" Schubert calculus because of the "modulo $(m+p)$" calculations. Another related problem is the eigenvalue completion problem [13]. We mention [3] as an alternative recent approach to the dynamic pole placement problem.

Homotopies for enumerative geometry are available in a separate module of the publicly available software PHCpack [44]. The Pieri homotopy algorithm consists in a combinatorial root count, the symbolic set up of the homotopies and the tracing of the solution curves. This last stage accounts for most of the computational work and is achieved using numerical continuation methods. See [1], [22] for introductions to and [2], [19] for recent surveys of numerical continuation. Computational experience suggests that the Pieri homotopies perform better than the Gröbner and SAGBI homotopies described in [16]. An explanation for this might be that the Pieri homotopies are more finely tuned to the geometry of the intersection problem than these other, extrinsic methods. In addition we point out that those Gröbner and SAGBI homotopies are restricted to the hypersurface $(k_i = 1)$ intersection conditions.

In our study of the Pieri homotopy algorithm, we first specialized the general description of [16] into the simpler case of hypersurface intersection conditions. Our presentation below reflects this progression from the special to the general. In the next section we show how to find the two lines in projective 3-space that meet four given lines. This example is then generalized to hypersurface and general intersection conditions in the third and fourth section. In the fifth section we describe our quantum Pieri homotopies and their use in solving the dynamic pole placement problem. The

key step is the extension of our notion of localization patterns to degree q-maps of p-planes. The development of numerical algorithms for the pole placement problem is formulated as an open problem in [33]. At the end of this paper we list computational experiences to illustrate the efficiency of our solution method.

2. The geometry of the Pieri deformations. We can illustrate the geometry of the Pieri deformations on the simplest example of determining the two lines meeting four given lines in projective 3-space.

Recall that projective 3-space is the set of 1-dimensional subspaces of $\mathbb{C}^4$. We write $\mathbb{C}^4$ as the span $\langle \mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_4 \rangle$ of the standard basis vectors $\mathbf{e}_1 = (1,0,0,0)$, $\mathbf{e}_2 = (0,1,0,0)$, $\mathbf{e}_3 = (0,0,1,0)$, and $\mathbf{e}_4 = (0,0,0,1)$. Denote the four given lines by L_1, L_2, L_3, and L_4. Without loss of generality we assume that L_3 is spanned by the first two basis vectors, $L_3 = \langle \mathbf{e}_1, \mathbf{e}_2 \rangle$, and L_4 by the last two basis vectors, $L_4 = \langle \mathbf{e}_3, \mathbf{e}_4 \rangle$. In general, the other two lines L_1 and L_2 do not meet each other. This general situation is at the right of Figure 2.1. Here we are visualizing the real positive orthant of projective 3-space as the interior of the tetrahedron spanned by $\mathbf{e}_1$, $\mathbf{e}_2$, $\mathbf{e}_3$, and $\mathbf{e}_4$.



FIG. 2.1. *In $\mathbb{P}^3$ two thick lines meet four given lines L_1, L_2, L_3, and L_4 in a point. At the left we see a special configuration and the general configuration is at the right.*

The special position of L_3 and L_4 allows the following special choice of coordinates for lines X which meet them:

$$X = \langle \mathbf{x}_1, \mathbf{x}_2 \rangle = \begin{bmatrix} x_{11} & 0 \\ x_{21} & 0 \\ 0 & x_{32} \\ 0 & x_{42} \end{bmatrix} \tag{2.1}$$

Here the two columns of X, its first and second generators, are chosen to be its points of intersection with L_3 and L_4 respectively. Lines expressed in these coordinates already satisfy the intersection conditions imposed by L_3 and L_4. Since the columns of X may be scaled independently we have exactly the two degrees of freedom needed to meet the two remaining intersection conditions imposed by L_1 and L_2.

If we take $L_2 = \langle \mathbf{e}_1, \mathbf{e}_3 \rangle$, as at the left of Figure 2.1, we solve the problem as follows. The lines passing through both L_2 and L_3 form the plane $\langle \mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3 \rangle$ which intersects L_4 at $\mathbf{e}_3$. Similarly we see that lines through L_2 and L_4 form the plane $\langle \mathbf{e}_1, \mathbf{e}_3, \mathbf{e}_4 \rangle$ which must intersect L_3 at $\mathbf{e}_1$. Thus we see that any line other than L_2 which meets L_2, L_3, and L_4 must contain either $\mathbf{e}_1$ or $\mathbf{e}_3$. In the first case, consider

the set of lines passing through $\mathbf{e}_1$ and intersecting L_4. These lines form a plane $\langle \mathbf{e}_1, \mathbf{e}_3, \mathbf{e}_4 \rangle$ which must intersect L_1 in a unique point $\mathbf{a}$ and thus the line $\langle \mathbf{e}_1, \mathbf{a} \rangle$ is the unique line through $\mathbf{e}_1$ intersecting L_1, L_2, L_3, and L_4. In the second case, the set of lines through $\mathbf{e}_3$ which intersect L_3 form a plane which intersects L_1 in a unique point $\mathbf{a}'$. So we find that the line $\langle \mathbf{e}_3, \mathbf{a}' \rangle$ is the unique line passing through $\mathbf{e}_3$ and intersecting L_1, L_2, L_3, and L_4.

An algebraic way to look at this argument goes as follows. Any line which meets both L_3 and L_4 must be expressible in the form (2.1). In order for X to meet $L_2 = \langle \mathbf{e}_1, \mathbf{e}_3 \rangle$ we must be able to find non-trivial linear combinations such that $\lambda_1 \mathbf{x}_1 + \lambda_2 \mathbf{x}_2 = \gamma_1 \mathbf{e}_1 + \gamma_2 \mathbf{e}_3$. That is the system

$$\begin{bmatrix} 1 & x_{11} & 0 & 0 \\ 0 & x_{21} & 0 & 0 \\ 0 & 0 & 1 & x_{32} \\ 0 & 0 & 0 & x_{42} \end{bmatrix} \begin{bmatrix} -\gamma_1 \\ \lambda_1 \\ -\gamma_2 \\ \lambda_2 \end{bmatrix} = 0 \tag{2.2}$$

must have a non-trivial solution. This can clearly happen only when $x_{21} = 0$ or $x_{42} = 0$. The first case then describes the plane of lines containing $\mathbf{e}_1$ and meeting L_4 and gives rise to the solution $\langle \mathbf{e}_1, \mathbf{a} \rangle$ as above, while $x_{42} = 0$ yields the solution $\langle \mathbf{e}_3, \mathbf{a}' \rangle$.

Schubert's principle [34], [50], of "conservation of number" suggests that the number of solutions to the general problem will be the same as the number of the solutions in the special case we just looked at, i.e.: 2. In order to adjust this argument to allow us to *find* the solutions as well as count them we must consider the process of deforming L_2 to special position.

The special position we use for L_2 is given by $S = \langle \mathbf{e}_1, \mathbf{e}_3 \rangle$, so our deformed problem is: find all X such that $\det(X|L_1) = 0$, $\det(X|S) = 0$, $\det(X|L_3) = 0$, and $\det(X|L_4) = 0$. Note that the last two equations are already satisfied when X is chosen as in (2.1). Specializing the second condition can be achieved by moving the line S into L_2 taking a convex combination $(1-t)S + tL_2$ of the generating matrices S and L_2 with t varying from 0 to 1. Fixing the first condition and enforcing non-trivial intersection with the moving line yields the homotopy:

$$H(X,t) = \begin{cases} \det(X|(1-t)S + tL_2) = 0 & \text{where} \quad X = \langle \mathbf{x}_1, \mathbf{x}_2 \rangle, \\ \det(X|L_1) = 0 & \qquad\quad \mathbf{x}_1 \in L_3, \mathbf{x}_2 \in L_4. \end{cases} \tag{2.3}$$

Numerical continuation methods trace the solution paths starting at the two solutions $X = \langle \mathbf{e}_1, \mathbf{a} \rangle$ and $X = \langle \mathbf{e}_3, \mathbf{b} \rangle$ calculated above for $t = 0$ to the solutions of the original problem at $t = 1$.

The straightforward application of this geometric principle of conservation of number may fail if our special position is too special in the sense that it is a singular point in the parameter space of configurations. This is a very subtle condition that is a property of the special position as a member of the set of problem instances and can not be determined by considering the geometry of the special position itself. Therefore we need to examine the algebraic formulation of the deformation near $t = 0$. As local coordinates we choose $x_{11} = 1$ and $x_{32} = 1$. Then $\det(X|S) = 0 \Leftrightarrow x_{21} x_{42} = 0$. So, at $t = 0$, as either $x_{21} = 0$ or $x_{42} = 0$, the structure of the matrix of partial derivatives of the system $H(X, t = 0)$ is

$$\begin{bmatrix} x_{42} & x_{21} \\ C_{13} x_{42} - C_{14} & C_{13} x_{21} - C_{23} \end{bmatrix} \tag{2.4}$$

with C_{ij} the 2-by-2 minor by selecting rows i and j from the matrix representation of L_4. For L_1 in general position, the two start solutions are regular and admit well-defined deformations. For general L_2, this regularity is maintained up to $t = 1$.

3. Pieri-type deformations for hypersurface conditions. In this section we extend the approach of the last section to produce an algorithm for computing all p-planes in $\mathbb{C}^{m+p}$ that intersect mp given m-planes L_i in general position. In order to have a uniform description of the algorithm, we will skip the step of specializing the last two m-planes, and the last step of using linear algebra to intersect with the first m-plane. Instead we will describe the algorithm solely in terms of the continuation step which involved deforming L_2 to S. Later we shall see that adding these linear algebra steps back in amounts to truncating the algorithm as we describe it and does yield a significant computational savings.

We represent p-planes in $\mathbb{C}^{m+p}$ by $(m+p) \times p$-matrices whose columns form a set of generators. Solutions to our problem are represented by variable $(m+p) \times p$-matrices. We move the input data into special positions such that the solution planes have matrix representations with zero coordinates at specific positions. To specify those positions, we need the following definition.

DEFINITION 3.1. A *localization pattern* is an element of $\{0, \star\}^{(m+p)\times p}$ such that all stars in a column are contiguous and the row indices in which the bottommost and topmost stars occur strictly increase as functions of the column number. These row indices are called the *top* and *bottom pivots* respectively. A p-plane *fits a localization pattern* if it can be represented by a matrix of generators with zero entries everywhere the localization pattern prescribes them.

Consider for example the case where $p = 2, m = 4$ and we are looking for all 2-planes which meet eight 4-planes. We can take the last two planes as $L_7 = \langle \mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_4 \rangle$ and $L_8 = \langle \mathbf{e}_3, \mathbf{e}_4, \mathbf{e}_5, \mathbf{e}_6 \rangle$. For any 2-plane that meets both these planes non-trivially we may choose its generators such that one generator lies in L_7 and the other in L_8. Such a 2-plane is represented by a variable matrix

$$\begin{bmatrix} x_{11} & 0 \\ x_{21} & 0 \\ x_{31} & x_{32} \\ x_{41} & x_{42} \\ 0 & x_{52} \\ 0 & x_{62} \end{bmatrix} \quad \text{which fits the pattern} \quad \begin{bmatrix} \star & 0 \\ \star & 0 \\ \star & \star \\ \star & \star \\ 0 & \star \\ 0 & \star \end{bmatrix}. \tag{3.1}$$

Planes which satisfy this localization pattern already meet L_7 and L_8. Equivalently, any specialization of the variables x_{ij} in the pattern will result in a p-plane which has non-trivial intersection with L_7 and L_8. Thus our problem is to find values of these variables such that the plane represented by (3.1) meets the remaining 4-planes. We achieve this by specializing the input planes as follows.

DEFINITION 3.2. Given a localization pattern X, *the special m-plane for the top (bottom) pivots of X* is the m-plane S_X spanned by the standard basis vectors not indexed by top (bottom) pivots.

It turns out that p-planes which meet fit a localization pattern X and meet its special m-plane S_X must fit a set of localization patterns easily derived from X in the following way.

DEFINITION 3.3. A *top (bottom) child* of a localization pattern X is a localization pattern obtained by turning a topmost (bottommost) star of X to a zero.

For the bottom localization pattern in Figure 3.1 having bottom pivots [3 5], the special 4-plane is $\langle \mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_4, \mathbf{e}_6 \rangle$. Lemma 3.4 states that all 2-planes meeting this special 4-plane and fitting the bottom pattern must fit one of its two children.



FIG. 3.1. *A localization pattern with bottom pivots* [3 5] *and its two children* [2 5] *and* [3 4]. *At the right we see the bottom pivots used as a shorthand notation.*

LEMMA 3.4. (Hypersurface Pieri rule) *Let S_X be the special m-plane for top (bottom) pivots of the localization pattern X. Every p-plane that fits X and meets S_X also fits one of the (at most p) children of X. Conversely every p-plane that fits a child of X fits X and meets S_X.*

Proof. A p-plane that fits X meets S_X if, and only if, for all specific values for the variables corresponding to the stars in the pattern X, we can find a linear combination of the columns of the p-plane which lies in S_X. This means that the intersection of the p-plane with S_X can be represented as a linear combination of the columns of S_X. Finding this linear combination is equivalent to finding a non-trivial element of the kernel of $[X|S_X]$. Since the m columns of S_X have pivots distinct from those of the p columns of X, the matrix $[X|S_X]$ is an $(m+p) \times (m+p)$-matrix with $m+p$ distinct pivot rows. So we can rearrange $[X|S_X]$ into a triangular matrix, which is singular precisely when one of the diagonal elements coming from X is specialized to zero. □

The repeated application of Lemma 3.4 builds up a poset diagram which allows us to count the solution planes as shown in Figure 3.2. The poset starts at the trivial localization pattern X_0 whose i-th column has its top pivot in row i and its bottom pivot in row $m+i$. While there are p-planes that do not fit X_0, the set of p-planes that do fit X_0 includes those for which all Plücker coordinates are non-zero. This is a dense open subset of the Grassmannian and with the general position assumption on our input planes it will suffice to count the number of solution planes which fit X_0. This start pattern, X_0 has exactly $mp+p$ stars and our original list of conditions contains exactly mp given m-planes. Initially we apply Theorem 3.5 starting at $X = X_0$ and $n = mp$:

THEOREM 3.5. *Consider a localization pattern X with $p+n$ stars. For $n = 0$, X counts for one solution. For $n > 0$, the number of p-planes fitting X and meeting n general m-planes equals the sum of the number of solution planes fitting the children of X and meeting $n-1$ general m-planes.*

Theorem 3.5 is proven by induction and follows from Lemma 3.7 establishing the correctness of the Pieri homotopy. Applying the Pieri homotopy algorithm following the root counting procedure through the poset yields an effective algorithm for actually finding the solutions.

DEFINITION 3.6. Let X be a localization pattern and S_X the special m-plane for top (bottom) pivots of X. Suppose that $n-1$ general intersection conditions are satisfied for children Y of X i.e.: $\det(Y|L_i) = 0$, for $i = 1, 2, \ldots, n-1$. To satisfy the nth intersection condition, the *Pieri homotopy* $H(X,t) = \mathbf{0}$ is:

$$(3.2)\quad H(X,t) = \begin{cases} \det(X|(1-t)S_X + tL_n) = 0 & \\ \det(X|L_i) = 0 \quad i = 1,2,\ldots,n-1 & \end{cases} \qquad t \in [0,1].$$



FIG. 3.2. *Combinatorial root count for $p = 2$, $m = 4$. The brackets at the left are the bottom pivots. The top pivots remain* [1 2]. *The trivial localization pattern X_0 corresponds to* [5 6]. *The roots are counted at the right, starting at the top and adding up the numbers at the leaves while moving down to the root of the poset yielding* 14 *solutions.*

To illustrate the Pieri homotopy, we consider the situation as in going down to [3 5], with the three 2-planes that fit the left child [2 5] and two 2-planes that fit the right child [3 4]. Locally the situation is pictured in Figure 3.1, with the global root counting procedure in Figure 3.2. At $t = 0$, $n = 4$, and we have already folded in the first four intersection conditions and are moving the special 4-plane $\langle \mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_4, \mathbf{e}_6 \rangle$ to the fifth input plane. In general we have the situation as in the following Lemma.

LEMMA 3.7. *Consider a localization pattern X with $p + n$ stars and n complex m-planes L_i in general position. Suppose we are given all p-planes that meet L_i, $i = 1, 2, \ldots, n-1$ and fit one of the children of X. Then the Pieri homotopy defines regular paths of p-planes, that start at the given p-planes and end at those p-planes which meet all n general m-planes L_i while fitting X.*

Proof. Our working space is a product of projective spaces, with as many spaces as there are columns in X. To describe this multi-projective space more precisely, we count the stars in X. For X defined by p-tuples of top and bottom pivots, respectively $\alpha, \beta \in \mathbb{N}^p$, we embed the p-planes into $\mathbb{P}^{d_1} \times \mathbb{P}^{d_2} \times \cdots \times \mathbb{P}^{d_p}$ with $d_i = \beta_i - \alpha_i$, $i = 1, 2, \ldots, p$. To fix an affine coordinate chart we set one coordinate to 1 in every column by scaling the corresponding generator of the p-plane.

Essentially we are applying a multi-homogeneous homotopy [23, 24] in a general situation. By the assumption of this Lemma, all p-planes at $t = 0$ are regular solutions, as the m-planes L_i are in general position. This general position is maintained for all $t \in [0, 1]$, ensuring the smoothness of the solution paths.

We still have to show that we will find all p-planes. Suppose that at $t = 1$ there are more solutions than the number of paths we started with. Going backwards with those additional solutions from $t = 1$ to $t = 0$ we either move to a singular solution, or to a solution at infinity. But at $t = 0$, all solutions are regular. By Lemma 3.4, all solutions have been found and since the localization patterns admit any affine chart, there are no solutions at infinity for $t = 0$. □

We remark that the root count in Figure 3.2 by changing only bottom pivots is just one possibility. One could alternatingly change bottom and top pivots in building up the levels of the poset. As was done in section 2, linear algebra can be applied to intersect with the first input plane. Another, more significant optimization includes the special position of the last two input planes. Choosing a basis so that the last two m-planes are already in special position $L_{mp-1} = \langle \mathbf{e}_1, \ldots, \mathbf{e}_m \rangle$ and $L_{mp} = \langle \mathbf{e}_{p+1}, \ldots, \mathbf{e}_{m+p} \rangle$ amounts to enforcing a different starting pattern for the poset. This

has two advantages. The first is that it really forces any pattern meeting all original planes to meet the starting localization pattern regardless of the genericity of the remaining input planes. The second is that it saves the work of doing the two final and most costly (because of highest dimension) continuation steps.

In contrast to the approach of [16], our description treats the hypersurface Pieri homotopy algorithm separately from the case of general intersection conditions. This allows us to simplify the definition of special m-planes. However, when all input m-planes are spanned by real matrices, we need the more general choice of special m-planes as defined in [16] to enforce complex arithmetic and avoid quadratic turning points along the solution paths. Also for the general intersection conditions we need the definition of [16] for special m-planes. So in Definition 3.2 we replace each of the standard basis vectors $\mathbf{e}_i$ by vectors $\mathbf{b}_i$ that have random complex entries in rows i and higher (lower) for bottom (top) pivots i. This adaption of Definition 3.2 we assume for the rest of the paper.

4. General intersection conditions. In the general case we are looking for p-planes in $\mathbf{C}^{m+p}$ which non-trivially meet a list of generic subspaces $L_1, L_2, \ldots, L_n$, where each L_i has dimension $m+1-k_i$. It turns out each of these conditions removes k_i degrees of freedom, and thus that we will expect a finite number of solutions when $k_1 + k_2 + \cdots + k_n = mp$.

In specializing the ith $(m+1-k_i)$-plane L_i we move it to the intersection of k_i special m-planes. This deformation of the solution p-planes while moving L_i from general to special position proceeds in k_i steps, each time specializing some of the coordinates of our current representation for the solution planes to zero. In modeling this deformation we need to be aware of the left and right sides of our localization patterns when working respectively with bottom and top pivots.

DEFINITION 4.1. For the localization patterns X^1 and X^2 we write $X^1 \supset_{c_1} X^2$ to indicate that X^2 is the child of X^1 obtained by turning the pivot in column c_1 of X^1 to zero. We say that the *chain of length j*

$$C = X^1 \supset_{c_1} X^2 \supset_{c_2} \cdots \supset_{c_{j-1}} X^j \tag{4.1}$$

is *bottom-left* if $c_i \leq c_{i-1}$, $i = 2, \ldots, j$ for bottom pivots c_i, or *top-right* if $c_i \geq c_{i-1}$, $i = 2, \ldots, j$ for top pivots c_i. If C is bottom-left (resp. top-right), we will call any child X^{j+1} of X^j a *bottom-left child* (resp. *top-right child*) of C if the chain $X^1 \supset_{c_1} \cdots \supset_{c_{j-1}} X^j \supset_{c_j} X^{j+1}$ is also bottom-left (resp. top-right).

In other words, in a bottom-left (top-right) chain, we never take away a bottom (top) pivot in a column to the right (left) of one we have already taken. Note that when $C = X^1$ is a chain of length one, we have not taken any pivots away yet and all children of X^1 are bottom-left (top-right) children of C.

Notice that the pivots of $X^1 \supset_{c_1} X^2$ are the same except in column c_1 and that the new pivot of X^2 can not have been a pivot of X^1 (otherwise X^2 would not be a localization pattern). Thus $S_{X^1} \cap S_{X^2}$ has dimension $m-1$. Similarly we can see that the intersection $S_{X^1} \cap S_{X^2}$ with S_{X^3} drops the dimension by one whenever X^3 is a child of X^2 obtained by deleting a pivot in a column with index less than or equal to c_1. This justifies the dimension of the following definition.

DEFINITION 4.2. For a bottom-left (top-right) chain $C = X^1 \supset_{c_1} X^2 \supset_{c_2} \cdots \supset_{c_{j-1}} X^j$, and an index $1 \leq k \leq j$ we define a *special $(m+1-k)$-plane* S_C^k to be $S_C^k = \cap_{i=1}^k S_{X^i}$, where the S_{X^i} are the special m-planes associated to the X^i in C.

From now on we assume that we work with bottom pivots.

LEMMA 4.3. (General Pieri rule) *Let C be a bottom-left chain of length j ending at X^j.*

1. *All p-planes fitting X^j and meeting S_C^j must fit at least one of X^j's children.*
2. *A child X^{j+1} meets S_C^j iff X^{j+1} is a bottom-left child of C, i.e. iff $X^j \supset_{c_j} X^{j+1}$ where $c_{j+1} \leq c_j$.*

Proof. Any p-plane which fits X^j and meets S_C^j must also meet S_{X^j} and thus must be a child of X^j by Lemma 3.4. This proves the first claim.

Recall from the proof of Lemma 3.4 that X^{j+1} meets S_C^j if, and only if, given any specification of the variables of X^j for which the pivot elements are non-zero it is possible to find a linear combination of the columns which lies in S_C^j. Clearly any such linear combination must involve the decremented column c_{j+1} and no columns after it, since all pivots of vectors in S_C^j must be non-pivots of X^j.

If $c_{j+1} \leq c_j$, then the first $c_j - 1$ columns of X^{j+1} have remained unchanged all the way through the chain, and the last has only had successive pivots removed. Thus S_C^j must contain a basis vector with a pivot row in each non-pivot of the first c_{j+1} columns of X^j. So if we take these basis vectors along with the first c_{j+1} columns of X^j we get a set of r_j column vectors with r_j distinct pivots all in rows r_j. Since X^{j+1} is obtained from X^j by deleting the pivot of column c_{j+1} we find that the (c_{j+1})th column must be in the span of the remaining columns, and this linear relation can be rewritten to express a vector of S_C^j as a linear combination of columns of X^{j+1}.

On the other hand, if $c_{j+1} > c_j$, then at least one non-pivot of these first c_{j+1} columns of X^j must have been a pivot previously and S_C^j must have no vector with a pivot in this row. In other words we are looking for a solution to a subsystem of the previous system where we have removed some number of columns (aside from that corresponding to column c_{j+1} of X^{j+1}). Looking at just the pivot rows appearing we get either a square triangular system with non-zeros on the diagonal (if the pivot of column c_{j+1} did not appear) or a non-square system with non-zero pivots in each column. In either case no non-trivial solution is possible. □

THEOREM 4.4. *Let X^1 be a localization pattern with $p + \sum_{i=1}^{n-1} k_i$ stars where the k_i are the codimension conditions specified in the problem (i.e.: $dim(L_i) = m+1-k_i$). Further suppose that $C = X^1 \supset_{c_1} X^2 \supset_{c_2} \cdots \supset_{c_{j-1}} X^j$, $(1 \leq j \leq k_n)$ is a bottom-left chain starting at X_1. For $1 \leq k \leq j$ let U_C^k be a generic $(m+1-k_n)$-dimensional subspace of $S_C^k = \cap_{i=1}^k S_{X^i}$. Then the number of p-planes meeting U_C^{j-1} as well as $L_1, \ldots, L_{n-1}$ and fitting X^j is equal to the sum of the numbers of p-planes meeting $U_C^j, L_1, \ldots, L_{n-1}$ and meeting Y as Y runs over all bottom-left children of C.*



FIG. 4.1. *Bottom pivots for $p = 2$, $m = 4$, $k_i = 2$, $i = 1, 2, 3, 4$. Only those arcs and nodes are drawn that connect localization patterns reachable by bottom-left chains. The corresponding root count is at the right.*

As with the hypersurface case, Theorem 4.4 is proven by induction and follows from the correctness of a homotopy. Before describing this homotopy we will show that the theorem provides a root count procedure generalizing that of the last section. Combining the homotopy with this counting procedure will yield an effective technique for producing all solutions. It is a little easier to understand the implications of the theorem for root counting if we restate it without reference to the intermediate steps where our input planes are partially specialized:

COROLLARY 4.5. *Suppose X is a localization pattern with $p+\sum_{i=1}^{n} k_i$ stars where the k_i are codimension conditions specified in the problem (i.e.: $dim(L_i) = m+1-k_i$). Then the number of p-planes fitting X and meeting $L_1, \ldots, L_n$ is equal to the total number of p-planes meeting $L_1, \ldots, L_{n-1}$ and fitting at least one descendant Y of X accessible as a bottom left child of a bottom-left chain of length k_n starting at X.*

Proof. To see that the corollary follows from Theorem 4.4, consider applying the theorem k_n times starting from the hypothesis of the corollary. Note that the first application of the theorem leaves us with the problem of counting the number of p-planes meeting $L_1, \ldots, L_{n-1}$, and also meeting an $m+1-k_n$ plane lying in S_X and fitting a child of X. Each of these numbers can now be counted by applying the theorem to the new case with $n = n-1$, $j = 2$, and $C = C^1 \supset C^2$. This process continues until all our chains have length k_n and the subspace $U_C^{j+1} = S_C^{k_n}$ in the conclusion of Theorem 4.4 is a special $m+1-k_n$ space and that Lemma 4.3 implies that any p-plane fitting a bottom-left child of C will meet S_C^j. □

Figure 4.1 illustrates the combinatorial counting method arising from Theorem 4.4 just as Figure 3.2 did for the hypersurface case. Starting at the root of the poset we apply the theorem to the case $n = 4$, $j = 1$, $C = [5\ 6]$ to see that the number of 2-planes which fit the pattern [5 6] and meet 4 general 3-planes L_1, L_2, L_3, L_4 is equal to the number of p-planes which fit [4 6], meet the 3 general 3-planes L_1, L_2, L_3 and meet a general 3-plane lying inside the special 4-plane $S_{[5\ 6]}$. A second application of the theorem in the case $n = 3$, $j = 2$, $C = [5\ 6] \supset_1 [4\ 6]$ now tells us that this number is equal to the number of p-planes fitting the pattern [3 6] and meeting the 3 general 3-planes L_1, L_2, L_3. Note that by Lemma 4.3 we do not get any solutions fitting the pattern [4 5] (so we omitted [4 5] in the Figure). Continuing, with $n = 3$, $j = 1$, $C = [3\ 6]$, we find that the number of 2-planes fitting the pattern [3 6] and meeting L_1, L_2, L_3 is equal to the sum of the numbers fitting the two children of [3 6] and meeting L_1, L_2, and a general 3-plane lying inside of $S_{[3\ 6]}$. Applying the theorem with $n = 3$, $j = 2$, to the chains $[3\ 6] \supset_1 [2\ 6]$ and $[3\ 6] \supset_2 [3\ 5]$ we find that the total number of 2-planes fitting the pattern defined by [3 6] and meeting L_1, L_2, and L_3 is equal to the sum of the numbers of 2-planes meeting L_1 and L_2 and fitting any of the localization patterns defined by [1 6], [2 5], and [3 4].

It is important to note that during the intermediate steps when the chains C have length $j > 1$, the number of solutions at a node depends on both the localization pattern and the chain. Thus we can only share nodes corresponding to the $j = 1$ case, i.e.: those nodes with no partially specified input planes. In Figure 4.1 these cases (with $j = 1$) are indicated by dotted horizontal lines. Also note that some chains may not make it k_n steps and may simply stop and fail to yield any solutions. Both these points are illustrated by considering the localization pattern [1 5] in Figure 4.1. Applying Theorem 4.4 with $n = 2$, $j = 2$, on the chains $[1\ 6] \supset_2 [1\ 5]$ yields an equivalence with the numbers of solutions fitting [1 4] and meeting L_1 while applying the theorem to with $n = 2$, $j = 2$, and the chain $[2\ 5] \supset_1 [1\ 5]$ yields no solutions at all since this chain has no bottom-left children. We are now ready to define the

homotopy which provides our effective proof of Theorem 4.4.

DEFINITION 4.6. Let X, n, j, and C be as in Theorem 4.4, and suppose that the subspaces U_C^j and U_C^{j-1} are represented by $(m+p)\times(m+1-k_n)$-dimensional matrices made up of random linear combinations of the S_C^j and S_C^{j-1} (except that $U_C^0 = L_n$). We define the *Pieri homotopy* $H_C^j(X,t) = \mathbf{0}$ to be

$$H_C^j(X,t) = \begin{cases} \text{all maximal minors of } [X|(1-t)U_C^j + tU_C^{j-1}] \\ \text{all maximal minors of } [X|L_i] = 0, \\ \qquad\qquad i = 1,2,\ldots,n-1 \end{cases} \qquad t \in [0,1]. \tag{4.2}$$

Note that while this homotopy may well have more equations than unknowns it will turn out to define a nonsingular path of solutions through each starting point. Here we will prove that these equations define a homotopy which can be used to find all solutions. We will defer a discussion of techniques for actually following these paths numerically to section six.

LEMMA 4.7. *Supposing that the solutions of $H_C^j(X,0) = \mathbf{0}$ are all nonsingular as X varies over all bottom-left extensions of C then the Pieri homotopy $H_C^j(X,t) = \mathbf{0}$ defines regular paths through these solutions which lead to all solutions of $H_C^j(X,1)$.*

Proof. As in the proof of Theorem 3.5, our working space is the product $\mathbb{P}^{d_1}\times\mathbb{P}^{d_2}\times\cdots\times\mathbb{P}^{d_p}$ of projective spaces where d_j is the number of stars in each column of X, and we are essentially applying a multi-homogeneous homotopy [23, 24]. By assumption all p-plane solutions at $t = 0$ are regular, and since U^j defines an $(m+1-k_n)$-dimensional subspace of S_C^{j-1} we see that U^{j-1} which defines a generic $(m+1-k_n)$-dimensional subspace of S_C^{j-1} must also define an intersection problem with a solution set consisting of a finite number of regular p-planes. This general position is maintained for all $t \in [0,1]$, ensuring the smoothness of the solution paths.

It remains to show that we will find all p-planes. This argument is exactly analogous to the proof of Theorem 3.5 except that we rely on Lemma 4.3 to show that all solutions found must fit some bottom-left child of C when we go backwards. ∎

Theorem 4.4 follows from the lemma by induction. We start at the tops of the counting poset where the localization patterns all have exactly p stars and no subspace conditions to meet and hence define unique solutions. The hypotheses of the homotopy lemma are then met and this forces the conditions to be met all the way down.

5. Control of linear systems and quantum Schubert calculus. This section is organized in three parts. We first rephrase the dynamic pole placement problem into a problem of enumerative geometry [31]. Thereafter we derive the critical dimension [47] for this problem to have a finite number of solutions. Lastly we state the root-counting theorem [29] and prove it inductively from the correctness of the Pieri homotopies.

Suppose we want to control a plant with n internal states $\mathbf{x} \in \mathbb{R}^n$ that takes m-inputs $\mathbf{u} \in \mathbb{R}^m$ and produces p-outputs $\mathbf{y} \in \mathbb{R}^p$ with a dynamic compensator that has q internal states $\mathbf{z} \in \mathbb{R}^q$. Schematically we picture this situation in Figure 5.1.

The evolution in time of the plant is described by a system of first-order differential equations:

$$\begin{cases} \dot{\mathbf{x}} = A\mathbf{x} + B\mathbf{u} & \text{with } \mathbf{x}\in\mathbb{R}^n, \mathbf{u}\in\mathbb{R}^m, \mathbf{y}\in\mathbb{R}^p, \\ \mathbf{y} = C\mathbf{x} & \text{and } A\in\mathbb{R}^{n\times n}, B\in\mathbb{R}^{n\times m}, C\in\mathbb{R}^{p\times n}. \end{cases} \tag{5.1}$$

The dynamic compensator obeys a qth order differential equation, described by the



FIG. 5.1. *Control of an m-input and p-output plant by a qth order dynamic compensator.*

system:

$$\text{(5.2)}\quad \begin{cases} \dot{\mathbf{z}} = F\mathbf{z} + G\mathbf{y} \\ \mathbf{u} = H\mathbf{z} + K\mathbf{y} \end{cases} \quad \text{with} \quad \mathbf{z} \in \mathbb{R}^q \quad \text{and} \quad \begin{matrix} F \in \mathbb{R}^{q\times q}, G \in \mathbb{R}^{q\times p}, \\ H \in \mathbb{R}^{m\times q}, K \in \mathbb{R}^{m\times p}. \end{matrix}$$

After elimination of $\mathbf{u}$ and $\mathbf{y}$ concatenation of (5.1) and (5.2) yields the following closed-loop system:

$$\text{(5.3)}\quad \begin{bmatrix} \dot{\mathbf{x}} \\ \dot{\mathbf{z}} \end{bmatrix} = \begin{bmatrix} A+BKC & BH \\ GC & F \end{bmatrix} \begin{bmatrix} \mathbf{x} \\ \mathbf{z} \end{bmatrix}$$

The behavior of this closed-loop system is determined by the $n+q$ eigenvalues of the matrix in (5.3). For a plant given by the matrix triplet (A,B,C) and $n+q$ eigenvalues, the dynamic pole placement problem asks for the matrix quadruples (F,G,H,K) which determine the dynamic compensators that yield closed-loop systems with a specific set of eigenvalues.

We can formulate the dynamic pole placement problem as a geometric problem. In rewriting the characteristic equation of (5.3) the subscripts in I denote the dimension of the identity matrix. There is only some hindsight involved in deriving (5.6). The rest of the equivalences are deduced by elementary row and column operations.

$$\text{(5.4)}\quad \det\left(s\begin{bmatrix} I_n & 0 \\ 0 & I_q \end{bmatrix} - \begin{bmatrix} A+BKC & BH \\ GC & F \end{bmatrix} \right) = 0$$

$$\text{(5.5)}\quad \Leftrightarrow \det\begin{bmatrix} sI_n - A - BKC & -BH \\ -GC & sI_q - F \end{bmatrix} = 0$$

$$\text{(5.6)}\quad \Leftrightarrow \det\begin{bmatrix} sI_n - A - BKC & -BH & BK & -B \\ -GC & sI_q - F & G & 0 \\ 0 & 0 & I_p & 0 \\ 0 & 0 & 0 & I_m \end{bmatrix} = 0$$

$$\text{(5.7)}\quad \Leftrightarrow \det\begin{bmatrix} sI_n - A & 0 & 0 & -B \\ 0 & sI_q - F & G & 0 \\ C & 0 & I_p & 0 \\ 0 & -H & K & I_m \end{bmatrix} = 0$$

$$\text{(5.8)}\quad \Leftrightarrow \det\begin{bmatrix} I_n & 0 & 0 & -(sI_n - A)^{-1}B \\ 0 & I_q & (sI_q - F)^{-1}G & 0 \\ C & 0 & I_p & 0 \\ 0 & -H & K & I_m \end{bmatrix} = 0$$

$$\text{(5.9)}\quad \Leftrightarrow \det\begin{bmatrix} I_n & 0 & 0 & -(sI_n - A)^{-1}B \\ 0 & I_q & (sI_q - F)^{-1}G & 0 \\ 0 & 0 & I_p & C(sI_n - A)^{-1}B \\ 0 & 0 & H(sI_q - F)^{-1}G + K & I_m \end{bmatrix} = 0$$

The last $m+p$ rows and columns of (5.9) represent the intersection of m-planes defined by the given triplet (A,B,C) with p-planes determined by the unknown quadruple (F,G,H,K). As these p-planes depend on the variable s introduced by the the term $(sI_q - F)^{-1}$, they have maximal minors of degree q and we call them degree q-maps. Thus the dynamic pole placement problem is equivalent to the computation of all degree q-maps into the Grassmannian of p-planes that meet n given m-planes at prescribed s-values. It is easy to see that for each specification of an eigenvalue λ_i the condition that the characteristic polynomial (5.9) vanish at $s = \lambda_i$ enforces one polynomial condition on the set of degree q-maps.

Note that when $q = 0$ we are solving the static pole placement problem ($\mathbf{u} = K\mathbf{y}$ and $F, G, H = 0$) and are looking for maps of degree 0 (i.e. constant maps) which meet a specific set of given m-planes. In this case the characteristic equation (5.9) has degree n and we can find solution planes whenever n is less than the dimension mp of the space of p-planes in $(m+p)$-dimensional space. At the critical dimension $n = mp$, we expect the number of such solutions to be finite. In other words when $q = 0$ and $n = mp$ this analysis transforms the static pole placement problem into the problem of finding all p-planes meeting a set of mp given m-planes dealt with in Section 3. For general q we will be able to use similar methods to solve it, but will need the following lemma which specifies the critical dimension and allows us to produce localization patterns for our maps.

LEMMA 5.1. *Let $q = dp + r$ with $d, r \in \mathbb{N}$ and $r < p$. The set of $(m+p) \times p$ degree q-maps is of dimension $mp+q(m+p)$ and has a dense open subset which can be represented by matrices with entries of maximum degree d in all but the last r columns and degree $d+1$ in the remaining r columns.*

Proof. Given a polynomial matrix we will define the degree of a column to be the maximum degree of any of its entries. Also note that two $(m+p) \times p$ polynomial matrices define the same map if there is an invertible $p \times p$-matrix $U(s)$ which takes one into the other by right multiplication. Lemma 3.1 from [47] tells us that for any $(m+p) \times p$ degree q-map $X'(s)$, there is a unique choice of column degrees $d_1 \leq \ldots \leq d_p$ (with $q = d_1 + \cdots + d_p$) such that the $X'(s)$ is equivalent to a matrix $X(s)$ with these column degrees. In other words the sets $U_{d_1,\ldots,d_p}$ of degree q-maps which can be represented by $(m+p) \times p$-matrices with column degrees $d_1, \ldots, d_p$ disjointly cover the set of degree q-maps.

In order to find the dimension of $U_{d_1,\ldots,d_p}$ we will want to find a canonical form for a general member. By a general member of $U_{d_1,\ldots,d_p}$ we mean, a matrix $X(s)$ whose entries are polynomials of appropriate degrees with no algebraic relations among the coefficients. If we write $a^k_{i,j}$ for the coefficient of s^k in the (i,j)th entry of $X(s)$, we can express our normal form by requiring $a^0_{i,i} = 1$ for $i = 1, \ldots, p$ and if for all $i < j$ we have $a^k_{i,j} = 0$ if $k \leq d_j - d_i$ and $a^0_{j,i} = 0$ if $d_j = d_i$. This normal form is illustrated in Figure 5.2 and can be achieved by the following algorithm:

for $i = 1, 2, \ldots, p$ do
  $X_i := \frac{1}{a^0_{i,i}} X_i$;
  for $j = i+1, i+2, \ldots, p$ do
    if $d_i = d_j$, then $X_i := X_i - \frac{a^0_{j,i}}{a^0_{j,j}} X_j$; *zero out* $a^0_{j,i}$
    for $k = 0, 1, \ldots, d_j - d_i$ do $X_j := X_j - \frac{a^k_{i,j}}{a^0_{i,i}} s^k X_i$. *zero out* $a^k_{i,j}$

The genericity of $X(s)$ ensures that the replacements in this algorithm each introduce no zeros besides the desired one in the coefficient $a^k_{i,j}$ so that the algorithm is

$$\begin{bmatrix} 1 & 0 & 0 & 0+0s & 0+0s+0s^2 \\ 0 & 1 & 0 & 0+0s & 0+0s+0s^2 \\ 0 & 0 & 1 & 0+0s & 0+0s+0s^2 \\ \star & \star & \star & 1+\star s & 0+0s+\star s^2 \\ \star & \star & \star & \star+\star s & 1+\star s+\star s^2 \\ \star & \star & \star & \star+\star s & \star+\star s+\star s^2 \\ \star & \star & \star & \star+\star s & \star+\star s+\star s^2 \end{bmatrix} \qquad \begin{bmatrix} 1 & 0 & 0+0s & 0+0s & 0+0s \\ 0 & 1 & 0+0s & 0+0s & 0+0s \\ \star & \star & 1+\star s & 0+\star s & 0+\star s \\ \star & \star & 0+\star s & 1+\star s & 0+\star s \\ \star & \star & 0+\star s & 0+\star s & 1+\star s \\ \star & \star & \star+\star s & \star+\star s & \star+\star s \\ \star & \star & \star+\star s & \star+\star s & \star+\star s \end{bmatrix}$$

$U_{0,0,0,1,2}$ $\qquad$ $U_{0,0,1,1,1}$

FIG. 5.2. *Standard forms for $m = 2$, $p = 5$ and $q = 3$ with 28 and 31 stars respectively*

well defined and will terminate having introduced exactly the zeros required for the canonical form. It is easy to check that each of the replacements appearing in the algorithm is equivalent to the application of an invertible transformation. Thus the result defines the same map as the input. Also note that replacing any column of a matrix in this form by a non-trivial linear combination of the other columns yields a matrix which is no longer in this form. Thus any two maps in this form are equivalent if and only if they have the same specifications for the remaining variables, i.e.: the result provides an affine chart defined on a Zariski dense open subset of $U_{d_1,\ldots,d_p}$.

Since, each column of degree d_i polynomials has $(d_i+1)(m+p)$ coefficients, the total number of parameters we started with was $(p+q)(m+p) = mp+q(m+p)+p^2$. Finally the algorithm specifies the p diagonal constant coefficients to 1 and introduces at least two zeros for each ordered pair $i < j$. In fact the number of zeros introduced is 2 if $d_i = d_j$ or $1 + d_j - d_i$ otherwise. Rewriting this last number as $2 + (d_j - d_i - 1)$ and adding up over all $\frac{p(p-1)}{2}$ ordered pairs $1 \leq i < j \leq p$ we find that the number of parameters specified is $p^2 + \sum_{i<j} max(d_j - d_i - 1, 0)$. The conclusion of the lemma now follows since the choice of column degrees specified is the only way to choose the d_i's so that they add up to q and no two differ from each other by more than 1. □

The normal form used above was easy to define for general $U_{d_1,\ldots,d_p}$, and was used in [47] where they use the $U_{d_1,\ldots,d_p}$ as cells in a cellation of the space of degree q-maps. For our purposes we need only work in the largest of these cells $U_{d,\ldots,d,d+1,\ldots,d+1}$. Since this cell has dimension $mp+q(m+p)$ and each root of the degree $n+q$ characteristic equation (5.9) enforces one polynomial condition on the space of degree q-maps we will expect no solutions whenever n is greater than the critical degree $mp+q(m+p)-q$ and a finite number of solutions when $n = mp+q(m+p)-q$. In both cases we expect no solutions of a generic dynamic pole placement problem to lie on these lower-dimensional pieces. In order to work on $U_{d,\ldots,d,d+1\ldots,d+1}$, however we will want to use a different normal form analogous to the form used in Section 3. To describe it we start by identifying $(m+p)$-vectors of degree d polynomials with their $(m+p)(d+1)$-element

coefficient vectors as follows:

$$\begin{bmatrix} a_1^0 + a_1^1 s + \cdots + a_1^d s^d \\ a_2^0 + a_2^1 s + \cdots + a_2^d s^d \\ \vdots \\ a_{m+p}^0 + a_{m+p}^1 s + \cdots + a_{m+p}^d s^d \end{bmatrix} \Leftrightarrow \begin{bmatrix} a_1^0 \\ a_2^0 \\ \vdots \\ a_{m+p}^0 \\ a_1^1 \\ \vdots \\ a_{m+p}^1 \\ \vdots \\ a_1^d \\ \vdots \\ a_{m+p}^d \end{bmatrix} \tag{5.10}$$

i.e.: the coefficient a_j^k of s^k in polynomial entry in row j in the representation on the left is stored in the $((k+1)(m+p)+j)$th position in the representation on the right.

DEFINITION 5.2. Let $q = dp + r$ with $d, r \in \mathbb{N}$ and $r < p$. A *localization pattern* for $(m+p) \times p$-maps of degree q is given by a table of of p columns of zeros and stars, where the first $p-r$ columns have dimension $(d+1)(m+p)$ and the remaining columns have dimensions $(d+2)$. As in the degree 0 case, all stars within a column should be contiguous and the row indices in which the bottommost and topmost stars occur strictly increase as a function of the column index. These row indices are called the *top* and *bottom* pivots respectively. It is further required that no two top (bottom) pivots differ by $m+p$ or more. We say that a map *fits* a localization pattern if it is equivalent to a map given by a matrix of polynomials with the column degrees specified by the pattern and zero coefficients everywhere the pattern requires them.

The dimension of the space of degree q-maps which meet a localization pattern X with n stars is $n-p$. To see this note that the requirements that no pivots differ by $m+p$ places ensures that no multiple of any column by a power of s has a bottommost star in a pivot row of any other column. This implies that it any linear combination involving more than one column has the lowest bottommost pivots and highest topmost pivots of the columns used. Thus only by scaling each column can we change the entries of a generic map fitting a localization pattern without producing a map which no longer fits that pattern. On the other hand we may freely scale each column to set one non-zero entry to one. This also tells us that patterns with two equal pivots or pivots that differ by $m+p$ or more will be fit by a space of dimension lower than $n-p$.

Given appropriate generality we can use invertible column operations to transform the normal form for $U_{d,\ldots,d,d+1,\ldots,d+1}$ into a localization pattern as follows: First we allow each column to be arbitrarily scaled and thus turn all ones into stars. Next, we use column operations with constant scalars on the first block of $p-r$ dimension $(d+1)(m+p)$ columns to move all the zeros below the diagonal to the bottom of the columns. We then use multiples of these columns by s to introduce $p-r$ rows of zeros to the tops of the block or r $(d+2)(m+p)$-dimensional columns. Finally we can use scalar column operations on this block to move the below diagonal elements of the normal form to the bottoms of the columns. This process is illustrated in Figure 5.3, and by counting diagonally in each block we see that the resulting localization pattern has top pivots $[1, 2, \ldots, p]$ and bottom pivots $[d(m+p)+m+r+1, \ldots, d(m+p)+$

$m+p, (d+1)(m+p)+m+1, \ldots, (d+1)(m+p)+m+1+r]$. We thus see that set of maps which fit this localization pattern has dimension $mp+q(m+p)$ and will account for all solutions of the problem of finding degree q-maps meeting $mp+q(m+p)$ generic m-planes at $mp+q(m+p)$ specified s values. In analogy to the approach used in Section 3, we will proceed by successively specializing the input planes to yield problems over smaller localization patterns involving fewer conditions.



FIG. 5.3. *Transforming standard form for $m=2$, $p=5$ and $q=3$ to top level localization pattern.*

When specializing the input planes, we also must specialize the values for s by fixing the interpolating points. For bottom pivots we focus on the highest-degree coefficients and move s to ∞ to select those coefficients. For top pivots the lowest-degree coefficients are specialized and we move s to 0. To deal with both situations we add an additional variable t to homogenize the polynomials. So we consider localization patterns of homogeneous polynomials and denote those by $X(s,t)$. The special m-planes of the hypersurface Pieri case are recycled when we evaluate $X(s,t)$ at $(s,t)=(1,0)$ for bottom pivots, or at $(s,t)=(0,1)$ for top pivots. The solutions to the problem in special position can be found in the child patterns that differ from $X(s,t)$ in exactly one position. The analogue to Definition 3.3 is immediate. The main root-counting theorem follows from the correctness of the Pieri homotopies.

THEOREM 5.3. *Consider a localization pattern $X(s,t)$ with $p+n$ stars. For $n=0$, $X(s,t)$ counts for one solution. For $n>0$, the number of maps fitting $X(s,t)$ and meeting n general m-planes at n values for (s,t) equals the sum of the number of solution maps fitting the children of $X(s,t)$ and meeting $n-1$ general m-planes at $n-1$ values for (s,t).*



FIG. 5.4. *Combinatorial root count for $p=2=m$, $q=1$. The brackets at the left are the bottom pivots. The top pivots remain [1 2]. The trivial localization pattern $X_0(s,t)$ corresponds to [4 7]. The dashed line marks the transition between $q=0$ and $q=1$. The root count is at the right.*

Note that in the definition of the Pieri homotopies we abuse notation in the sense that t denotes both the continuation parameter and the variable added to homogenize

the maps. This abuse is justified for bottom pivots because we move t from 0 to 1 to move the interpolation point from its specific value $(s,t) = (1,0)$ at ∞ to a general value $(s,t) = (s_n, 1)$. For top pivots we swap the roles of s and t.

DEFINITION 5.4. Let $X(s,t)$ be a localization pattern and S_X be the special m-plane for bottom pivots at $(s,t) = (1,0)$. Suppose that the children $Y(s,t)$ of $X(s,t)$ meet $n-1$ general m-planes L_i at (s_i,t_i), i.e.: $\det(Y(s_i,t_i)|L_i) = 0$, for $i = 1,2,\ldots,n-1$. To satisfy the nth intersection condition with L_n at (s_n,t_n), the *Pieri homotopy* is defined as

$$H(X(s,t),s,t) = \begin{cases} \det(X(s,t)|(1-t)S_X + tL_n) = 0 \\ (s-1)(1-t) + (s-s_n)t = 0 \\ \det(X(s_i,t_i)|L_i) = 0 \\ i = 1,2,\ldots,n-1 \end{cases} \qquad t \in [0,1]. \tag{5.11}$$

For top pivots s and t swap roles and we consider s_n^{-1} as the target value for s. When we remove entries from both top and bottom, two independent s-variables are used. Theorem 5.3 follows from the correctness of the Pieri homotopy, proven in Lemma 5.5.

LEMMA 5.5. *Consider a localization pattern $X(s,t)$ with $p+n$ stars and n complex m-planes L_i in general position at the values (s_i,t_i). Suppose we are given all maps that meet L_i at (s_i,t_i), $i = 1,2,\ldots,n-1$ and fit one of the children of $X(s,t)$. Then the Pieri homotopy defines regular solution paths as t varies from 0 to 1, that start at the given maps and end at those maps that meet all n general m-planes L_i at (s_i,t_i) while fitting $X(s,t)$.*

Proof. Observe that in the Pieri homotopy (5.11) at $(s,t) = (1,0)$ we select in the first equation with the special m-plane S_X in the expansion of $\det(X(1,0)|S_X) = 0$ those stars that are indexed by the bottom pivots. So the first equation in (5.11) looks like

$$H_1(X(s,t),s,t) = p(X(1,0)) + t(f(X(s,t),s,t) + O(t)), \tag{5.12}$$

where $p(X(1,0))$ denotes the product of all pivot stars and where $f(X(s,t),s,t)$ is a polynomial not divisible by t. Ordering the variables as (s,t,x_{ij}), the matrix of all partial derivatives of the homotopy has the following structure:

$$\begin{bmatrix} \frac{\partial H_1}{\partial s} & \frac{\partial H_1}{\partial t} & A \\ 1 & 1-s_n & B \\ 0 & 0 & C \end{bmatrix}. \tag{5.13}$$

Since the $n-1$ planes L_i are in general position the rank of C is $n-1$. Thus the Jacobian matrix (5.13) is of full rank if and only if $\frac{\partial H_1}{\partial s}\frac{\partial H_1}{\partial t} \neq 0$ at $(s,t) = (1,0)$, We can show that $f(X(1,0),1,0) \neq 0$, which implies $\frac{\partial H_1}{\partial t} \neq 0$. If $f(X(1,0),1,0)$ were equal to 0 then $f(X(s,t),s,t)$ would be identically zero and H_1 could not contain any term that is linear in t. This could only happen for very special choices of L_n. Hence at $(s,t) = (1,0)$, all solutions are regular.

To compactify our space we choose again a multi-homogenization according to the columns of the matrices. The regularity of the homotopy is maintained as t moves to 1 since L_n is in general position. Suppose we would have more solutions at $(s,t) = (s_n,1)$ than the number of paths we have started with. Going backwards, when in the Pieri homotopy we let t go from 1 to 0, we would arrive at solutions at $(s,t) = (1,0)$ that do not fit any of the child patterns of $X(s,t)$. But any other child pattern that is not provided in the root count does not allow enough freedom to meet the general intersection conditions L_i at (s_i,t_i) for $i = 1,2,\ldots,n-1$. ∎

We can reformulate the Pieri rule for the general case, to deal with general intersection conditions for any q, when the codimensions conditions k_i, $i = 1, 2, \ldots, n$ satisfy $k_1 + k_2 + \ldots + k_n = mp + q(m+p)$. In the case where $q > 0$ we work with brackets modulo $(m+p)$ to represent the top and bottom pivots.

Consider a chain C ending at the pattern $X(s,t)$. Let α denote the bracket with the indices of the basis vectors in the intersection of the special m-planes for the patterns in the chain C. Let β denote the pivots modulo $(m+p)$ for a child $Y(s,t)$ of $X(s,t)$. Define γ as the bracket obtained from concatenating α and β, sorted in increasing order. Note that γ may contain repeated entries. We say that $Y(s,t)$ is *a valid child of the chain* C if there exists an index i such that

$$|\gamma_i - \gamma_1| < i - 1, \quad \text{for bottom pivots } \beta \tag{5.14}$$

or

$$|\gamma_l - \gamma_i| < l - i, \quad \text{for top pivots } \beta, \tag{5.15}$$

where l is the index of the last element in γ. This rule is consistent with the general Pieri rule for $q = 0$ and allows us to compute intersection numbers for any q and for general codimension conditions.

6. Computational Experiments and Experiences. The Pieri homotopy algorithm is part of the module of PHCpack [44] that also provides the SAGBI homotopies. We compare with those homotopies and give comments on the organization of the code, numerical aspects, and applications. All timings reported concern a 166 MHz Pentium II processor with 64 Mb internal memory running Linux, unless indicated otherwise.

6.1. Implementation Issues. In setting up the Pieri homotopy algorithm, we restricted to the case where the input planes are all in general position. In this case, all the deformations are free from multiplicities. To solve a nongeneric, specific instance of the problem instance we first solve a generic case and then apply a cheater's homotopy to obtain solutions to the original problem. The paradigm of cheater's homotopy ([20], [21]) or coefficient-parameter polynomial continuation ([25], [26]) makes it straightforward to set up this homotopy. This paradigm ensures that singularities can only occur at the end of the solution paths, and only when the original problem has singular solutions.

The general intersection conditions are represented by overdetermined polynomial systems. A direct method to solve an overconstrained polynomial system of N equations in n unknowns is to multiply it by an $n \times N$-matrix of random complex numbers to construct a square system, as is proposed in [36]. This reduction to a square system destroys the geometric structure and creates many excess solution paths to follow. In our situation, the overdetermined polynomial systems we have to deal with are the Pieri homotopies for general intersection conditions, which are overdetermined homotopies. From the geometry we know which solution paths we have to follow so that we maintain the optimal number of solution paths, see [16, Section 4]. To control the correctness and to detect numerical failures, we evaluate the original overconstrained system at the computed end points of the paths. The complexity of Newton's method and path-following algorithms for overdetermined systems is analyzed in [9].

The geometric intersection conditions treated in Section 4 are expressed by determinantal equations, which are a special kind of polynomial systems. Just as sparse

polynomial systems allow fewer perturbations on the coefficient structure than general polynomial systems (whence they have a better numerical condition [8] and can be more easily solved), we expect that the polynomial systems arising from these geometric intersection conditions are numerically easier to solve. We have done some tests on working directly with the matrix structures, but abandoned this approach because the general linear algebra operations turn out to be slower than the nested Horner schemes that are used to evaluate the expanded determinantal forms. Nevertheless, we expect computational progress from Newton's method adapted to this specific geometric situation [11], in combination with secant methods [49].

6.2. Computational and numerical experiments. The description in [16] leads to path following in a chain-by-chain fashion as defined by Pieri trees. Contemplating the posets in Figures 3.2 and 4.1 one sees that one can avoid the tracing of many solution paths when the continuation matches the poset structure. To solve a generic complex instance encoded by $(m,p) = (4,3)$, the chain-by-chain Pieri algorithm takes 4 h 15 m 58 s 220 ms and has to trace 2310 solution paths. The new poset-oriented Pieri homotopies only take 1 h 7 m 20 s 160 ms (see Table 6.1) to trace 1663 paths. This savings are only partially explained by the reduction in the number of paths. The poset structure allows us to organize the symbolic homotopy constructors in a much more economical way.

In Table 6.1 we summarize the computational results with pivoting done in a mixed fashion. Here we are removing pivots from both the top and the bottom, as in the original chain-by-chain original description in [16]. In this case, the Pieri homotopy has two moving equations:

$$H(X,t) = \begin{cases} \det(X|(1-t)S_X^t + tL_n) &= 0 \\ \det(X|(1-t)S_X^b + tL_{n-1}) &= 0 \\ \det(X|L_i) &= 0 \\ & i = 1,2,\ldots,n-2 \end{cases} \qquad t \in [0,1], \tag{6.1}$$

where S_X^t and S_X^b are the special m-planes respectively for top and bottom pivots of X. The input 4-planes for the real target problem are osculating a rational normal curve, as prescribed in one of the Shapiro-Shapiro conjectures ([32], [39], [45]). For these inputs, all solution 3-planes turn out to be real on the examples we ran, in accordance with the Shapiro-Shapiro conjectures. Dividing the total time on the last line in Table 6.1 by 462, we arrive at 48 s 476 ms cpu time as cost for one solution, which is the cost for one feedback law.

Note that one can change basis to bring two input 4-planes into special position and thus save the time spent at stage $n = 12$ in Table 6.1. Another optimization involves using linear algebra instead of continuation to satisfy the intersection conditions at the top level of the poset, see [16, Proposition 3.1]. But this optimization only cuts off milliseconds.

The input planes are represented by matrices of generators which are equivalent up to a transformation of basis. We have noticed that an orthonormal representation of the input planes – filtered through the QR-algorithm [14] – is responsible for considerable computational savings and an increased numerical accuracy of the solution planes. The difference between random and orthonormal inputs becomes especially significant for higher-dimensional problems. For instance, the case $(m,p) = (5,3)$ leads to a fifteen-dimensional polynomial system of cubics with 6006 solutions. A 450MHz Intel Pentium machine needs 16 h 17 m 16 s cpu time on random inputs, whereas it only takes 10 h 12 m 38 s on orthonormal input planes.

n	#paths	user cpu time
2	18	800 ms
4	72	6 s 918 ms
6	187	1 m 8 s 40 ms
8	462	8 m 27 s 460 ms
10	462	18 m 46 s 390 ms
12	462	38 m 42 s 360 ms
total	1663	1 h 7 m 20 s 160 ms
target	462	5 h 4 m 46 s 610 ms
total	2125	6 h 13 m 15 s 840 ms

TABLE 6.1

Number of paths and timings for $(m,p) = (4,3)$ at every even level n, up to the total resolution time for a generic complex intersection problem. The first total is added to the time needed for solving a real target instance and results in the last line.

6.3. Comparison with SAGBI Homotopies. The SAGBI homotopies were proposed in [16] to compute all p-planes that meet mp given m-planes in general position. The second author implemented those homotopies to verify large instances of the Shapiro-Shapiro conjectures. See [45] for a description and [32] and [39, 41] for other tests and related work on these conjectures.

To solve a generic complex instance, encoded by $(m,p) = (4,3)$, the Pieri homotopy algorithm takes 1 h 7 m 20 s 160 ms (see Table 6.1). As reported in [45], SAGBI homotopies need 2 h 23 m 27 s 630 ms for the same problem using the localization patterns that lead to polynomials of lowest degree, and 5 h 23 m 36 s 840 ms with localization patterns used by Pieri homotopies. The common optimization of bringing two input planes into special position to diminish the dimension with two forces those localization patterns that lead to higher-degree polynomials, increasing the complexity of SAGBI homotopies.

SAGBI homotopies suffer from two drawbacks, not shared by the Pieri homotopies. Firstly, they cannot solve the more general intersection problem treated in Section 4. Secondly, a polyhedral continuation is needed to start up the SAGBI homotopies. These continuations happen in the top dimension, which makes them computationally intensive. In addition, these homotopies are both highly nonlinear in the continuation parameter, which leads to numerical instabilities as the problems get larger. Although the techniques in [12] help, they always involve a lower bound that may be too high for the machine precision.

To conclude the comparison, we contrast the analytic geometry as used in the SAGBI homotopies with the intrinsic geometry of the Pieri homotopies. With the SAGBI homotopies we loose all geometric information after the transformation of the problem into a polynomial system. The Pieri homotopies use the polynomial equations merely as a means to compute and are better able to exploit the evolving geometry.

6.4. Reality Issues. The Pieri homotopies arose from the question [37, 38] whether a problem of enumerative geometry can have all its solutions real. The developed software may be useful for verifying instances of conjectures such as the Shapiro-Shapiro conjectures, see [32], [39, 41], and [45]. Note that for the quantum Schubert calculus a counterexample to one of the analogues of the Shapiro-Shapiro conjectures is given in [40] for the case $q = 1$, $m = 2 = p$. For engineering applications,

where the machines are specified with real matrices, only real feedback laws are of interest. Several explicit root counting formulas are given in [29] which allow one to quickly determine the parity of the intersection numbers.

We end this paper describing some computational experiences with random real inputs. Table 6.2 gives the number of paths and timings reported for solving the case of $m = 3$, $p = 2$, and $q = 1$. For this random real instance, 25 real and 30 complex maps were found. The timings are for a 350 Mhz Intel II Pentium processor, with 64 Mb internal memory, running SunOs 5.7.

n	#paths	user cpu time
1	3	$<$ 1ms
3	8	240ms
5	21	1s 690ms
7	55	18s 390ms
9	55	35s 560ms
11	55	1m 16s 560ms
total	197	2m 12s 880ms
target	55	2m 25s 280ms
total	252	4m 39s 760ms

TABLE 6.2

Number of paths and timings for $(m,p) = (3,2)$ for $q = 1$, at every odd level n. The first total concerns the time needed for the quantum Pieri homotopy algorithm on a random complex instance. Note that this is less than the time needed to solve a random real target system.

In the last example we indicate what could be done when the number of solution is even, following the spirit of Frank Sottile's original ideas [37]. In the hypersurface case for $(m,p) = (4,3)$, there are 462 solution 3-planes. Since 462 is even, there might be a chance that all of them are complex. But suppose you specialize the input somewhat, so that we have four sets of three 4-planes that each meet at one line (=2-plane) common to their set. Then we may replace the original hypersurface intersection conditions by the requirement that the solution 3-planes meet those four common 2-planes. Each of these four 2-planes gives a codimension 3 condition and as $12 = 3+3+3+3$ we can apply the general Pieri algorithm which gives exactly one solution, which is real for real inputs. Note that this involves the resolution of a polynomial system of 84 cubic equations in 12 variables. Table 6.3 summarizes the computations.

Acknowledgments. We are indebted to Frank Sottile for the many explanations that helped us through the first implementation and for the hint that Pieri homotopies exist for the quantum Schubert calculus. The explanations of Joachim Rosenthal were most helpful. We thank MSRI for the opportunity of sharing an office for one full year. Frank Sottile and Mengnien Wu gave valuable comments on the preprint version of this paper. We are grateful for the remarks of the anonymous referees.

REFERENCES

[1] E.L. Allgower and K. Georg. *Numerical Continuation Methods, an Introduction*, volume 13 of *Springer Ser. in Comput. Math.* Springer–Verlag, Berlin Heidelberg New York, 1990.

[2] E.L. Allgower and K. Georg. Numerical Path Following. In P.G. Ciarlet and J.L. Lions, editors, *Techniques of Scientific Computing (Part 2)*, volume 5 of *Handbook of Numerical Analysis*, pages 3–203. North-Holland, 1997.

n	#paths	user cpu time
2	1	20ms
4	1	190ms
6	1	570ms
8	1	6s 800ms
10	1	4s 910ms
12	1	22s 780ms
total	6	38s 190ms
target	1	19s 500ms
total	7	1m 25s 340ms

TABLE 6.3

Number of paths and timings for $(m,p) = (4,3)$, $q = 0$, and $k_i = 3$, for $i = 1,2,3,4$, listed at every even level. The timings for the general Pieri homotopy algorithm, on a random complex instances are added up into the first total. The last total includes the resolution of a random real target system.

[3] S. Ariki. Generic pole assignment via dynamic feedback. *SIAM J. Control and Optimization*, 36(1):379–390, 1998.

[4] A. Bertram. Quantum Schubert calculus. *Adv. Math.*, 128(2):289–305, 1997.

[5] R.W. Brockett and C.I. Byrnes. Multivariate Nyquist criteria, root loci, and pole placement: a geometric viewpoint. *IEEE Trans. Automat. Control*, 26:271–284, 1981.

[6] C.I. Byrnes. Pole assignment by output feedback. In H. Nijmacher and J.M. Schumacher, editors, *Three Decades of Mathematical Systems Theory*, volume 135 of *Lecture Notes in Control and Inform. Sci.*, pages 13–78. Springer–Verlag, Berlin, 1989.

[7] D.A. Cox and S. Katz. *Mirror Symmetry and Algebraic Geometry*, volume 68 of *Mathematical Surveys and Monographs*. AMS, Providence, R.I., 1999.

[8] J.-P. Dedieu. Condition number analysis for sparse polynomial systems. In F. Cucker and M. Shub, editors, *Foundations of Computational Mathematics*, pages 75–101. Springer, 1997.

[9] J.P. Dedieu and M. Shub. Newton and predictor-corrector methods for over-determined systems of equations. IBM Technical Report, IBM Research Division, 1998. To appear in *Math. Comp.*

[10] D.F. Delchamps. *State Space and Input-Output Linear Systems*. Springer–Verlag, Berlin, 1988.

[11] A. Edelman, T. Arias, and S.T. Smith. The geometry of algorithms with orthogonality constraints. *SIAM J. Matrix Anal. Appl.*, 20(2):303–353, 1998.

[12] T. Gao, T.Y. Li, J. Verschelde, and M. Wu. Balancing the lifting values to improve the numerical stability of polyhedral homotopy continuation methods. To appear in *Applied Math. Comput.*

[13] I. Gohberg, M.A. Kaashoek, and F. van Schagen. *Partially Specified Matrices and Operators: Classification, Completion, Applications*, volume 79 of *Operator Theory, Advances and Applications*. Birkhäuser, Boston, 1995.

[14] G. Golub and C. Van Loan. *Matrix Computations*. The Johns Hopkins University Press, Baltimore and London, third edition, 1996.

[15] W.V.D. Hodge and D. Pedoe. *Methods of Algebraic Geometry, Volume 2*. Cambridge University Press, 1994. Paperback reprint of the 1952 edition.

[16] B. Huber, F. Sottile, and B. Sturmfels. Numerical Schubert calculus. *J. of Symbolic Computation*, 26(6):767–788, 1998.

[17] T. Kailath. *Linear Systems*. Prentice-Hall, Inc., Englewood Cliffs, N.J., 1980.

[18] S. Kleiman and D. Laksov. Schubert calculus. *American Mathematical Monthly*, 79(10):1061–1082, 1972.

[19] T.Y. Li. Numerical solution of multivariate polynomial systems by homotopy continuation methods. *Acta Numerica*, 6:399–436, 1997.

[20] T.Y. Li, T. Sauer, and J.A. Yorke. The cheater's homotopy: an efficient procedure for solving systems of polynomial equations. *SIAM J. Numer. Anal.*, 26(5):1241–1251, 1989.

[21] T.Y. Li and X. Wang. Nonlinear homotopies for solving deficient polynomial systems with parameters. *SIAM J. Numer. Anal.*, 29(4):1104–1118, 1992.

[22] A. Morgan. *Solving polynomial systems using continuation for engineering and scientific prob-*

lems. Prentice-Hall, Englewood Cliffs, N.J., 1987.

[23] A. Morgan and A. Sommese. Computing all solutions to polynomial systems using homotopy continuation. *Appl. Math. Comput.*, 24(2):115–138, 1987.

[24] A. Morgan and A. Sommese. A homotopy for solving general polynomial systems that respects m-homogeneous structures. *Appl. Math. Comput.*, 24(2):101–113, 1987.

[25] A.P. Morgan and A.J. Sommese. Coefficient-parameter polynomial continuation. *Appl. Math. Comput.*, 29(2):123–160, 1989. Errata: *Appl. Math. Comput.* 51:207(1992).

[26] A.P. Morgan and A.J. Sommese. Generically nonsingular polynomial continuation. In E.L. Allgower and K. Georg, editors, *Computational Solution of Nonlinear Systems of Equations*, pages 467–493, Providence, R.I., 1990. AMS.

[27] M. Pieri. Formule di coincidenza per le serie algebriche ∞^n di coppie di punti dello spazio a n dimensioni. *Rend. Circ. Mat. Palermo*, 5:252–268, 1891.

[28] M.S. Ravi and J. Rosenthal. A smooth compactification of the space of transfer functions with fixed McMillan degree. *Acta Applicandae Mathematicae*, 34:329–352, 1994.

[29] M.S. Ravi, J. Rosenthal, and X. Wang. Dynamic pole placement assignment and Schubert calculus. *SIAM J. Control and Optimization*, 34(3):813–832, 1996.

[30] M.S. Ravi, J. Rosenthal, and X. Wang. Degree of the generalized Plücker embedding of a quot scheme and quatum cohomology. *Math. Ann.*, 311:11–26, 1998.

[31] J. Rosenthal. On dynamic feedback compensation and compactifications of systems. *SIAM J. Control and Optimization*, 32(1):279–296, 1994.

[32] J. Rosenthal and F. Sottile. Some remarks on real and complex output feedback. *Systems and Control Lett.*, 33(2):73–80, 1998. See `http://www.nd.edu/~rosen/pole` for a description of computational aspects of the paper.

[33] J. Rosenthal and J. C. Willems. Open problems in the area of pole placement. In V.D. Blondel, E.D. Sontag, M. Vidyasagar, and J.C. Willems, editors, *Open Problems in Mathematical Systems and Control Theory*, Communication and Control Engineering Series, pages 181–191. Springer–Verlag, Berlin, 1998.

[34] H. Schubert. *Kalkül der abzählenden Geometrie*. Teubner, Leipzig, 1879. Reprinted in 1979 by Springer–Verlag with an Introduction by S.L. Kleiman.

[35] H. Schubert. Beziehungen zwischen den linearen Räumen auferlegbaren charakteristischen Bedingungen. *Math. Ann.*, 38:588–602, 1891.

[36] A.J. Sommese and C.W. Wampler. Numerical algebraic geometry. In J. Renegar, M. Shub, and S. Smale, editors, *The Mathematics of Numerical Analysis*, volume 32 of *Lectures in Applied Mathematics*, pages 749–763, 1996. Proceedings of the AMS-SIAM Summer Seminar in Applied Mathematics, Park City, Utah, July 17-August 11, 1995, Park City, Utah.

[37] F. Sottile. Enumerative geometry for real varieties. In J. Kollár, R. Lazarsfeld, and D. R. Morrison, editors, *Algebraic Geometry - Santa Cruz 1995 (University of California, Santa Cruz, July 1995)*, volume 62, Part I of *Proceedings of Symposia in Pure Mathematics*, pages 435–447. AMS, 1997.

[38] F. Sottile. Pieri's formula via explicit rational equivalence. *Can. J. Math.*, 49(6):1281–1298, 1997.

[39] F. Sottile. Real Schubert calculus: polynomial systems and a conjecture of Shapiro and Shapiro. Preprint #1998-066, MSRI, 1998. Accepted for publication in *Experimental Mathematics*.

[40] F. Sottile. Real rational curves in Grassmannians. Preprint #1999-025, MSRI, 1999. To appear in the *Journal of the Americal Mathematical Society*.

[41] F. Sottile. The special Schubert calculus is real. *Electronic Research Announcements of the AMS*, 5:35–39, 1999. Available at `http://www.ams.org/era`.

[42] F. Sottile and B. Sturmfels. A SAGBI basis for the quantum Grassmannian. Preprint #1999-043, MSRI, 1998.

[43] R.P. Stanley. Some combinatorial aspects of the Schubert calculus. In D. Foata, editor, *Combinatoire et Représentation du groupe symétrique, Strasbourg, 1976*, volume 579 of *Lecture Notes in Mathematics*, pages 217–251. Springer–Verlag, Berlin, 1977.

[44] J. Verschelde. Algorithm 795: PHCpack: A general-purpose solver for polynomial systems by homotopy continuation. To appear in *ACM Trans. Math. Softw.*, vol 25, no 2, 1999. Paper and software available at the author's web-pages.

[45] J. Verschelde. Numerical evidence for a conjecture in real algebraic geometry. Preprint #1998-064, MSRI, 1998. Accepted for publication in *Experimental Mathematics*. Paper and software available at the author's web-pages.

[46] X. Wang, M.S. Ravi, and J. Rosenthal. Algebraic and combinatorial aspects of the dynamic pole assignment problem. In U. Helmke, R. Mennicken, and J. Saurer, editors, *Systems and Networks: Mathematical Theory and Applications*, volume 79 of *Mathematical Research*,

pages 547–550. Akademie Verlag, Berlin, 1994. Proc. of the International Symposium MTNS 93 held in Regensburg Germany. Vol II: Invited and contributed papers.

[47] X. Wang and J. Rosenthal. A cell structure for the set of autoregressive systems. *Linear Algebra and its Applications*, 205–206:1203–1225, 1994.

[48] J.C. Willems. Paradigms and puzzles in the theory of dynamical systems. *IEEE Transactions on Automatic Control*, 36(3):259–294, 1991.

[49] J.C. Yakoubsohn. Finding zeros of analytic functions: α-theory for secant type methods. *J. Complexity*, 15(2):239–281, 1999.

[50] H.G. Zeuthen. *Lehrbuch der abzählenden Methoden der Geometrie*, volume XXXIX of *Lehrbücher der mathematischen Wissenschaften*. Teubner, Leipzig, 1914.