

Sum of Squares

Nonnegative Polynomials

positive semidefinite matrices

Applications

control of nonlinear systems and global optimization

Sum of Squares and Radical Ideals

real radicals and a real Nullstellensatz

Moment Matrices

counting the number of real solutions

- 1 Nonnegative Polynomials
positive semidefinite matrices
- 2 Applications
control of nonlinear systems and global optimization
- 3 Sum of Squares and Radical Ideals
real radicals and a real Nullstellensatz
- 4 Moment Matrices
counting the number of real solutions

MCS 563 Lecture 30
Analytic Symbolic Computation
Jan Verschelde, 4 April 2011

Sum of Squares

Nonnegative Polynomials

positive semidefinite matrices

Applications

control of nonlinear systems and global optimization

Sum of Squares and Radical Ideals

real radicals and a real Nullstellensatz

Moment Matrices

counting the number of real solutions

- 1 Nonnegative Polynomials
positive semidefinite matrices
- 2 Applications
control of nonlinear systems and global optimization
- 3 Sum of Squares and Radical Ideals
real radicals and a real Nullstellensatz
- 4 Moment Matrices
counting the number of real solutions

sum of squares

Nonnegative
Polynomialspositive semidefinite
matrices

Applications

control of nonlinear
systems and global
optimizationSum of
Squares and
Radical Idealsreal radicals and a
real NullstellensatzMoment
Matricescounting the number
of real solutions

Consider a real polynomial $p \in \mathbb{R}[\mathbf{x}]$.

If p can be written as a sum of squares, i.e.:

- if there exist r polynomials $q_i \in \mathbb{R}[\mathbf{x}]$, $i = 1, 2, \dots, r$,
- such that $p = q_1^2 + q_2^2 + \dots + q_r^2$,

then $p(\mathbf{x}) \geq 0$, for all $\mathbf{x} \in \mathbb{R}$.

Moreover, for any $\epsilon > 0$, $p + \epsilon > 0$
is positive and has no real roots.

Not every nonnegative polynomial is a sum of squares.

positive semidefinite matrices

Nonnegative Polynomials

positive semidefinite
matrices

Applications

control of nonlinear
systems and global
optimization

Sum of Squares and Radical Ideals

real radicals and a
real Nullstellensatz

Moment Matrices

counting the number
of real solutions

We make the connection between
sum of squares and positive semidefinite matrices.

The matrix A is positive semidefinite
if for all vectors \mathbf{x} : $\mathbf{x}^T A \mathbf{x} \geq 0$.

A symmetric positive semidefinite matrix A
can be decomposed as $A = R^T R$,
where R is an upper triangular matrix.

This is known as the Cholesky decomposition of A .

representing polynomials

Nonnegative Polynomials

positive semidefinite matrices

Applications

control of nonlinear systems and global optimization

Sum of Squares and Radical Ideals

real radicals and a real Nullstellensatz

Moment Matrices

counting the number of real solutions

We use matrices to represent polynomials.

Take as p for example a fourth degree polynomial, which we may represent as

$$p(x) = [x^2 \ x \ 1] \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{12} & a_{22} & a_{23} \\ a_{13} & a_{23} & a_{33} \end{bmatrix} \begin{bmatrix} x^2 \\ x \\ 1 \end{bmatrix} = \mathbf{x}^T A \mathbf{x}.$$

If A is positive semidefinite, then its Cholesky decomposition $A = R^T R$ gives

$$p(x) = \mathbf{x}^T A \mathbf{x} = \mathbf{x}^T R^T R \mathbf{x} = (R \mathbf{x})^T (R \mathbf{x}) = y_1^2 + y_2^2 + y_3^3,$$

with $\mathbf{y} = R \mathbf{x}$.

semidefinite programming

Nonnegative
Polynomialspositive semidefinite
matrices

Applications

control of nonlinear
systems and global
optimizationSum of
Squares and
Radical Idealsreal radicals and a
real NullstellensatzMoment
Matricescounting the number
of real solutions

The positive definite cone (or PSD cone)
of all real symmetric positive semidefinite n -by- n matrices is

$$PSD(\mathbb{R}^n) = \{ A \in \mathbb{R}^{n \times n} \mid A^T = A \text{ and } \forall \mathbf{x} : \mathbf{x}^T A \mathbf{x} \geq 0 \}.$$

That $PSD(\mathbb{R}^n)$ is a cone follows from $\mathbf{x}^T A \mathbf{x} \geq 0$.

Semidefinite programming (or SDP for short) concerns

SDP decision: given r linear functions l_1, l_2, \dots, l_r ,
does there exist an $A \in PSD(\mathbb{R}^n)$ such that
 $l_i(A) = 0$, for $i = 1, 2, \dots, r$?

SDP optimization: for $r + 1$ linear functions l_0, l_1, \dots, l_r ,
minimize $l_0(A)$ subject to $A \in PSD(\mathbb{R}^n)$ and
 $l_i(A) = 0$, for $i = 1, 2, \dots, r$.

Interior-point methods provide efficient algorithms.

Nonnegative
Polynomialspositive semidefinite
matrices

Applications

control of nonlinear
systems and global
optimizationSum of
Squares and
Radical Idealsreal radicals and a
real NullstellensatzMoment
Matricescounting the number
of real solutions

For the polynomial $p(x) = x^4 - x^2 - 2x + 2$, we consider

$$\mathbf{x}^T A \mathbf{x} = a_{11}x^4 + 2a_{12}x^3 + (2a_{13} + a_{22})x^2 + a_{23}x + a_{33}.$$

Now we want to find those $A \in PSD(\mathbb{R}^n)$ satisfying the linear equations $a_{11} = 1$, $a_{12} = 0$, $2a_{13} + a_{22} = -1$, $a_{23} = -1$, and $a_{33} = 2$.

This SDP decision problem has a unique solution so that p can be written as a sum of squares.

Consider for example the set defined by $1 \leq x^2 + y^2 \leq 2$.
This is clearly not a convex set.

Can we use convex optimization to compute with this set?

$$\ell : \mathbb{R}^2 \rightarrow \mathbb{R}^3 : (x, y) \mapsto (x, y, x^2 + y^2)$$

The idea is to view the set defined by $1 \leq x^2 + y^2 \leq 2$
as the projection of the convex set

$$\{ (x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 \leq z, 1 \leq z \leq 4 \}.$$

This lifting is a useful relaxation technique.

Sum of Squares

Nonnegative Polynomials

positive semidefinite matrices

Applications

control of nonlinear systems and global optimization

Sum of Squares and Radical Ideals

real radicals and a real Nullstellensatz

Moment Matrices

counting the number of real solutions

- 1 Nonnegative Polynomials
positive semidefinite matrices
- 2 Applications
control of nonlinear systems and global optimization
- 3 Sum of Squares and Radical Ideals
real radicals and a real Nullstellensatz
- 4 Moment Matrices
counting the number of real solutions

control of nonlinear systems

Nonnegative
Polynomialspositive semidefinite
matrices

Applications

control of nonlinear
systems and global
optimizationSum of
Squares and
Radical Idealsreal radicals and a
real NullstellensatzMoment
Matricescounting the number
of real solutions

Consider a nonlinear system

$$\frac{\partial \mathbf{x}(t)}{\partial t} = f(\mathbf{x}(t)),$$

where $f(\mathbf{x})$ is a real polynomial system in \mathbf{x} .

To prove asymptotic stability of $\mathbf{x} = \mathbf{0}$
as a fixed point of this nonlinear system,
we must find a Lyapunov function $V(\mathbf{x})$ satisfying

$$\text{for all } \mathbf{x} \neq \mathbf{0} : V(\mathbf{x}) > \mathbf{0} \text{ and } \frac{\partial V(\mathbf{x}(t))}{\partial t} = \left(\frac{\partial V}{\partial \mathbf{x}} \right)^T f(\mathbf{x}) < 0,$$

for all nonzero \mathbf{x} in a neighborhood of the origin.

certificate for positivity

Nonnegative
Polynomialspositive semidefinite
matrices

Applications

control of nonlinear
systems and global
optimizationSum of
Squares and
Radical Idealsreal radicals and a
real NullstellensatzMoment
Matricescounting the number
of real solutions

Expressing $V(\mathbf{x})$ as a sum of squares gives an explicit way of showing its positivity.

To automate the search for a Lyapunov function, parameters λ are introduced, so $V = V(\mathbf{x}, \lambda)$.

For \mathbf{z} a monomial vector, we then represent V as a quadratic form in \mathbf{z} : $V(\mathbf{x}, \lambda) = \mathbf{z}^T Q(\lambda) \mathbf{z}$.

The parameters λ are then determined so that $Q(\lambda)$ is positive semidefinite.

global optimization

Nonnegative
Polynomialspositive semidefinite
matrices

Applications

control of nonlinear
systems and global
optimizationSum of
Squares and
Radical Idealsreal radicals and a
real NullstellensatzMoment
Matricescounting the number
of real solutions

Via semidefinite programming we may find the global minimum of a polynomial function f on \mathbb{R}^n .

Without semidefinite programming, we could compute all critical points of the system defined by all partial derivatives

$$\frac{\partial f}{\partial x_i}(\mathbf{x}) = 0, \quad \text{for } i = 1, 2, \dots, n.$$

The number of solutions of this system grows exponentially in n .

Solving this system of partial derivatives is feasible only for a very modest number of variables and for low degree polynomials.

a relaxation

Instead, we consider a relaxation of the problem:

SOS Relaxation: Find the largest $\lambda \in \mathbb{R}$ such that

$f(x_1, x_2, \dots, x_n) - \lambda$ is a sum of squares.

The dimension of the vector space N for which we consider $PSD(\mathbb{R}^N)$ is proportional to the number of monomials in f , i.e. it grows like $\binom{n+d}{d}$, where $d = \deg(f)$.

Although also this dimension grows exponentially, the growth is more moderate compared to the system of partial derivatives.

Sum of Squares

Nonnegative Polynomials

positive semidefinite matrices

Applications

control of nonlinear systems and global optimization

Sum of Squares and Radical Ideals

real radicals and a real Nullstellensatz

Moment Matrices

counting the number of real solutions

- 1 Nonnegative Polynomials
positive semidefinite matrices
- 2 Applications
control of nonlinear systems and global optimization
- 3 Sum of Squares and Radical Ideals
real radicals and a real Nullstellensatz
- 4 Moment Matrices
counting the number of real solutions

Nonnegative
Polynomialspositive semidefinite
matrices

Applications

control of nonlinear
systems and global
optimizationSum of
Squares and
Radical Idealsreal radicals and a
real NullstellensatzMoment
Matricescounting the number
of real solutions

The definition of radical of an ideal I is

$$\sqrt{I} = \{ f \in \mathbb{R}[\mathbf{x}] \mid f^k \in I \text{ for some integer } k \geq 1 \}.$$

Hilbert's Nullstellensatz states $\sqrt{I} = I(V)$ for a variety V .

In words:

the radical of an ideal is the vanishing ideal of the solutions.

Lemma 1 gives a criterion for an ideal to be radical, limiting the powers to squares, justifying the \sqrt{I} notation.

Lemma 1

Let I be an ideal in $\mathbb{R}[\mathbf{x}]$. I is radical if and only if

$$\text{for all } f \in \mathbb{R}[\mathbf{x}] : f^2 \in I \Rightarrow f \in I. \quad (1)$$

Proof. First, consider the \Rightarrow in the *if and only if*, assuming I is radical. If I is radical, then for all integer $k \geq 1$: $f^k \in I$ implies $f \in I$. Take $k = 2$ and (1) follows.

Second, to show the \Leftarrow in the *if and only if*, assuming (1) holds, we show that $f \in I$, if $f^k \in I$, for some integer $k \geq 1$. For example, let $k = 4$: if $f^4 \in I$, then $(f^2)^2 \in I$ implies $f^2 \in I$, using (1). Applying (1) once more results in $f \in I$. For $k = 2^\ell$, applying (1) ℓ times yields $f \in I$. If k is not a power of 2, we multiply f^k sufficiently many times with f till we have $f^{2^\ell} \in I$. □

Nonnegative
Polynomialspositive semidefinite
matrices

Applications

control of nonlinear
systems and global
optimizationSum of
Squares and
Radical Idealsreal radicals and a
real NullstellensatzMoment
Matricescounting the number
of real solutions

The real radical of an ideal I is

$$\sqrt[\mathbb{R}]{I} = \left\{ f \in \mathbb{R}[\mathbf{x}] \mid f^{2k} + \sum_{j=1}^m p_j^2 \in I \text{ for some } k \geq 1, p_j \in \mathbb{R}[\mathbf{x}] \right\}.$$

Similar to Lemma 1, we have a criterion for an ideal to be real radical in Lemma 2.

critterion on real radical

Nonnegative
Polynomials

positive semidefinite
matrices

Applications

control of nonlinear
systems and global
optimization

Sum of
Squares and
Radical Ideals

real radicals and a
real Nullstellensatz

Moment
Matrices

counting the number
of real solutions

Lemma 2

Let I be an ideal in $\mathbb{R}[\mathbf{x}]$. I is real radical if and only if

$$\text{for all } p_j \in \mathbb{R}[\mathbf{x}] : \sum_{j=1}^m p_j^2 \in I \Rightarrow p_j \in I. \quad (2)$$

Proof. First, consider the \Rightarrow in the *if and only if*, assuming I is real radical. If I is real radical, i.e.: $I = \sqrt[\mathbb{R}]{I}$, then for all

integer $k \geq 1$ and $p_j \in \mathbb{R}[\mathbf{x}]$: $f^{2k} + \sum_{j=1}^m p_j^2 \in I$ implies $f \in I$.

Take $k = 2$ and (2) follows.

Second, to show the \Leftarrow in the *if and only if*, assuming (2)

holds. Let $f, p_j \in \mathbb{R}[\mathbf{x}]$: $f^{2k} + \sum_{j=1}^m p_j^2 \in I$ holds. By (2):

$f^k, p_j \in I$. As (2) implies (1), Lemma 1 leads to $f \in I$.

Hilbert's Nullstellensatz

Nonnegative Polynomials

positive semidefinite matrices

Applications

control of nonlinear systems and global optimization

Sum of Squares and Radical Ideals

real radicals and a real Nullstellensatz

Moment Matrices

counting the number of real solutions

For a polynomial system $f(\mathbf{x}) = \mathbf{0}$, with $f = (f_1, f_2, \dots, f_N)$, $f_i \in \mathbb{C}[\mathbf{x}]$, for $i = 1, 2, \dots, N$, Hilbert's Nullstellensatz implies

$$f^{-1}(\mathbf{0}) = \emptyset \quad \Leftrightarrow \quad \langle f \rangle = \langle 1 \rangle.$$

So if the system $f(\mathbf{x}) = \mathbf{0}$ has no solutions, then there are polynomials $g_i \in \mathbb{C}[\mathbf{x}]$, $i = 1, 2, \dots, N$, such that

$$1 = g_1 f_1 + g_2 f_2 + \dots + g_N f_N.$$

The polynomials g_i provide a certificate that $f(\mathbf{x}) = \mathbf{0}$ has no solutions.

Real Nullstellensatz

Nonnegative
Polynomialspositive semidefinite
matrices

Applications

control of nonlinear
systems and global
optimizationSum of
Squares and
Radical Idealsreal radicals and a
real NullstellensatzMoment
Matricescounting the number
of real solutions

Theorem (Real Nullstellensatz)

The system of real polynomial equations and inequalities

$$\left\{ \begin{array}{l} f_1(\mathbf{x}) = 0, f_2(\mathbf{x}) = 0, \dots, f_r(\mathbf{x}) = 0 \\ g_1(\mathbf{x}) \geq 0, g_2(\mathbf{x}) \geq 0, \dots, g_s(\mathbf{x}) \geq 0 \\ h_1(\mathbf{x}) > 0, h_2(\mathbf{x}) > 0, \dots, h_t(\mathbf{x}) > 0 \end{array} \right.$$

has a solution for $\mathbf{x} \in \mathbb{R}^n$, or there exists an identity

$$\begin{aligned} & \sum_{i=1}^r \alpha_i f_i + \sum_{\nu \in \{0,1\}^s} \left(\sum_j b_{j\nu}^2 \right) \cdot g_1^{\nu_1} g_2^{\nu_2} \cdots g_s^{\nu_s} \\ & + \sum_{\nu \in \{0,1\}^t} \left(\sum_j c_{j\nu}^2 \right) \cdot h_1^{\nu_1} h_2^{\nu_2} \cdots h_t^{\nu_s} + \sum_k d_k^2 + \prod_{l=1}^t h_l^{u_l} = 0, \end{aligned}$$

where $u_j \in \mathbb{N}$ and $a_i, b_{j\nu}, c_{j\nu}$, and d_k are polynomials.

Sum of Squares

Nonnegative Polynomials

positive semidefinite matrices

Applications

control of nonlinear systems and global optimization

Sum of Squares and Radical Ideals

real radicals and a real Nullstellensatz

Moment Matrices

counting the number of real solutions

- 1 Nonnegative Polynomials
positive semidefinite matrices
- 2 Applications
control of nonlinear systems and global optimization
- 3 Sum of Squares and Radical Ideals
real radicals and a real Nullstellensatz
- 4 Moment Matrices
counting the number of real solutions

Nonnegative
Polynomialspositive semidefinite
matrices

Applications

control of nonlinear
systems and global
optimizationSum of
Squares and
Radical Idealsreal radicals and a
real NullstellensatzMoment
Matricescounting the number
of real solutions

We can represent a polynomial with support A as

$$f(\mathbf{x}) = \sum_{\mathbf{a} \in A} c_{\mathbf{a}} \mathbf{x}^{\mathbf{a}} \in \mathbb{R}[\mathbf{x}] \text{ by its coefficient vector } \mathbf{c}_A$$

with respect to some fixed monomial basis $(\mathbf{x}^{\mathbf{a}})_{\mathbf{a} \in A}$.

The corresponding dual basis uses differentials $(\partial^{\mathbf{a}})_{\mathbf{a} \in A}$.

Given a sequence $\mathbf{y} = (y_{\mathbf{a}})_{\mathbf{a} \in A} \in \mathbb{R}^m$, for $m = \#A$, we consider a linear functional $\Lambda \in (\mathbb{R}[\mathbf{x}])^*$ as $\Lambda = \sum_{\mathbf{a} \in A} y_{\mathbf{a}} \partial^{\mathbf{a}}$.

Then \mathbf{y} is *the coordinate sequence* $\mathbf{y} = (\Lambda(\mathbf{x}^{\mathbf{a}}))_{\mathbf{a} \in A}$.

Given the linear form Λ , we define the *moment matrix* as

$$M(\mathbf{y}) = (\Lambda(\mathbf{x}^{\mathbf{a}+\mathbf{b}}))_{\mathbf{a}, \mathbf{b}} = (y_{\mathbf{a}+\mathbf{b}})_{\mathbf{a}, \mathbf{b}},$$

with rows and columns indexed by $\mathbf{a}, \mathbf{b} \in A$.

Nonnegative
Polynomialspositive semidefinite
matrices

Applications

control of nonlinear
systems and global
optimizationSum of
Squares and
Radical Idealsreal radicals and a
real NullstellensatzMoment
Matricescounting the number
of real solutions

For a polynomial f as denoted above, we have the inner product

$$\langle f, f \rangle_{\Lambda} = \mathbf{c}_A^T M(\mathbf{y}) \mathbf{c}_A = \Lambda(f^2).$$

By construction, the positive semidefiniteness of $M(\mathbf{y})$ is equivalent to $\Lambda(f^2) \geq 0$.

An interpretation of the moment matrix is the matrix representing the quasi-Hankel operator:

$$H_{\Lambda} : \mathbb{R}[\mathbf{x}] \rightarrow (\mathbb{R}[\mathbf{x}])^* : h \mapsto h \cdot \Lambda.$$

We have $h \cdot \Lambda(f) = \Lambda(hf)$ for all $f \in \mathbb{R}[\mathbf{x}]$.

The kernel of the moment matrix is

$$\ker M(\mathbf{y}) = \{ f \in \mathbb{R}[\mathbf{x}] \mid M(\mathbf{y})\mathbf{c}_A = \mathbf{0} \}.$$

The kernel $\ker M(\mathbf{y})$ is an ideal in $\mathbb{R}[\mathbf{x}]$.

If $M(\mathbf{y})$ is positive semidefinite, then $\ker M(\mathbf{y})$ is real radical.

In computations, we consider polynomials of degree at most d , denoted by $\mathbb{R}[\mathbf{x}]_d$. Given a linear form $\Lambda = \sum_{\mathbf{a} \in A} y_{\mathbf{a}} \partial^{\mathbf{a}}$,

the matrix

$$M_{\lfloor d/2 \rfloor}(\mathbf{y}) = \Lambda(\mathbf{x}^{\mathbf{a}+\mathbf{b}})_{\mathbf{a},\mathbf{b}} = (y_{\mathbf{a}+\mathbf{b}})_{\mathbf{a},\mathbf{b}}$$

is *the truncated moment matrix*.

The method is to find \mathbf{y} for which the rank of the moment matrix is maximum.

This rank equals the number of real solutions.

The Moment-Matrix algorithm:

Input: $I = \langle f_1, f_2, \dots, f_N \rangle$, with $\# V_{\mathbb{R}}(I) < \infty$.

Output: a border or Gröbner basis for $J := \ker M_d(\mathbf{y})$,
with $V_{\mathbb{C}}(J) = V_{\mathbb{C}}(I)$.

1. $d := D := \max_{i=1}^N \deg(f_i)$;
2. Find a generic \mathbf{y} ;
3. if $(\text{rank} M_s(\mathbf{y}) = \text{rank} M_{s-1}(\mathbf{y}), \text{ for some } D \leq s \leq \lfloor d/2 \rfloor$
or $\text{rank} M_s(\mathbf{y}) = \text{rank} M_{s-t}(\mathbf{y}) \text{ for some } t \leq s \leq \lfloor d/2 \rfloor)$
4. then return a basis for the column space of $M_{d-1}(\mathbf{y})$;
5. else $d := d + 1$; go to step 2;
6. end if.

Summary + Exercises

Interior-point methods are useful for real root finding of polynomial systems.

Exercises:

- 1 Verify that we may write every fourth degree polynomial as $\mathbf{z}^T \mathbf{A} \mathbf{z}$ with $\mathbf{z} = [x^2 \ x \ 1]$, although not necessarily in a unique way. Can you generalize this representation for polynomials of any degree?
- 2 Suppose we want to compute the global minimum of a cubic polynomial f in 10 variables. Use Bézout's theorem to bound the number of solutions of the system of all partial derivatives. Compare this number with the number of monomials in f .
- 3 Install SOSTOOLS on your computer and use it for example through its Macaulay 2 interface in the paper of Peyrl and Parrilo. Report running times for the examples given by Peyrl and Parrilo.

one last exercise

Nonnegative
Polynomialspositive semidefinite
matrices

Applications

control of nonlinear
systems and global
optimizationSum of
Squares and
Radical Idealsreal radicals and a
real NullstellensatzMoment
Matricescounting the number
of real solutions

- 4 Consider the family of homogeneous polynomials

$$M_{jk}(x_1, x_2, x_3) = jx_3^6 + x_1^2 x_2^2 (jx_1^2 + jx_2^2 - kx_3^2),$$

where j and k are positive integers.

Give examples of numerical evaluations of M_{jk} for some instances of j and k at well chosen values for x_1 , x_2 , and x_3 to illustrate the importance of positivity on the accuracy of the evaluation.