

A NOTE ON SPACES OF ASYMPTOTIC DIMENSION ONE

KOJI FUJIWARA AND KEVIN WHYTE

ABSTRACT. Let X be a geodesic metric space with $H_1(X)$ uniformly generated. If X has asymptotic dimension one then X is quasi-isometric to an unbounded tree. As a corollary, we show that the asymptotic dimension of the curve graph of a compact, oriented surface with genus $g \geq 2$ and one boundary component is at least two.

INTRODUCTION

Our main results show that, under fairly weak restrictions, a geodesic metric space of asymptotic dimension one is quasi-isometric to a tree. Before stating precise results we review some definitions.

If X is a set and $X = \cup_i O_i$ a covering, we say that the *multiplicity* of the covering is at most n if any point $x \in X$ is contained in at most n elements of $\{O_i\}_i$.

Recall that the *covering dimension* of a topological space X is the minimal d such that every open covering has a refinement with multiplicity at most $d + 1$. The notion of asymptotic dimension is a coarse analogue introduced by Gromov in [Gr93]:

Let X be a metric space, and $X = \cup_i O_i$ a covering. For $D \geq 0$, we say that the *D -multiplicity* of the covering is at most n if for any $x \in X$, the closed D -ball centered at x intersects at most n elements of $\{O_i\}_i$. The multiplicity is exactly the 0-multiplicity.

The *asymptotic dimension* of the metric space X is at most n if for any $D \geq 0$, there exists a covering $X = \cup_i O_i$ such that the diameter of O_i is uniformly bounded (i.e. there exists C such that for all i , $\text{diam} O_i \leq C$), and the D -multiplicity of the covering is at most $n + 1$. We say that the asymptotic dimension of X , *asdim* X , is n if the asymptotic dimension of X is at most n , but it is not at most $n - 1$. If such n does not exist, then we define the asymptotic dimension of X to be infinite.

It is easy to see that if two metric spaces are quasi-isometric, then they have the same asymptotic dimension. A geodesic metric space has asymptotic dimension zero if and only if it is bounded. Since unbounded trees

Date: 2006.10.11.

This work has been done when the first author was visiting the second author at University of Illinois at Chicago, and during his visit at Max-Planck-Institute for Mathematics in Bonn. He would like to thank both institutions for hospitality.

The second author was partially supported by an NSF Career award (DMS-0204576).

have asymptotic dimension one, all spaces quasi-isometric to unbounded trees also have asymptotic dimension one, [Gr93].

Let X be a geodesic metric space. We say $H_1(X)$ is *uniformly generated* if there is an $L > 0$ so that $H_1(X)$ is generated by loops of length at most L .

Theorem 0.1. *Let X be a geodesic metric space with $H_1(X)$ uniformly generated. If X has asymptotic dimension one then X is quasi-isometric to an unbounded tree.*

Remark: This theorem relies on a result of Manning ([Man]) characterizing spaces quasi-isometric to trees, see Theorem 1.2. The restriction on $H_1(X)$ is essential, see example 1.4.

Let G be a finitely generated group, and Γ the Cayley graph with respect to a finite generating set. Since the asymptotic dimension is invariant by quasi-isometry, the asymptotic dimension of G is defined to be the asymptotic dimension of Γ .

Corollary 0.2. *Let G be a finitely presented group. The asymptotic dimension of G is one if and only if G contains a non-trivial free group as a subgroup of finite index.*

We are informed by Dranishnikov that this has been known [JS] (cf. [Ge], [D]).

Corollary 0.3. *Let X be geodesic metric space which is Gromov hyperbolic. If X has asymptotic dimension one then X is quasi-isometric to an unbounded tree.*

This result does not require that X be proper. One important class of examples which are not proper are curve complexes. Let $S = S_{g,p}$ be a compact, orientable surface such that g is the genus and p is the number of the connected components of the boundary of S . We assume that $3g - 4 + p > 0$. The *curve complex* of S , defined by Harvey [Ha], are the flag complexes with 1-skeleton the *curve graph* of S , $C(S)$. The curve graph is the graph whose vertices are isotopy classes of essential, nonperipheral, simple closed curves in S , with two distinct vertices joined by an edge if the corresponding curves can be realized by disjoint curves. We remark that the curve complex of S is quasi-isometric to the curve graph of S , so that they have same asymptotic dimension.

Masur and Minsky [MaMi] show the remarkable result that $C(S)$ is Gromov hyperbolic.

It is known that $C(S)$ has finite asymptotic dimension ([BeF]). Since $C(S)$ is unbounded it cannot have asymptotic dimension zero. No upper bound or non-trivial lower bound is known in general. In the case $p = 1$, Schleimer shows in [Sch] that $C(S)$ is one-ended, from which it follows that it is not quasi-isometric to a tree. Thus we can improve the lower bound on asymptotic dimension here:

Corollary 0.4. *Let S be a compact, oriented surface with genus $g \geq 2$ and one boundary component. Let $C(S)$ be the curve graph of S . Then the asymptotic dimension of $C(S)$ is at least two.*

Acknowledgements. The first author would like to thank Greg Bell for discussion. We appreciate comments and information by Dranishnikov.

1. PROOFS

Theorem 0.1 follows from two key ingredients. The first is the classical argument that surfaces have covering dimension two (see, for instance, [Mu], section 55). The fact we need is the following:

Lemma 1.1. *Let S be a compact surface with boundary S^1 . Let A , B , and C be points on ∂S , dividing it into arcs AB , AC , and BC . If $\{O_i\}_i$ is an open cover of S with multiplicity two then there is some i for which O_i intersects all three segments AB , AC , and BC .*

The second ingredient is a characterization of spaces quasi-isometric to trees due to Manning:

Theorem 1.2 (Th 4.6 [Man]). *Let Y be a geodesic space. Suppose that there exists a constant K with the following property:*

let a, b be points in Y , γ a geodesic from a to b , and α a path from a to b . Then the K -ball at the midpoint m of the geodesic γ has non-empty intersection with α .

Then Y is quasi-isometric to a simplicial tree.

We now prove Theorem 0.1:

Proof. The following lemma, essentially contained in [BW], explains the role of the uniform generation assumption:

Lemma 1.3. *Let X be a complete geodesic metric space. The following are equivalent:*

- X has uniformly generated H_1 .
- X is quasi-isometric to a complete geodesic metric space, Y , with $H_1(Y) = 0$.

(These are also equivalent to the condition $H_1^{uf}(X) = 0$ which is the form which appears in [BW].)

Proof. Let Y be a complete geodesic metric space with $H_1(Y) = 0$, and let $f : X \rightarrow Y$ be a (K, C) -quasi-isometry. Suppose l is any loop in X . Divide l into segments $[a_0, a_1], [a_1, a_2], \dots, [a_{n-1}, a_n]$, where $a_0 = a_n$ and the length of each segment is at most 1.

Let $b_i = f(a_i)$. We have $d(b_i, b_{i+1}) \leq K + C$. For each i choose a geodesic segment in Y connecting b_i to b_{i+1} . Call the resulting loop l' . By

assumption, l' bounds, so we have $l' = \partial S$ for some surface S mapping to Y . Triangulate S so that the image of each simplex has diameter at most 1 in Y . For each vertex v in S let $\alpha(v)$ be a point of X which maps close to the image of v in Y . For each edge e of S with endpoints u and v , choose a geodesic segment, $\alpha(e)$, in X connecting $\alpha(u)$ to $\alpha(v)$.

Thus we have α mapping the one-skeleton of S to X . For each simplex σ of S we have a loop in X of length at most D (depending only on K and C). The boundary component of S is mapped to a loop l'' which is within Hausdorff distance D of l . The map α exhibits that l'' is a sum of classes of length at most D , and the condition on Hausdorff distance implies the same for $l - l''$. Thus l is a sum of loops of length at most D , proving that $H_1(X)$ is uniformly generated.

Conversely, suppose $H_1(X)$ is generated by loops of length at most D . Let A be a maximal collection of points in X with $d(a, a') \geq D$ for all $a \neq a'$. Let $R = 3D$, and let Y be the space:

$$X \cup_{a \in A} \text{cone}(B(a, R))$$

In words, Y is X with each R -ball centered at $a \in A$ coned to a point. We give Y the induced path metric where each cone line has length R . The inclusion of X is then isometric, and has coarsely dense image, and so is a quasi-isometry. By construction, any loop l in X of length at most D is contained $B(a, R)$ for some a , and hence is null homotopic in Y . Since these generate $H_1(X)$, we have $H_1(X) \rightarrow H_1(Y)$ is the zero map.

Let l be any loop in Y . By compactness of the circle, l only passes through finitely many cone points. By pushing l off of the cones whose cone points it misses, we may arrange for l to visit the interior of only finitely many cones. For each such cone $\text{cone}(B(a, R))$, let u and v be the first and last places l visits this cone. By a further homotopy we may assume l travels along the cone line from u to the cone point and then along the cone line back from the cone point to v . Choose a path p_a of length at most $2R$ in $B(a, R)$ connecting u and v . Let l' be the loop with all the trips into the cones cut out and replaced with the p_a . The disks $c(p_a)$ give a homotopy between l and l' in Y . Since $l' \subset X$ we know $[l] = [l'] = 0$ in Y . Thus $H_1(Y) = 0$ as claimed. \square

Remark: The above proof goes through verbatim to show that uniform generation of $\pi_1(X)$ is equivalent to X quasi-isometric to a simply connected space.

Returning to the proof of Theorem 0.1, we assume X is quasi-isometric to a geodesic space Y with $H_1(Y) = 0$ using the lemma. Since Y has asymptotic dimension one there is a cover O_i of Y with 1-multiplicity two, and all O_i of diameter at most some $D < \infty$. Let U_i be the open subset of Y consisting of the points within a distance of 1 of O_i . By definition this cover has multiplicity at most two, and each U_i has diameter at most $D + 2$.

We check that Manning's condition holds in Y with $K = \frac{3}{2}(D + 2)$. Let γ , a , b , m , and α be as in the statement of theorem 1.2. Let l be the loop $\gamma \circ \alpha^{-1}$. Since $H_1(Y) = 0$ there is a surface S with $\partial S = l$ and a continuous map $f : S \rightarrow Y$ with $f|_{\partial S} = l$. Applying lemma 1.1 to the covering $f^{-1}(U_i)$ shows that there is some i such that U_i has non-trivial intersection with α and with the segments \overline{am} and \overline{mb} of γ . Since U_i has diameter at most $D + 2$ this means there are points p on α and s and t on \overline{am} and \overline{mb} which are pairwise within a distance $D + 2$. Since γ is a geodesic, one of s or t must be within $\frac{1}{2}(D + 2)$ of m , so α must pass through the K ball around m as desired. We showed that Y , therefore X as well, is quasi-isometric to a tree, which is unbounded. \square

Remark: The proof only uses that X has a cover by sets of bounded diameter with L_0 -multiplicity two for L_0 large enough to generate all of $H_1(X)$. Since it follows that X is quasi-isometric to a tree, we know that there are covers by uniformly bounded sets of L -multiplicity two for all L .

Theorem 0.1 does not hold without the assumption on $H_1(X)$:

Example 1.4. *Let X be a graph with vertex set \mathbb{N} and two edges connecting n and $n + 1$, both of length 2^n . Thus X is a string of circles of increasing radii, each touching the next at a single point. It is clear that $H_1(X)$ is not uniformly generated and in particular that X is not quasi-isometric to a tree.*

For any $D > 0$, choose n such that $2^n \gg D$. Let U be the union of the vertices $\{1, 2, \dots, n\}$ of X and all adjacent edges except for the $\frac{1}{2}D$ -ball around the vertex $n + 1$. For $i > 0$ let A_i be the ball of radius D around the vertex $n + i$, let B_i and C_i be the subsets of the two edges from $n + i$ to $n + i + 1$ not within $\frac{1}{2}D$ of a vertex. It is easy to see that collection $\{U, A_i, B_i, C_i\}$ is a covering of X with D -multiplicity two. By subdividing the B_i and C_i to a collection of intervals of length $2D$ covering each, one gets a covering with D -multiplicity by sets of diameter at most 2^n . Thus X has asymptotic dimension one.

Proof of Cor 0.2. A well known corollary of work of Stallings ([Sta68]) and Dunwoody ([Dun85]) is that a group quasi-isometric to a tree is virtually free. Thus corollary 0.2 follows immediately from Theorem 0.1 since the fundamental group of the Cayley graph is generated by translates of the relators, and so is generated by loops of length at most the length of the longest relator. \square

Remark: In fact, we get something a priori stronger: if $\Gamma = \langle F | R \rangle$ where R_{ab} is a finitely generated Γ -module and Γ has asymptotic dimension one then Γ is virtually free. However, since every virtually free group is finitely presented, this does not include any new examples.

Proof of Cor 0.3. To prove Corollary 0.3, we need to check that $H_1(X)$ is uniformly generated. For any graph G we have that $H_1(G)$, and, indeed, $\pi_1(G)$, are generated by isometrically embedded loops. If we divide such a loop into three equal pieces we get a geodesic triangle whose sides come close only at the corners. If G is δ -hyperbolic, this implies any such loop can have length at most 6δ , which proves the corollary. \square

REFERENCES

- [BeF] G. Bell, K. Fujiwara. The asymptotic dimension of a curve graph is finite, preprint, 2005 September. Arxiv, math.GT/0509216.
- [BW] J. Block and S. Weinberger, *Large Scale Homology Theories and Geometry*, AMS/IP Studies in Advanced Mathematics, volume 2, 1997, pp. 522-569.
- [D] A. N. Dranishnikov. Cohomological approach to asymptotic dimension. preprint, 2006 August, math.MG/0608215.
- [Dun85] M. J. Dunwoody, *The accessibility of finitely presented groups*, Invent. Math. **81** (1985), 449–457.
- [Ge] A. Gentimis. Asymptotic dimension of finitely presented group, preprint 2005.
- [Gr87] M. Gromov. Hyperbolic groups. In *Essays in group theory*, 75–263. Springer, New York, 1987.
- [Gr93] M. Gromov, *Asymptotic invariants of infinite groups*, Geometric Group Theory, London Math. Soc. Lecture Note Ser. (G. Niblo and M. Roller, eds.), no. 182, 1993.
- [Ha] W. J. Harvey. Boundary structure of the modular group. In *Riemann surfaces and related topics: Proceedings of the 1978 Stony Brook Conference*, 245–251, 1981. Princeton Univ. Press.
- [JS] Tadeusz Januszkiewicz, Jacek Swiatkowski. Filling invariants in systolic complexes and groups. preprint 2005.
- [Man] J. Manning, Fox Geometry of pseudocharacters. *Geom. Topol.* **9** (2005), 1147–1185.
- [MaMi] H. Masur, Y.N. Minsky. Geometry of the complex of curves. I. Hyperbolicity. *Invent. Math.* **138** (1999), no. 1, 103–149.
- [Mu] J.R. Munkres, "Topology", 2nd Ed. Prentice Hall 2000,
- [Sch] S. Schleimer. Notes on the complex of curves. preprint, 2005.
- [Sta68] J. Stallings, *On torsion free groups with infinitely many ends*, *Ann. of Math.* **88** (1968), 312–334.

MATH INSTITUTE, TOHOKU UNIVERSITY, SENDAI, 980-8578 JAPAN
E-mail address: fujiwara@math.tohoku.ac.jp

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT CHICAGO, CHICAGO, IL
E-mail address: kwhyte@math.uic.edu