

MCS 590 – Mathematical Foundations of Data Science
Fall 2017
Problem Set 1

Lev Reyzin

Due: 9/29/17 at the beginning of class

Instructions: Atop your problem set, please write your name and list your collaborators.

Problems

Prove all your answers.

1. For what value of d is the volume of the d -dimensional unit ball maximized?
2. Suppose we are given n unit vectors in R^n divided into two sets P, Q with the guarantee that there exists a hyperplane $a \cdot x = 0$ such that every point in P is on one side of it and every point in Q is on the other. Furthermore, assume that the ℓ_2 distance of each point to the hyperplane is at least γ (this is sometimes called the “margin”). Show that random projection (as defined in the book) to $O(\log n/\gamma^2)$ dimensions will have the property that with high probability, the two sets of points will still remain separated by a hyperplane with margin $\gamma/2$.
3. Show the following:
 - i. If A is a symmetric matrix with distinct singular values, then the left and right singular vectors are the same and $A = VDV^T$.
 - ii. If A is a real valued matrix, $B = AA^T$ is positive semi-definite.
4. A Markov chain is said to be symmetric if for all i and j , $p_{ij} = p_{ji}$. What is the stationary distribution of a connected symmetric Markov chain?
5. Given a Markov chain on an undirected graph, we modify the chain as follows: at the current state, we stay there with probability $1/2$; with the other probability $1/2$, we move as in the old chain. Show that the new chain has the same stationary distribution.