

On the Chromatic Thresholds of Hypergraphs

József Balogh ^{*} Jane Butterfield [†] Ping Hu [‡] John Lenz [§]
Dhruv Mubayi [¶]

March 8, 2011

Abstract

Let \mathcal{F} be a family of r -uniform hypergraphs. The *chromatic threshold* of \mathcal{F} is the infimum of all non-negative reals c such that the subfamily of \mathcal{F} comprising hypergraphs H with minimum degree at least $c \binom{|V(H)|}{r-1}$ has bounded chromatic number. This parameter has a long history for graphs ($r = 2$), and in this paper we begin its systematic study for hypergraphs.

Łuczak and Thomassé recently proved that the chromatic threshold of near bipartite graphs is zero, and our main contribution is to generalize this result to r -uniform hypergraphs. For this class of hypergraphs, we also show that the exact Turán number is achieved uniquely by the complete $(r + 1)$ -partite hypergraph with nearly equal part sizes. This is one of very few infinite families of nondegenerate hypergraphs whose Turán number is determined exactly. In an attempt to generalize Thomassen's result that the chromatic threshold of triangle-free graphs is $1/3$, we prove bounds for the chromatic threshold of the family of 3-uniform hypergraphs not containing $\{abc, abd, cde\}$, the so-called generalized triangle.

In order to prove upper bounds we introduce the concept of *fiber bundles*, which can be thought of as a hypergraph analogue of directed graphs. This leads to the notion of *fiber bundle dimension*, a structural property of fiber bundles which is based on the idea of Vapnik-Chervonenkis dimension in hypergraphs. Our lower bounds follow from explicit constructions, many of which use a generalized Kneser hypergraph. Using methods from extremal set theory, we prove that these generalized Kneser hypergraphs have unbounded chromatic number. This generalizes a result of Szemerédi for graphs and might be of independent interest. Many open problems remain.

Keywords: hypergraphs, chromatic threshold, exact Turán number, VC-dimension

^{*}University of Illinois, Urbana-Champaign, University of California, San Diego, jobal@math.uiuc.edu. This material is based upon work supported by NSF CAREER Grant DMS-0745185, UIUC Campus Research Board Grant 11067, and OTKA Grant K76099.

[†]University of Illinois, Urbana-Champaign, jbutter2@math.uiuc.edu. Research supported in part by the Dr. Lois M. Lackner Mathematics Fellowship.

[‡]University of Illinois, Urbana-Champaign, pinghu1@math.uiuc.edu.

[§]University of Illinois, Urbana-Champaign, jlenz2@math.uiuc.edu.

[¶]Department of Mathematics, Statistics, and Computer Science, University of Illinois, Chicago IL 60607, email: mubayi@math.uic.edu. Research supported in part by NSF Grant 0969092.

1 Introduction

An r -uniform hypergraph on n vertices is a collection of r -subsets of V , where V is a set of n elements. If $r = 2$ then we call it a graph. The r -sets in a hypergraph are called **edges**, and the n elements of V are called **vertices**. For a hypergraph H let $V(H)$ denote the set of vertices. We denote the set of edges by either $E(H)$ or simply H . The **chromatic number** of a hypergraph H , denoted $\chi(H)$, is the least integer k for which there exists a map $f : V(H) \rightarrow [k]$ such that if E is an edge in the hypergraph then there exist $v, u \in E$ for which $f(v) \neq f(u)$. For a vertex v in a hypergraph H we let $d(v)$ denote the number of edges in H that contain v . We let $\delta(H) = \min\{d(v) : v \in V(H)\}$, called the **minimum degree** of H .

Definition. Let \mathcal{F} be a family of r -uniform hypergraphs. The **chromatic threshold** of \mathcal{F} , is the infimum of the values $c \geq 0$ such that the subfamily of \mathcal{F} consisting of hypergraphs H with minimum degree at least $c \binom{|V(H)|}{r-1}$ has bounded chromatic number.

We say that F is a subhypergraph of H if there is an injection from $V(F)$ to $V(H)$ such that every edge in F gets mapped to an edge of H . Notice that this is only possible if both H and F are r -uniform for some r . If H is an r -uniform hypergraph, then the family of H -free hypergraphs is the family of r -uniform hypergraphs that do not contain H as a (not necessarily induced) subgraph.

The study of the chromatic thresholds of graphs was motivated by a question of Erdős and Simonovits [6]: “If G is non-bipartite, what bound on $\delta(G)$ forces G to contain a triangle?” This question was answered by Andrásfai, Erdős, and Sós [3], who showed that the answer is $2/5 |V(G)|$, achieved by the blowup of C_5 . Andrásfai, Erdős, and Sós’s [3] result can be generalized to construct triangle-free graphs with chromatic number at least k and large minimum degree. As k increases, these constructions have minimum degree approaching $1/3$. This led to the following conjecture: if $\delta(G) > (1/3 + \epsilon) |V(G)|$ and G is triangle-free, then $\chi(G) < k_\epsilon$, where k_ϵ is a constant depending only on ϵ .

Note that the conjecture is equivalent to the statement that the family of triangle-free graphs has chromatic threshold $1/3$. The conjecture was proven by Thomassen [35]. Subsequently, there have been three more proofs of the conjecture: one by Łuczak [22] using the Regularity Lemma, a result of Brandt and Thomassé [4] proving that one can take $k_\epsilon = 4$, and a recent proof by Łuczak and Thomassé [23] using the concept of Vapnik-Chervonenkis dimension (which is defined later in this paper).

For other graphs, Goddard and Lyle [14] proved that the chromatic threshold of the family of K_r -free graphs is $(2r - 5)/(2r - 3)$ while Thomassen [36] showed that the chromatic threshold of the family of C_{2k+1} -free graphs is zero for $k \geq 2$. Recently, Łuczak and Thomassé [23] gave another proof that the class of C_{2k+1} -free graphs has chromatic threshold zero for $k \geq 2$, as well as several other results about related families, such as Petersen-free graphs. The main result of Allen, Böttcher, Griffiths, Kohayakawa and Morris [1] is to determine the chromatic threshold of the family of H -free graphs for all H .

We finish this section with some definitions. For an r -uniform hypergraph H and a set of vertices $S \subseteq V(H)$, let $H[S]$ denote the r -uniform hypergraph consisting of exactly those edges of H that are completely contained in S . We call this the hypergraph **induced by** S . A set of vertices $S \subseteq V(H)$ is called **independent** if $H[S]$ contains no edges and

strongly independent if there is no edge of H containing at least two vertices of S . A hypergraph is s -partite if its vertex set can be partitioned into s parts, each of which is strongly independent.

If \mathcal{H} is a family of r -uniform hypergraphs, then the family of \mathcal{H} -free hypergraphs is the family of r -uniform hypergraphs that contain no member of \mathcal{H} as a (not necessarily induced) subgraph. For an r -uniform hypergraph H and an integer n , let $ex(n, H)$ be the maximum number of edges an r -uniform hypergraph on n vertices can have while being H -free and let

$$\pi(H) = \lim_{n \rightarrow \infty} \frac{ex(n, H)}{\binom{n}{r}}.$$

We call $\pi(H)$ the **Turán density** of H .

Let $T_{r,s}(n)$ be the complete n -vertex, r -uniform, s -partite hypergraph with part sizes as equal as possible. When $s = r$, we write $T_r(n)$ for $T_{r,r}(n)$. Let $t_r(n)$ be the number of edges in $T_r(n)$. We say that an r -uniform hypergraph H is **stable** with respect to $T_r(n)$ if $\pi(H) = r!/r^r$ and for any $\epsilon > 0$ there exists $\delta > 0$ such that if G is an H -free r -uniform hypergraph with at least $(1 - \delta)t_r(n)$ edges, then there is a partition of $V(G)$ into U_1, U_2, \dots, U_r such that all but at most ϵn^r edges of G have exactly one vertex in each part.

Let $\text{TK}^r(s)$ be the r -uniform hypergraph obtained from the complete graph K_s by enlarging each edge with $r - 2$ new vertices. The **core vertices** of $\text{TK}^r(s)$ are the s vertices of degree larger than one. For $s > r$, let $\mathcal{TK}^r(s)$ be the family of r -uniform hypergraphs such that there exists a set S of s vertices where each pair of vertices from S are contained together in some edge. The set S is called the set of **core vertices** of the hypergraph. For $s \leq r$, let $\mathcal{TK}^r(s)$ be the family of r -uniform hypergraphs such that there exists a set S of s vertices where for each pair of vertices $x \neq y \in S$, there exists an edge E with $E \cap S = \{x, y\}$ (the definition is different when $s \leq r$ so that a hypergraph which is just a single edge is not in $\mathcal{TK}^r(s)$). It is obvious that $\text{TK}^r(s) \in \mathcal{TK}^r(s)$.

2 Results

Motivated by the above results, we investigate the chromatic thresholds of the families of A -free hypergraphs for some r -uniform hypergraphs A . One of our main results concerns a generalization of cycles to hypergraphs.

Definition. Let H be an r -uniform hypergraph. We say that H is **near r -partite** if there exists a partition $V_1 \cup \dots \cup V_r$ of $V(H)$ such that all edges of H either cross the partition (have one vertex in each V_i) or are contained entirely in V_1 , and in addition $H[V_1]$ is a partial matching. The edges in $H[V_1]$ are called the **special edges**. Say that H is **mono near r -partite** if $H[V_1]$ contains exactly one edge.

Our main theorem claims that for an infinite family of hypergraphs H the chromatic threshold of the family of H -free hypergraphs is 0. This is the first (non-trivial) family of hypergraphs whose chromatic threshold is determined.

Theorem 1. *Let H be an r -uniform, near r -partite hypergraph. If H is $\mathcal{TK}^r(3)$ -free, then the chromatic threshold of the family of H -free hypergraphs is zero.*

The proof of Theorem 1 requires a slightly weaker condition on H , so we actually prove a statement slightly stronger than Theorem 1. The proof does not require that H is $\mathcal{TK}^r(3)$ -free, just that any copy of a hypergraph in $\mathcal{TK}^r(3)$ in H has at most one core vertex in a special edge of H .

For a subfamily of the hypergraphs considered in Theorem 1 we determine the exact extremal hypergraph. For many hypergraphs H (for example the Fano plane), at first only asymptotic extremal results were proved and later the precise structure of extremal hypergraphs was determined. We prove that if a mono near r -partite hypergraph H has Turán density $r!/r^r$ and is stable with respect to $T_r(n)$, then its unique extremal hypergraph is the complete r -partite hypergraph. Similar phenomena occur for graphs; see Simonovits [33], where for critical graphs the Erdős-Stone Theorem [8] was sharpened.

Definition. Let H be an r -uniform hypergraph. We say that H is **critical** if

- H is mono near r -partite,
- the special edge of H has at least $r - 2$ vertices of degree one,
- $\pi(H) = r!/r^r$,
- H is stable with respect to $T_r(n)$.

Theorem 2. *Let H be an r -uniform critical hypergraph. Then there exists some n_0 such that for $n > n_0$, $T_r(n)$ is the unique H -free hypergraph with the most edges.*

A particularly interesting critical family is one that generalizes cycles to hypergraphs.

Definition. Let C_m^r be the r -uniform hypergraph with m edges on n vertices v_1, \dots, v_n for which

1. the n vertices are arranged consecutively in a circle,
2. each edge contains r consecutive vertices,
3. if $m = 2k + 1$ for some integer $k > 0$ then $n = rk + (r - 1)$, and if $m = 2k$ then $n = rk$,
4. edges E_i and E_j share vertices if and only if $i \in \{j - 1, j + 1\}$ or $i = 1$ and $j = m$,
5. for $i \leq m - 1$, if i is odd then $|E_i \cap E_{i+1}| = 1$; if i is even then $|E_i \cap E_{i+1}| = r - 1$, and
6. if m is even then $|E_1 \cap E_m| = 1$; if m is odd then $|E_1 \cap E_m| = r - 1$.

We say that C_m^r is **odd** if m is odd, and **even** otherwise.

Lemma 3. *If m is odd then C_m^r is not r -partite.*

Proof. Suppose $m = 2k + 1$ for some integer K . Notice that edge E_{2k+1} consists of the vertices $v_{rk+1}, v_{rk+2}, \dots, v_{rk+r-1}, v_1$. Suppose $f : V \rightarrow [r]$ is an r -coloring of the vertices of C_{2k+1}^r such that each color class induces a strongly independent set. Then vertices $v_1, v_{r+1}, v_{2k+1}, \dots, v_{rk+1}$ must all have the same color. In particular, $f(v_1) = f(v_{rk+1})$, which is a contradiction because v_1 and v_{rk+1} are both in E_{2k+1} . \square

It is easy to see that C_{2k+1}^r is mono near r -partite. A theorem of Keevash and the last author [18], combined with a theorem of Pikhurko [28], the supersaturation result of Erdős and Simonovits [7], and the hypergraph removal lemma of Gowers, Nagle, Rödl, and Skokan [15, 26, 29, 30, 34] prove that C_{2k+1}^3 and C_{2k+1}^4 are critical. It remains an open question whether C_{2k+1}^r is critical for $r \geq 5$.




Figure 1: Hypergraph Cycles

Theorem 4. *The cycles C_{2k+1}^3 and C_{2k+1}^4 are critical for every $k \geq 2$.*

Theorems 1, 2, and 4 together with the simple observation that C_{2k+1}^r is near r -partite and $\mathcal{TK}^r(3)$ -free for all $r, k \geq 2$ proves the following corollary, which extends the results in [36] and [23] that the chromatic threshold of the family of C_{2k+1} -free graphs is zero.

Corollary 5. *For $r = 3$ or $r = 4$, there exists some n_0 such that for $n > n_0$, the unique n -vertex, r -uniform, C_{2k+1}^r -free hypergraph with the largest number of edges is $T_r(n)$. For all $r, k \geq 2$, the chromatic threshold of the family of C_{2k+1}^r -free hypergraphs is zero.*

Note that Łuczak and Thomassé [23] proved Theorem 1 for graphs, and they conjectured that the family of H -free graphs has chromatic threshold zero if and only if H is near acyclic and triangle free. (A graph G is **near acyclic** if there exists an independent set S in G such that $G - S$ is a forest and every odd cycle has at least two vertices in S .) This conjecture is announced to have been verified in [1]. We pose a similar question for hypergraphs.

Problem 6. *Characterize the r -uniform hypergraphs H for which the chromatic threshold of the family of H -free hypergraphs has chromatic threshold zero.*

Another way to generalize the triangle to 3-uniform hypergraphs is the hypergraph F_5 , which is the hypergraph with vertex set $\{a, b, c, d, e\}$ and edges $\{a, b, c\}$, $\{a, b, d\}$, and $\{c, d, e\}$. Frankl and Füredi [9] proved that $ex(n, F_5)$ is achieved by $T_3(n)$ for $n > 3000$ (recently Goldwasser has determined $ex(n, F_5)$ for all n). We prove the following bounds on the chromatic threshold of the family of F_5 -free 3-uniform hypergraphs.

Theorem 7. *The chromatic threshold of the family of F_5 -free 3-uniform hypergraphs is between $6/49$ and $(\sqrt{41} - 5)/8 \approx 7/40$.*

The rest of the paper is organized as follows. First, in Section 3 we define and motivate fiber bundles and fiber bundle dimension, the main tools in the proofs of Theorem 1 and 7.

Next, in Section 4 we show the power of fiber bundle dimension by giving a relatively short proof of Theorem 1. We prove our key theorem about fiber bundle dimension, Theorem 8, in Section 5. In Section 6, we prove that C_{2k+1}^r is critical (Theorem 4), and then prove Theorem 2. The proof of Theorem 7 is given in Section 7. The final section gives lower bounds for several other families of hypergraphs, along with conjectures and open problems. The lower bounds all follow from specific constructions, some of which use a generalized Kneser hypergraph; this graph is defined and discussed in Section 8.

Throughout this paper, we occasionally omit the floor and ceiling signs for the sake of clarity.

3 Fiber Bundles and Fiber Bundle Dimension

The proofs of Theorems 1 and 7 are based on an idea pioneered by Łuczak and Thomassé [23] to color graphs, which itself was based on the Vapnik-Chervonenkis dimension. Let H be a hypergraph. A subset X of $V(H)$ is **shattered** by H if for every $Y \subseteq X$, there exists an $E \in H$ such that $E \cap X = Y$. Introduced in [32] and [37], the **Vapnik-Chervonenkis dimension** (or VC-dimension) is the maximum size of a vertex subset shattered by H .

Definition. A **fiber bundle** is a tuple (B, γ, F) such that B is a hypergraph, F is a finite set, and $\gamma : V(B) \rightarrow 2^F$. That is, γ maps vertices of B to collections of subsets of F , which we can think about as hypergraphs on vertex set F . The hypergraph B is called the **base hypergraph** of the bundle and F is the **fiber** of the bundle. For a vertex $b \in V(B)$, the hypergraph $\gamma(b)$ is called the **fiber over b** .

We should think about a fiber bundle as taking a base hypergraph and putting a hypergraph “on top” of each base vertex. There is one canonical example of a fiber bundle. Given a hypergraph B , define the **neighborhood bundle of B** to be the bundle (B, γ, F) where $F = V(B)$ and γ maps $b \in V(B)$ to $\{A \subseteq F : A \cup \{b\} \in E(B)\}$.

Why define and use the language of fiber bundles? We can consider that in some sense fiber bundles are a generalization of directed graphs to hypergraphs, where we think of $\gamma(x)$ as the “out-neighborhood” of x . In the neighborhood bundle, $\gamma(x)$ is related to the neighbors of x so we can consider the neighborhood bundle as some sort of directed analogue of the undirected hypergraph B , where each edge is directed “both ways”. By thinking of the “out-neighborhood” of x as $\gamma(x)$ and not requiring any dependency between $\gamma(x)$ and $\gamma(y)$ for $x \neq y$, we have no dependency between the neighborhood of x and the neighborhood of y , which is one of the defining differences between directed and undirected graphs. Note that the definition of a fiber bundle differs from the usual definition of *directed hypergraph* used in the literature, which is the reason we use the term “fiber bundle” instead of “directed hypergraph.”

A fiber bundle (B, γ, F) is (r_b, r_γ) -**uniform** if B is an r_b -uniform hypergraph and $\gamma(b)$ is an r_γ -uniform hypergraph for each $b \in V(B)$. Given $X \subseteq V(B)$, the **section of X** is the hypergraph with vertex set F and edges $\bigcap_{x \in X} \gamma(x)$. In other words, the section of X is the collection of subsets of F that appear in every fiber over x for $x \in X$. Motivated by a definition of Łuczak and Thomassé [23], we define the dimension of a fiber bundle. Let H be a hypergraph and define $\dim_H(B, \gamma, F)$ to be the maximum integer d such that there exist d

disjoint edges E_1, \dots, E_d of B (i.e. a matching) such that for every $x_1 \in E_1, \dots, x_d \in E_d$, the section of $\{x_1, \dots, x_d\}$ contains a copy of H . Our definition of dimension will coincide with the definition of paired VC-dimension in [23] when (B, γ, F) is $(2, 1)$ -uniform and $H = \{\{x\}\}$, the complete 1-uniform, 1-vertex hypergraph.

Let A be an r -uniform hypergraph. Our method of proving an upper bound on the chromatic threshold of the family of A -free hypergraphs, used in Theorems 1 and 7, is the following. Let G be an A -free r -uniform hypergraph with minimum degree at least $c \binom{|V(G)|}{r-1}$. We now need to show that G has bounded chromatic number, which we do in two steps. Let (G, γ, F) be the neighborhood bundle of G . First, we show that the dimension of (G, γ, F) is bounded by showing that if the dimension is large then we can find A as a subhypergraph. Given that $\dim_H(G, \gamma, F)$ is bounded, we use the following theorem to bound the chromatic number of G . In most applications, we will let H be an $(r-1)$ -uniform, $(r-1)$ -partite hypergraph.

Theorem 8. *Let $r_b \geq 2$, $r_\gamma \geq 1$, $d \in \mathbb{Z}^+$, $0 < \epsilon < 1$, and H be an r_γ -uniform hypergraph with zero Turán density. Then there exists constants $K_1 = K_1(r_b, r_\gamma, d, \epsilon, H)$ and $K_2 = K_2(r_b, r_\gamma, d, \epsilon, H)$ such that the following holds. Let (B, γ, F) be any (r_b, r_γ) -uniform fiber bundle where $\dim_H(B, \gamma, F) < d$ and for all $b \in V(B)$,*

$$|\gamma(b)| \geq \epsilon \binom{|F|}{r_\gamma}.$$

If $|F| \geq K_1$, then $\chi(B) \leq K_2$.

The above theorem is sufficient for our purposes, but our proof of Theorem 8 proves something slightly stronger. The conclusion of the above theorem can be reworded to say that either F is small, the chromatic number of B is bounded, or $\dim_H(B, \gamma, F)$ is large, which means that we can find d hyperedges E_1, \dots, E_d such that every section of $x_1 \in E_1, \dots, x_d \in E_d$ contains a copy of H . In fact, the proof shows that if F is large and the chromatic number of B is large, we can guarantee not only one copy of H but at least $\Omega(|F|^h)$ copies of H in each section, where h is the number of vertices in H .

We conjecture a similar statement for all r_γ -uniform hypergraphs H , instead of just those hypergraphs with a Turán density of zero.

Conjecture 9. *Let $r_b \geq 2$, $r_\gamma \geq 1$, $d \in \mathbb{Z}^+$, $0 < \epsilon < 1$, and H be an r_γ -uniform hypergraph. Then there exists a constants $K_1 = K_1(r_b, r_\gamma, d, \epsilon, H)$ and $K_2 = K_2(r_b, r_\gamma, d, \epsilon, H)$ such that the following holds. Let (B, γ, F) be any (r_b, r_γ) -uniform fiber bundle where $\dim_H(B, \gamma, F) < d$ and for all $b \in V(B)$,*

$$|\gamma(b)| \geq (\pi(H) + \epsilon) \binom{|F|}{r_\gamma}.$$

If $|F| \geq K_1$, then $\chi(B) \leq K_2$.

The motivation behind defining and using the language of fiber bundles rather than using the language of hypergraphs is that in the course of the proof of Theorem 8, we will modify B and γ and apply induction. As mentioned above, fiber bundles can be thought of as a directed version of a hypergraphs. When applying Theorem 8 in Sections 4 and 7, we

start with the neighborhood bundle, which carries no “extra” information beyond just the hypergraph B . But if we tried to prove Theorem 8 in the language of hypergraphs, we would run into trouble when we needed to modify γ . In the neighborhood bundle, γ is related to the neighborhood of a vertex and if we restricted ourselves to neighborhood bundles or just used the language of hypergraphs, modifying $\gamma(x)$ would imply that some $\gamma(y)$ ’s would change at the same time. The notion of a fiber bundle allows us to change the “out-neighborhood” of x independently of changing the “out-neighborhood” of $y \neq x$, and this power is critical in the proof of Theorem 8.

4 Chromatic threshold for near r -partite hypergraphs

In this section we show an application of Theorem 8 by proving Theorem 1. Fix $\epsilon > 0$ and let G be an n -vertex, r -uniform, H -free hypergraph with $\delta(G) \geq \epsilon \binom{n}{r-1}$. We would like to use Theorem 8 to bound the chromatic number, so we need to choose an appropriate bundle. We will not use the neighborhood bundle of G , but rather a closely related bundle. Once we have defined this bundle, we show it has bounded dimension by proving that if the dimension is large then we can find a copy of H in G .

As preparation, we need the following lemma, which tells us something about the structure of near r -partite graphs.

Lemma 10. *Let H be an r -uniform, near r -partite, $\mathcal{TK}^r(3)$ -free hypergraph. Let E_1, \dots, E_k be the special edges of H . For $x \in V(H)$, let $\gamma(x) = \{E - x : x \in E \in H\}$. Let N_1, \dots, N_{r^k} be the r^k possible hypergraphs $\gamma(x_1) \cap \gamma(x_2) \cap \dots \cap \gamma(x_k)$ where $x_1 \in E_1, \dots, x_k \in E_k$. Then*

- $V(N_i) \cap V(N_j) = \emptyset$ for $i \neq j$ (let $V(A) = \cup_{E \in A} E$).
- N_i is $(r-1)$ -partite for every i .

Proof. Assume $x \in V(N_i) \cap V(N_j)$ and let E_t be a special edge such that N_i selects y from E_t and N_j selects z from E_t with $y \neq z$. Then x, y, z are the core vertices of some hypergraph in $\mathcal{TK}^r(3)$, a contradiction. Secondly, N_i is $(r-1)$ -partite because $H \setminus E_1 \setminus \dots \setminus E_k$ is r -partite by the definition of near r -partite. \square

Proof of Theorem 1. Let H be an r -uniform, near r -partite, m -vertex, $\mathcal{TK}^r(3)$ -free hypergraph, and let $\epsilon > 0$ be fixed. Let G be an n -vertex, H -free hypergraph with $\delta(G) \geq \epsilon \binom{n}{r-1}$. We need to show that the chromatic number of G is bounded by a constant depending only on ϵ and H .

First, choose an equitable partition X_1, \dots, X_r of $V(G)$ such that the sizes of X_1, \dots, X_r are as equal as possible and for every $x \in V(G)$ the number of edges containing x and one vertex from each X_i is at least $\frac{1}{2r^r} \epsilon \binom{n}{r-1}$. (This can be done by randomly choosing the partition X_1, \dots, X_r .) We will show how to bound the chromatic number of $G[X_1]$; the same argument can be applied to bound the chromatic number of each $G[X_i]$ and thus the chromatic number of G .

Define the $(r, r-1)$ -uniform fiber bundle (B, γ, F) as follows. Let $B = G[X_1]$, let $F = X_2 \cup \dots \cup X_r$, and for $x \in X_1$ define

$$\gamma(x) = \{\{x_2, \dots, x_r\} \subseteq F : x_2 \in X_2, \dots, x_r \in X_r, \{x, x_2, \dots, x_r\} \in G\}.$$




Figure 2: The structure guaranteed by dimension d .

Then $\gamma(x)$ has size at least $\frac{1}{2r^r} \epsilon \binom{n}{r-1}$. Let L be the complete $(r-1)$ -uniform, $(r-1)$ -partite hypergraph on $(rm)^m$ vertices. Let V_1, \dots, V_r be the r -partition of $V(H)$ guaranteed by the definition of near r -partite and let d be the size of V_1 . Using that the Turán density of a complete $(r-1)$ -uniform $(r-1)$ -partite hypergraph is zero, we apply Theorem 8 to show that there exists constants $K_1 = K_1(r, \epsilon, H)$ and $K_2 = K_2(r, \epsilon, H)$ such that one of the following holds: either $|F| \leq K_1$, $\chi(B) \leq K_2$, or $\dim_L(B, \gamma, F) \geq d$. Since $|F| = (1 - 1/r) |V(G)|$, if either of the first two possibilities occur then the chromatic number of $G[X_1]$ is bounded. It must therefore be the case that $\dim_L(B, \gamma, F) \geq d$.

We now show this implies that G contains a copy of H , which follows from Lemma 10. Since $\dim_L(B, \gamma, F) \geq d$, there are d edges E_1, \dots, E_d such that for each $x_1 \in E_1, \dots, x_d \in E_d$, we have that $\gamma(x_1) \cap \dots \cap \gamma(x_d)$ contains a complete $(r-1)$ -uniform, $(r-1)$ -partite hypergraph on $(rm)^m$ vertices, see Figure 2. Since $m = |V(H)|$, from each $\gamma(x_1) \cap \dots \cap \gamma(x_d)$ we can pick a copy of the complete $(r-1)$ -uniform, $(r-1)$ -partite hypergraph on m vertices so that all these copies are vertex disjoint. Assume $V_1 = A_1 \cup \dots \cup A_\ell \cup \{a_{\ell+1}\} \cup \dots \cup \{a_{\ell'}\}$, where A_1, \dots, A_ℓ are the special edges of H . Using Lemma 10, we can embed a copy of H in G by mapping A_i to E_i for $1 \leq i \leq \ell$, mapping a_i to any vertex in E_i for $\ell + 1 \leq i \leq \ell'$, and mapping N_i to a subhypergraph of the corresponding complete $(r-1)$ -uniform, $(r-1)$ -partite hypergraph on m vertices. \square

5 Coloring hypergraphs with bounded dimension

5.1 An insightful attempt at proving Theorem 8

The proof of Theorem 8, which appears in Sections 5.2 and 5.3, is complex, but it started as a simple idea built on three key ideas. In this section we attempt to motivate the essential ingredients behind the proof.

Consider the following proof strategy for Theorem 8: assume the chromatic number of B is large and give an algorithm that produces d edges which witness that $\dim_H(B, \gamma, F) \geq d$. The first key idea is to use the greedy algorithm, which selects the edges one by one while maintaining that all sections are large enough to force a copy of H . Initially, we can pick

any edge E_1 , since any section of $x \in E_1$ is the set $\gamma(x)$, and by assumption $\gamma(x)$ is large enough to force a copy of H . So where could the greedy algorithm get stuck? Assume the greedy algorithm selected E_1, \dots, E_i but cannot continue. That is, for every other edge E , there exists some section S of $x_1 \in E_1, \dots, x_i \in E_i$ and there exists some $x \in E$ such that $S \cap \gamma(x)$ is too small to force a copy of H .

The second key idea in the proof is to assume that every edge of B has small overlap; that is, we assume that (B, γ, F) satisfies the condition that for every $x \neq y \in E \in B$, the number of edges in the hypergraph $\gamma(x) \cap \gamma(y)$ is small. With this assumption, the r^i sections S_1, \dots, S_{r^i} of $x_1 \in E_1, \dots, x_i \in E_i$ are almost disjoint. Let $S_{r^{i+1}} = \binom{F}{r-1} \setminus S_1 \setminus \dots \setminus S_{r^i}$ so that $S_1, \dots, S_{r^{i+1}}$ almost form a partition. Recall that the greedy algorithm could not continue because for every edge E disjoint from E_1, \dots, E_i , there existed some S_j and some $x \in E$ with $S_j \cap \gamma(x)$ small. Because $S_1, \dots, S_{r^{i+1}}$ is almost a partition, this implies that there is some other $S_{j'}$ with $S_{j'} \cap \gamma(x)$ large. In other words, the greedy algorithm cannot continue if there exists sets $S_1, \dots, S_{r^{i+1}}$ such that for every edge E , there exists some S_j and some $x \in E$ such that $S_j \cap \gamma(x)$ is large.

The third key idea is to use this in a density increment argument, similar to that used in the proof of the Regularity Lemma. If we have a partition P of (B, γ, F) (a partition will be formally defined later), we apply the greedy algorithm from the last two paragraphs in each part of P to refine the partition; the sets S_1, \dots, S_t are used to refine each part. If we define the density of a partition correctly, we can show that each time when we apply the greedy algorithm the density will increase by a constant amount and add only a constant number of new parts in the partition.

The last part of the proof is to reduce the full theorem to the case where (B, γ, F) satisfies the condition that for every $x \neq y \in E \in B$, $\gamma(x) \cap \gamma(y)$ is small.

5.2 A conditional proof of Theorem 8

Condition 1. Throughout this section, fix $r_b, r_\gamma \geq 1$, $d \in \mathbb{Z}^+$, $0 < \epsilon < \frac{1}{4}r_b^{-d}$, and H an r_γ -uniform hypergraph with zero Turán density. Also, let (B, γ, F) be an (r_b, r_γ) -uniform fiber bundle for which $\dim_H(B, \gamma, F) < d$ and if $b \in V(B)$, then $|\gamma(b)| \geq \epsilon \binom{|F|}{r_\gamma}$.

In this section, we will prove the following theorem.

Theorem 11. *Let (B, γ, F) be a fiber bundle satisfying Condition 1. Then there exists constants $0 < \lambda = \lambda(r_b, r_\gamma, d, \epsilon, H) < \epsilon$, $L_1 = L_1(r_b, r_\gamma, d, \epsilon, H)$, and $L_2 = L_2(r_b, r_\gamma, d, \epsilon, H)$ such that the following holds. Assume $|F| \geq L_1$ and for every $x \neq y \in E \in B$, $|\gamma(x) \cap \gamma(y)| \leq \lambda \binom{|F|}{r_\gamma}$. If $r_b = 1$, then $|B| \leq L_2$ and if $r_b \geq 2$ then $\chi(B) \leq L_2$.*

The only differences between this theorem and Theorem 8 are that we allow $r_b = 1$ and also assume a restriction on $\gamma(x) \cap \gamma(y)$ for $x \neq y \in E \in B$. We will remove the need for this assumption in the next section.

Define the following constants.

$$\alpha = \frac{1}{1000} \left(\frac{\epsilon}{4} \right)^{d+1}, \quad \eta = \frac{1}{4} \epsilon^2 \alpha, \quad \beta = \alpha^{1/\eta}, \quad L_2 = \lceil r_b d (r_b^d + 2)^{1/\eta} \rceil.$$

Next, pick L_1 large enough so that if $|F| \geq L_1$ and $S \subseteq \binom{F}{r_\gamma}$ with $|S| \geq 4^{-d} \epsilon^{d+1} \beta \binom{|F|}{r_\gamma}$, then S contains a copy of H . Lastly, pick $\lambda = \frac{\epsilon^{d+1} \beta}{d r_b^2 4^d}$.

Condition 2. If (B, γ, F) is a fiber bundle, then for every edge E in B and every $x \neq y \in E$, the section of x, y has at most $\lambda_{(r_\gamma)}^{(|F|)}$ edges.

If (B, γ, F) is a fiber bundle, a **partition** P of (B, γ, F) is a family $P = \{(X_1, S_1), \dots, (X_p, S_p)\}$ such that X_1, \dots, X_p is a partition of $V(B)$ and S_1, \dots, S_p is a partition of $\binom{F}{r_\gamma}$, where we allow $X_i = \emptyset$ or $S_i = \emptyset$. A partition Q is a refinement of a partition P if for each $(X, S) \in P$, there exist $(Y_1, T_1), \dots, (Y_q, T_q) \in Q$ such that $X = \cup Y_i$ and $S = \cup T_i$. For $X \subseteq V(B)$ and $S \subseteq 2^F$, the **density of** (X, S) is

$$d(X, S) = \begin{cases} 1 & S = \emptyset \text{ or } X = \emptyset, \\ \min \left\{ \frac{|\gamma(x) \cap S|}{|S|} : x \in X \right\} & \text{otherwise,} \end{cases}$$

and define

$$d(P) = \min \{d(X, S) : (X, S) \in P\}.$$

A partition P is a **partial coloring** if for every $(X, \emptyset) \in P$, we have that $B[X]$ independent. The **rank** of a partition P is the minimum of $|S|$ over all $(X, S) \in P$ with $S \neq \emptyset$.

Recall from the sketch in Section 5.1 that the general structure of our proof is to show how to refine a partial coloring P to a partial coloring Q where $d(Q) \geq \eta + d(P)$. This will imply that we can only refine the partition a constant number of times, at which point we will have a coloring with a bounded number of parts. The key lemma to facilitate this refinement is the following.

Lemma 12. *Let (B, γ, F) be a fiber bundle satisfying Conditions 1 and 2. Let $X \subseteq V(B)$ and $S \subseteq \binom{F}{r_\gamma}$ with $X \neq \emptyset$, $d(X, S) \geq \epsilon$, $|F| \geq L_1$, and $|S| \geq \beta_{(r_\gamma)}^{(|F|)}$. Then there exists a partition Y_1, \dots, Y_n, Z of X and a partition T_1, \dots, T_n of S such that $n \leq r_b^d + 1$ and*

- $|T_i| \geq \alpha |S|$,
- $d(Y_i, T_i) \geq \min \{1, \eta + d(X, S)\}$,
- $B[Z]$ is independent.

This lemma has an easy corollary.

Corollary 13. *Let (B, γ, F) be a fiber bundle satisfying Conditions 1 and 2 and $|F| \geq L_1$. Let P be a partial coloring of (B, γ, F) where P has rank at least $\alpha^k_{(r_\gamma)}^{(|F|)}$ with $k \leq \frac{1}{\eta}$. Then there exists a refinement Q of P such that*

- $|Q| \leq (r_b^d + 2) |P|$,
- Q is also a partial coloring,
- the rank of Q is at least $\alpha^{k+1}_{(r_\gamma)}^{(|F|)}$,
- $d(Q) \geq \min \{1, \eta + d(P)\}$.

Proof. For each pair $(X, S) \in P$ with $X \neq \emptyset$ and $S \neq \emptyset$, apply Lemma 12. Since $k \leq \frac{1}{\eta}$, $|S| \geq \alpha^k \binom{|F|}{r_\gamma} \geq \alpha^{1/\eta} \binom{|F|}{r_\gamma} \geq \beta \binom{|F|}{r_\gamma}$. Lemma 12 produces Y_1, \dots, Y_n, Z and T_1, \dots, T_n with $n \leq r_b^d + 1$. We replace the pair (X, S) with the pairs $(Y_1, T_1), \dots, (Y_n, T_n), (Z, \emptyset)$. The resulting partition satisfies all the required properties. \square

We can now easily prove Theorem 11.

Proof of Theorem 11. By assumption, (B, γ, F) satisfies Conditions 1 and 2. Start with the partition $P = \left\{ (V(B), \binom{F}{r_\gamma}) \right\}$ and apply Corollary 13 repeatedly until the partition satisfies $d(P) = 1$. Since the value of $d(P)$ increases by η at each step, the partition is refined at most $1/\eta$ times, and so the resulting partition P has at most $(r_b^d + 2)^{1/\eta}$ parts. Consider a part $(X, S) \in P$. If $S = \emptyset$, then since P is a partial coloring $B[X]$ must be independent, so $\chi(B[X]) = 1$. If $S \neq \emptyset$, we know that $|S| \geq \beta \binom{|F|}{r_\gamma}$, which by the choice of β and L_1 forces a copy of H in S . Since $d(X, S) = 1$ we must have $S \subseteq \gamma(x)$ for every $x \in X$, so that a matching of size d in $B[X]$ witnesses that $\dim_H(B, \gamma, F) \geq d$. Therefore, the maximum size of a matching in $B[X]$ is $d - 1$. If $r_b = 1$, then $B[X]$ has at most $d - 1$ edges, so the total number of edges of B is $d(r_b^d + 2)^{1/\eta}$. If $r_b \geq 2$, then since the size of a maximal matching in $B[X]$ is $d - 1$, it is the case that $\chi(B[X]) \leq r_b(d - 1) + 1$. This implies that the chromatic number of B is at most $r_b d (r_b^d + 2)^{1/\eta}$. \square

All that remains is to prove Lemma 12. Before proving this lemma, we make some definitions. If $E_1, \dots, E_t \in B$ and $S \subseteq \binom{F}{r_\gamma}$, then the **minimum section density of E_1, \dots, E_t with respect to S** is

$$\delta(E_1, \dots, E_t, S) = \min \left\{ \frac{|\gamma(x_1) \cap \dots \cap \gamma(x_t) \cap S|}{|S|} : x_1 \in E_1, \dots, x_t \in E_t \right\}.$$

Notice that if E_1, \dots, E_d are disjoint, $\delta(E_1, \dots, E_d, S) > 0$, S is a constant fraction of $\binom{F}{r_\gamma}$, and F is large, then E_1, \dots, E_d witness that $\dim_H(B, \gamma, F) \geq d$. Define constants ψ_1, \dots, ψ_d with $\psi_1 = 1$ by $\psi_{m+1} = \frac{1}{4}\epsilon\psi_m$ for $1 \leq m \leq d - 1$.

For the proof of Lemma 12, recall the outline from Section 5.1. We will greedily select edges E_1, \dots, E_i as long as we can maintain that (using our new notation) $\delta(E_1, \dots, E_i, S)$ is large enough to force a copy of H . Since $\dim_H(B, \gamma, F) < d$, the greedy algorithm must terminate before choosing d edges. Once the greedy algorithm terminates, we will let the sets T_1, \dots, T_{n-1} (the sets that we must find to prove Lemma 12) be all sections of $x_1 \in E_1, \dots, x_i \in E_i$. If we let Y_i be the set of vertices y such that $|\gamma(y) \cap T_i| / |T_i|$ is at least $d(X, S) + \eta$, we will satisfy almost all of the requirements in Lemma 12. We need only prove that if Z are the vertices not in any Y_i then $B[Z]$ is independent. We do this by showing that if $B[Z]$ contained an edge E we could have continued the greedy algorithm by selecting E . In order for this to be true, we need the greedy algorithm to require a weaker lower bound on $\delta(E_1, \dots, E_{i+1}, S)$ than the algorithm required on $\delta(E_1, \dots, E_i, S)$. The constants ψ_i are used to define the greedy algorithm: the greedy algorithm will select edges while maintaining $\delta(E_1, \dots, E_i, S) \geq \epsilon\psi_i$. Since $\psi_{i+1}/\psi_i = \frac{1}{4}\epsilon$, we lose a fraction of ϵ in each step. The careful choice of L_1 guarantees that even after losing a fraction of ϵ for the d steps per refinement over a maximum of $1/\eta$ refinements we still have enough edges to force a copy of H .

Proof of Lemma 12. Start by greedily selecting disjoint edges E_1, \dots, E_i of $B[X]$ such that $\delta(E_1, \dots, E_i, S) \geq \epsilon\psi_i$. Since for every $x \in V(B)$

$$\frac{|\gamma(x) \cap S|}{|S|} \geq d(X, S) \geq \epsilon\psi_1,$$

the greedy algorithm can start with any edge E_1 in $B[X]$. Assume the greedy algorithm has selected E_1, \dots, E_m with $\delta(E_1, \dots, E_m, S) \geq \epsilon\psi_m$ but for every other edge E in $B[X]$ disjoint from E_1, \dots, E_m , we have $\delta(E_1, \dots, E_m, E, S) < \epsilon\psi_{m+1}$.

First, we prove that $\dim_H(B, \gamma, F) \geq m$. Since $d(E_1, \dots, E_m, S) \geq \epsilon\psi_m \geq \epsilon\psi_d$, we have that every section of $x_1 \in E_1, \dots, x_m \in E_m$ has size at least $\epsilon\psi_d |S| \geq \epsilon\psi_d \beta \binom{|F|}{r_\gamma}$. By the choice of L_1 , the section of x_1, \dots, x_m contains a copy of H , and so $m \leq d$. We make the following definitions.

- Let R_1, \dots, R_t be all r_b^m sections of $v_1 \in E_1, \dots, v_m \in E_m$ intersected with S . If some $I \subseteq F$ appears in more than one R_i , remove it from all but the least indexed R_i .
- For $1 \leq i \leq t$, start with $T_i = R_i$ and remove elements from T_i (recall that elements of T_i are subsets of F) until $|T_i|$ is smaller than $2\epsilon |S|$. (If R_i is already smaller than $2\epsilon |S|$, nothing needs to be removed.)
- Let $T_{t+1} = S \setminus T_1 \setminus \dots \setminus T_t$.
- For $1 \leq i \leq t+1$, define

$$Y_i = \left\{ x \in X : \frac{|\gamma(x) \cap T_i|}{|T_i|} \geq \min \{1, \eta + d(X, S)\} \right\}.$$

If some x appears in more than one Y_i , remove it from all but the least indexed Y_i .

- Let $Z = X \setminus Y_1 \setminus \dots \setminus Y_{t+1}$.

By the definition of Y_i , $d(Y_i, T_i) \geq \min\{1, \eta + d(X, S)\}$. Therefore, to finish the proof we need to check that $|T_i| \geq \alpha |S|$ and $B[Z]$ is independent.

Claim: $|T_i| \geq \alpha |S|$ for all $1 \leq i \leq t+1$.

Proof. Since $\delta(E_1, \dots, E_m, S) \geq \epsilon\psi_m$, before removing anything from R_i , each R_i has size at least $\epsilon\psi_m |S|$. Consider some $I \in R_i \cap R_j$ for some $j \neq i$. Since $j \neq i$, there must be some E_k such that R_i selects $\gamma(x)$ and R_j selects $\gamma(y)$ for $x \neq y \in E_k$. Thus $I \in \gamma(x) \cap \gamma(y)$, which has size at most $\lambda \binom{|F|}{r_\gamma}$ by Condition 2. In other words, every element removed from R_i is contained in $\gamma(x) \cap \gamma(y)$ for some $x \neq y \in E_k$. There are at most $mr_b(r_b - 1)$ choices of $x \neq y \in E_k$, so the maximum number of elements removed from R_i is $mr_b^2 \lambda \binom{|F|}{r_\gamma}$. Since $m \leq d$ and $|S| \geq \beta \binom{|F|}{r_\gamma}$, we remove at most $dr_b^2 \lambda \beta^{-1} |S|$ elements from R_i . By the choice of constants, $dr_b^2 \lambda \beta^{-1} \leq \frac{1}{2} \epsilon\psi_d \leq \frac{1}{2} \epsilon\psi_m$, so

$$|R_i| \geq \frac{1}{2} \epsilon\psi_m |S|. \tag{1}$$

Since $\psi_m \leq 1$ and we remove elements from R_i to form T_i only if $|R_i| \geq 2\epsilon|S|$, equation (1) implies

$$|T_i| \geq \frac{1}{2}\epsilon\psi_m|S|, \quad (2)$$

which is at least $\alpha|S|$ by the choice of α . Now consider the size of T_{t+1} . Since each T_i with $i \leq t$ has size at most $2\epsilon|S|$ and we assumed that $\epsilon < \frac{1}{4}t^{-1}$ in Condition 1, the set T_{t+1} has at least $\frac{1}{2}|S| \geq \alpha|S|$ elements. \square

Claim: $B[Z]$ is independent.

Proof. Assume E is an edge in $B[Z]$. We would like to show that there exists some $x \in E$ and some T_j such that

$$\frac{|\gamma(x) \cap T_j|}{|T_j|} \geq \min\{1, \eta + d(X, S)\}, \quad (3)$$

since this would show that $x \in Y_j$, contradicting that $x \in Z$. Assume E intersects some E_i for some $1 \leq i \leq m$, with $x \in E \cap E_i$. Since $x \in E_i$ there are many sections R_j that select x , since the sections R_j were formed by choosing one vertex from each of E_1, \dots, E_d . Fix some such section R_j that selects x , in which case $R_j \subseteq \gamma(x)$. Then $T_j \subseteq R_j \subseteq \gamma(x)$ and $|\gamma(x) \cap T_j| / |T_j| = 1$ so (3) is satisfied.

Now assume E is disjoint from E_1, \dots, E_m . Since the greedy algorithm could not continue, $\delta(E_1, \dots, E_m, E, S) < \epsilon\psi_{m+1}$, which implies that there exists some $v_1 \in E_1, \dots, v_m \in E_m, x \in E$ such that

$$|\gamma(v_1) \cap \dots \cap \gamma(v_m) \cap \gamma(x) \cap S| < \epsilon\psi_{m+1}|S|.$$

By the definition of T_i , there exists some T_i such that $T_i \subseteq \gamma(v_1) \cap \dots \cap \gamma(v_m) \cap S$. Therefore,

$$|\gamma(x) \cap T_i| < \epsilon\psi_{m+1}|S| \leq \frac{2\psi_{m+1}}{\psi_m}|T_i|,$$

where the last inequality uses (2). Assume that for every $j \neq i$, (3) fails. Then

$$|\gamma(x) \cap S| = |\gamma(x) \cap T_i| + \sum_{j \neq i} |\gamma(x) \cap T_j| \leq \frac{2\psi_{m+1}}{\psi_m}|T_i| + \sum_{j \neq i} (\eta + d(X, S))|T_j|.$$

Dividing through by $|\gamma(x) \cap S|$ we obtain

$$1 \leq \frac{2\psi_{m+1}}{\psi_m} \frac{|T_i|}{|S|} \frac{|S|}{|\gamma(x) \cap S|} + (\eta + d(X, S)) \left(1 - \frac{|T_i|}{|S|}\right) \frac{|S|}{|\gamma(x) \cap S|}.$$

Because $|S| / |\gamma(x) \cap S| \leq \frac{1}{d(X, S)} \leq \frac{1}{\epsilon}$,

$$1 \leq \frac{2\psi_{m+1}}{\psi_m \epsilon} \frac{|T_i|}{|S|} + \left(\frac{\eta}{\epsilon} + 1\right) \left(1 - \frac{|T_i|}{|S|}\right). \quad (4)$$

Let $w = |T_i|/|S|$. The right hand side of the above inequality is a weighted average of $\frac{2\psi_{m+1}}{\psi_m\epsilon}$ and $(1 + \frac{\eta}{\epsilon})$:

$$\frac{2\psi_{m+1}}{\psi_m\epsilon}w + \left(1 + \frac{\eta}{\epsilon}\right)(1 - w).$$

Since $2\psi_{m+1}/(\psi_m\epsilon) = \frac{1}{2} < 1 + \frac{\eta}{\epsilon}$, this will be maximized when w is as small as possible. By (2), $w \geq \alpha$, so

$$\frac{2\psi_{m+1}}{\psi_m\epsilon}\alpha + \left(1 + \frac{\eta}{\epsilon}\right)(1 - \alpha) < \frac{1}{2}\alpha + 1 + \frac{\eta}{\epsilon} - \alpha \leq 1 + \frac{\eta}{\epsilon} - \frac{1}{2}\alpha < 1.$$

This implies that for any $w \geq \alpha$, the inequality in (4) is false. This contradiction shows that there must be some $j \neq i$ such that $|\gamma(x) \cap T_j|/|T_j|$ is at least $\eta + d(X, S)$, which contradicts that E is contained in $B[Z]$. \square

Thus $B[Z]$ is independent and the proof is complete. \square

5.3 Fiber bundles with large overlap

In the previous section, we proved Theorem 8 restricted to fiber bundles that satisfy Condition 2. To prove Theorem 8, we will divide the edges of B into two pieces. Let B' be the subset of edges of B that satisfy Condition 2; that is for every $x \neq y \in E \in B'$, $\gamma(x) \cap \gamma(y)$ has density at most λ . We apply Theorem 11 to (B', γ, F) to bound the chromatic number of B' . For the remaining edges, we will merge x and y into a new vertex z if $\gamma(x) \cap \gamma(y)$ has density at least λ (we define $\gamma(z)$ to be $\gamma(x) \cap \gamma(y)$). Let (M, ψ, F) be the fiber bundle after merging all such vertices. Since all edges of $B - B'$ have some such pair x, y , all edges of M will have size at most $r_b - 1$. Then we apply induction on r_b to bound the chromatic number of M . To be able to apply induction, we need to verify that $\dim_H(M, \psi, F) \leq \dim_H(B, \gamma, F)$ and that there is a lower bound on the density of $\gamma(m)$ for $m \in V(M)$. The definition of $\gamma(z) = \gamma(x) \cap \gamma(y)$ satisfies both of these requirements. First, $\dim_H(M, \psi, F) \leq \dim_H(B, \gamma, F)$ because any copy of H in $\gamma(z)$ will be in both $\gamma(x)$ and $\gamma(y)$. Also, there is a lower bound on the density of $\gamma(z)$ because we only merge if $\gamma(x) \cap \gamma(y)$ has density at least λ . The magic in this proof is that Condition 2, the extra requirement needed in the previous section, fits exactly with the requirements to be able to apply induction after merging.

For technical reasons, our induction statement needs to be slightly stronger than Theorem 8; we no longer assume B is a uniform hypergraph. Instead, we allow the edges of B to have size between one and r_b . This is because after merging, all we know is that the edges have size between one and $r_b - 1$. This is also why we need to allow 1-uniform hypergraphs in Theorem 11.

Theorem 14. *Let $r_b, r_\gamma \geq 1$, $d \in \mathbb{Z}^+$, $0 < \epsilon < 1$, and H be an r_γ -uniform hypergraph with zero Turán density. Then there exists constants $K_1 = K_1(r_b, r_\gamma, d, \epsilon, H)$ and $K_2 = K_2(r_b, r_\gamma, d, \epsilon, H)$ such that the following holds. Let (B, γ, F) be any fiber bundle where the edges of B have size between 1 and r_b , $\dim_H(B, \gamma, F) < d$, and for all $b \in V(B)$, $\gamma(b)$ is an r_γ -uniform hypergraph and $|\gamma(b)| \geq \epsilon \binom{|F|}{r_\gamma}$. Let A be the set of edges of size 1 in B . If $|F| \geq K_1$, then $|A| \leq K_2$ and $\chi(B - A) \leq K_2$.*

Proof. The proof is by induction on r_b . When $r_b = 1$, we can directly apply Theorem 11, since Condition 2 is trivially satisfied. So assume $r_b \geq 2$, and let $\lambda = \lambda(r_b, r_\gamma, d, \epsilon, H)$ be the constant from Theorem 11. Define a subhypergraph B' of B as follows:

$$B' = \left\{ E \in B : |E| = r_b \text{ and } \forall x \neq y \in E \quad |\gamma(x) \cap \gamma(y)| \leq \lambda \binom{|F|}{r_\gamma} \right\}.$$

Then (B', γ, F) is an (r_b, r_γ) -uniform fiber bundle to which we can apply Theorem 11 to bound the chromatic number of B' . To complete the proof, we will bound the chromatic number of $B - B'$.

Initially, let $B_0 = B - B'$ and $\gamma_0 = \gamma$. At stage i , assume we have defined (B_i, γ_i, F) . Let E be some edge of B_i with $|E| = r_b$, where there exists $x \neq y \in E$ be such that $|\gamma_i(x) \cap \gamma_i(y)| \geq \lambda \binom{|F|}{r_\gamma}$. We form $(B_{i+1}, \gamma_{i+1}, F)$ by merging the vertices x and y . More precisely, let z be a new vertex and define

$$\begin{aligned} V(B_{i+1}) &= V(B_i) - x - y + z, \\ E(B_{i+1}) &= \{E \in B_i : E \cap \{x, y\} = \emptyset\} \cup \{E - x - y + z : E \in B_i, E \cap \{x, y\} \neq \emptyset\}, \\ \gamma_{i+1}(w) &= \begin{cases} \gamma_i(w) & w \neq z, \\ \gamma_i(x) \cap \gamma_i(y) & w = z. \end{cases} \end{aligned}$$

We do this until every edge of B_i has size at most $r_b - 1$; say this occurs at step s .

Through this modification, the dimension cannot increase. Consider step i , when we merge the vertices x and y in (B_i, γ_i, F) to form $(B_{i+1}, \gamma_{i+1}, F)$. Let S be any section in (B_i, γ_i, F) that selects x or y , and let S' be a section in $(B_{i+1}, \gamma_{i+1}, F)$ that is identical to S except that it selects z instead of x or y . If S' contains a copy of H , then this copy of H is in $\gamma(z)$, which implies it is in both $\gamma(x)$ and $\gamma(y)$. Therefore, H is in S .

For each vertex z we add, we have $|\gamma(z)| \geq \lambda \binom{|F|}{r_\gamma}$. Therefore, we can apply induction on r_b to (B_s, γ_s, F) using $\epsilon = \lambda$ to bound the chromatic number of B_s . We now consider “un-merging” the vertices of B_s to obtain a coloring of B_0 . Consider un-merging $z \in V(B_i)$ to obtain $x, y \in V(G_{i-1})$. If z was contained in an edge of size one, we color x and y using two new colors. Otherwise, we color x and y the same color as z . After the un-merging, we will have a proper coloring of $B_0 = B - B'$. By induction, there is a bounded number of edges of size one, so we will use a bounded number of new colors.

The colorings on B_0 and B' can be combined to obtain a coloring of B using a bounded number of colors. Let K_1 (the minimum size of F) be the maximum between the required size from Theorem 11 and the size required by induction. Lastly, any edge of size one in B will also appear in B_s , and by induction B_s has a bounded number of edges of size one. \square

6 Extremal results for critical hypergraphs

In this section, we prove Theorems 2 and 4. First, it is easy to see that C_{2k+1}^r is mono near r -partite; the edge E_{2k+1} in C_{2k+1}^r will be the special edge. Also, E_{2k+1} has $r - 2$ vertices, $v_{rk+2}, \dots, v_{rk+r-1}$, that have degree one. Thus to complete the proof of Theorem 4 we need only prove that C_{2k+1}^3 and C_{2k+1}^4 are stable with respect to $T_3(n)$ and $T_4(n)$. One tool we will use is the hypergraph removal lemma of Gowers, Nagle, Rödl, and Skokan [15, 26, 29, 30, 34].

Theorem 15. *For every integer $r \geq 2$, $\epsilon > 0$, and r -uniform hypergraph H , there exists a $\delta > 0$ such that any r -uniform hypergraph with at most $\delta n^{|V(H)|}$ copies of H can be made H -free by removing at most ϵn^r edges.*

The second tool we will use is supersaturation, proved by Erdős and Simonovits [7]. There are several equivalent formulations of supersaturation, the one we will use is the following.

Theorem 16. *[7, Corollary 2] Let K_{t_1, \dots, t_r}^r be the complete r -uniform, r -partite hypergraph with part sizes t_1, \dots, t_r . Let $t = \sum t_i$. For every $\epsilon > 0$, there exists a $\delta = \delta(r, t, \epsilon)$ such that any r -uniform hypergraph with at least ϵn^r edges contains at least δn^t copies of K_{t_1, \dots, t_r}^r .*

For any hypergraph H , let $H(t)$ denote the hypergraph obtained from H by blowing up each vertex into an independent set of size t . An easy extension of supersaturation is the following (see Theorem 2.2 in the survey by Keevash [16]).

Corollary 17. *For every $r, t \geq 2$, $\epsilon > 0$, and r -uniform hypergraph H , there exists an n_0 such that if $n \geq n_0$ and G is an n -vertex, r -uniform hypergraph which contains at least $\epsilon n^{|V(H)|}$ copies of H , then G contains a copy of $H(t)$.*

Next, we will need stability results for F_5 and the book $B_{4,2}$, proved by Keevash and the last author [18] and Pikhurko [28] respectively. Let the **book** $B_{r,m}$ be the r -uniform hypergraph with vertices $x_1, \dots, x_{r-1}, y_1, \dots, y_r$ and hyperedges $\{x_1, \dots, x_{r-1}, y_i\}$ for $1 \leq i \leq m$ and $\{y_1, \dots, y_r\}$. Note that $F_5 = B_{3,2}$.

Theorem 18. *[18] F_5 is stable with respect to $T_3(n)$.*

Theorem 19. *[28] $B_{4,2}$ is stable with respect to $T_4(n)$.*

The last piece of the proof of Theorem 4 is the following lemma.

Lemma 20. *If H is an r -uniform hypergraph that is stable with respect to $T_r(n)$, and F is a subhypergraph of $H(t)$ for some t , then F is also stable with respect to $T_r(n)$.*

Proof. If F is contained in $T_r(n)$, then for large n there is no r -uniform, F -free hypergraph with at least $(1 - \delta)t_r(n)$ edges, so F is vacuously stable with respect to $T_r(n)$. So assume F is not a subhypergraph of $T_r(n)$. Let h denote the number of vertices in H and let $\epsilon > 0$ be fixed. We need to show how to define δ such that if G is a F -free hypergraph with at least $t_r(n) - \delta n^r$ edges, it differs from $T_r(n)$ in at most ϵn^r edges.

Since H is stable with respect to $T_r(n)$, there exists an $\alpha \leq \epsilon/2$ such that if G' has at least $t_r(n) - 2\alpha n^r$ edges and contains no copy of H , then G' differs from $T_r(n)$ in at most $\epsilon n^r/2$ edges. By Theorem 15, there exists $\beta = \beta(\alpha)$ such that if there are at most βn^h copies of H in G then by deleting at most αn^r edges of G we can remove all copies of H . Lastly, choose $\delta \ll \beta$.

Now, fix some G that contains no copy of F and has at least $t_r(n) - \delta n^r$ edges. Because G contains no copy of F it contains no copy of $H(t)$. Therefore, by Corollary 17 there are at most βn^h copies of H in G . By Theorem 15, we may therefore delete αn^r edges in order to find a subhypergraph G' of G that contains no copy of H . Notice that G' has at least $t_r(n) - (\delta + \alpha)n^r$ edges, and $(\delta + \alpha) < 2\alpha$, so G' differs from $T_r(n)$ in at most $\epsilon n^r/2$ edges. Therefore, G differs from $T_r(n)$ in at most $(\alpha + \epsilon/2)n^r$ edges, and $\alpha + \epsilon/2 < \epsilon$. \square

It is easy to see that C_{2k+1}^r is a subhypergraph of $B_{r,2}(k)$. Thus Theorem 18 combined with Lemma 20 shows that C_{2k+1}^3 is stable with respect to $T_3(n)$ and similarly Theorem 19 combined with Lemma 20 shows that C_{2k+1}^4 is stable with respect to $T_4(n)$, which completes the proof of Theorem 4. Frankl and Füredi [10] determined $\pi(B_{r,2})$ for $r = 5, 6$, and the values come from blowups of small designs. In particular, $\pi(B_{r,2}) \neq t_r(n)$ for $r = 5, 6$. This means a different method must be used when attempting to prove that C_{2k+1}^r is critical for $r \geq 5$.

Proof of Theorem 2. Let H be a critical n -vertex, r -uniform hypergraph. Suppose H has h vertices and assume that E is the special edge of H . Suppose G is an H -free, r -uniform, n -vertex hypergraph with $|G| \geq t_r(n)$. We would like to show that $G = T_r(n)$. Partition the vertices of G into parts X_1, \dots, X_r such that the number of edges with one vertex in each X_i is maximized. Let $\epsilon_1 = (2r)^{-h}$, let $\epsilon_2 = \epsilon_1/8r^3$, let $\delta = \delta(r, h, \epsilon_2)$ from Theorem 16, and let $\epsilon < 2^{-2r}\epsilon_1\epsilon_2\delta$. Organize r -sets of vertices into the following sets.

- Let M be the set of r -sets with one vertex in each of X_1, \dots, X_r that are not edges of G (the missing cross-edges).
- Let B be the collection of edges of G that have at least two vertices in some X_i (the bad edges).
- Let $G' = G - B + M$, so that G' is a complete r -partite hypergraph.
- Let $B_i = \{W \in B : |W \cap X_i| \geq 2\}$.

Since $B = \cup_i B_i$, there is some B_i which has size at least $\frac{1}{r}|B|$. Assume without loss of generality that $|B_1| \geq \frac{1}{r}|B|$. For $a \in X_1$, make the following definitions.

- $B_a = \{W \in B_1 : a \in W\}$.
- Let $C_{a,i}$ be the edges in B_a which have exactly two vertices in B_1 and exactly one vertex in each X_j with $j \geq 1$ and $j \neq i$.
- Let $D_a = B_a \setminus C_{a,2} \setminus \dots \setminus C_{a,r}$.

First, $|B| < \epsilon n^r$ because G is stable with respect to $T_r(n)$. Also, since $|G| \geq t_r(n)$, the number of r -sets in M is at most the number of edges in B , so $|M| \leq |B| < \epsilon n^r$.

In the rest of the proof, we will assume that B is non-empty and then count the r -sets in M in several different ways. Our counting will imply that $|M| \geq \epsilon n^r$, and this contradiction will force $B = \emptyset$ and so $G = T_r(n)$. We will count r -sets in M by counting embeddings of $H - E$ into G' that also map E to some element of B . Since G is H -free, each embedding must use at least one edge in M . Let Φ be the collection of embeddings $\phi : V(H) \rightarrow V(G')$ of $H - E$ into G' , by which we mean that ϕ is an injection and for all $F \in H$, $\phi(F) = \{\phi(x) : x \in F\} \in G'$. We say that $\phi \in \Phi$ is **W -special** if $\phi(E) = W$ and **a -avoiding** if $a \in V(G)$ and some degree one vertex in E is mapped to a . If $W \in B$ and ϕ is W -special, then ϕ must use at least one edge of M . Call one of these edges the **missing edge of ϕ** .

Claim 1: For $\phi \in \Phi$ and $v \in V(H)$, there are at least $\frac{1}{2r}n$ embeddings $\phi' \in \Phi$ where $\phi(x) = \phi'(x)$ for $x \neq v$ and $\phi(v) \neq \phi'(v)$.

Proof. This follows easily because G' is a complete r -partite hypergraph for which each class has size about n/r , and $\phi(v)$ can be replaced by any unused vertex in the X_i that contains $\phi(v)$. \square

Fix some $W \in B$, and consider when there exists a W -special embedding of $H - E$. Since $W \in B_i$ for some i , let $w_1 \neq w_2 \in W \cap X_i$. Then there exists an embedding of $H - E$ where w_1 and w_2 are used for the non degree one vertices in the special edge of H . Since the other vertices in the special edge have degree zero in $H - E$, the vertices in the special edge can then be embedded to W . Thus for any $W \in B$, by Claim 1 there are at least $\epsilon_1 n^{h-r}$ W -special embeddings of $H - E$, since we can vary any vertex of H not in W . The situation with a -avoiding is more complicated. If $W \in C_{a,i}$, then the only choice of w_1 and w_2 that we are guaranteed to have are the two vertices in $W \cap X_1$, one of which is a . Thus in a W -special embedding, the only way we can guarantee an embedding is by mapping a non-degree one vertex to a . Therefore, only when $W \in D_a$ can we guarantee that there exists at least $\epsilon_1 n^{h-r}$ W -special, a -avoiding embeddings of $H - E$.

Claim 2: For every $a \in X_1$, $|D_a| \leq \epsilon_2 n^{r-1}$.

Proof. Assume there exists some $a \in X_1$ with $|D_a| \geq \epsilon_2 n^{r-1}$. We count a -avoiding, W -special embeddings of $H - E$ into G' where $W \in D_a$. For each $W \in D_a$, we argued above that there are at least $\epsilon_1 n^{h-r}$ embeddings. Since $|D_a| \geq \epsilon_2 n^{r-1}$, the number of a -avoiding embeddings which are W -special for some $W \in D_a$ is at least $\epsilon_1 \epsilon_2 n^{r-1} \cdot n^{h-r} = \epsilon_1 \epsilon_2 n^{h-1}$.

Fix some $L \in M$. We want to count the number of a -avoiding embeddings which are W -special for some $W \in D_a$ and have missing edge L . An upper bound on the number of such embeddings will be the number of choices for W times the number of choices for the $h - |W \cup L|$ vertices of H mapped outside $W \cup L$. Since all these embeddings are a -avoiding, L cannot contain a . For each $0 \leq \ell \leq r$, there exists at least $\binom{r}{\ell}$ choices for the intersection between L and W , at most $n^{r-\ell-1}$ choices of $W \in D_a$ with $|W \cap L| = \ell$ (here it is crucial that $a \in W$ and $a \notin L$), and at most $n^{h-2r+\ell}$ choices for the vertices of H not in $W \cup L$. Thus each $L \in M$ can kill at most $2^{-r} n^{h-r-1}$ embeddings. Since there are at least $\epsilon_1 \epsilon_2 n^{h-1}$ embeddings, M must have size at least $2^{-r} \epsilon_1 \epsilon_2 n^r$, contradicting the choice of ϵ . \square

Claim 3: For every $a \in X_1$ and every $2 \leq i \leq r$, $|C_{a,i}| \leq \epsilon_2 n^{r-1}$.

Proof. Assume there exists some a and i with $|C_{a,i}| \geq \epsilon_2 n^{r-1}$. The proof is similar to the proof of Claim 2, except now we cannot count a -avoiding embeddings. In the previous claim, we used the a -avoiding property to imply that the missing edge does not contain a . In this proof, we will instead guarantee that the missing edge cannot contain a by only counting embeddings which map all neighbors of $\phi^{-1}(a)$ into G .

Let v be one of the non degree one vertices in the special edge of H , and define $H_v = \{F \in H : v \in F, F \neq E\}$, that is all edges of H containing v which are not the special edge. Let $Z_a = \{F \in G \setminus B : a \in F\}$, that is all cross-edges of G which contain a . We now count embeddings $\phi \in \Phi$ which are W -special for some $W \in C_{a,i}$, map v to a , and all edges of H_v are mapped to edges in Z_a . For these embeddings, since edges in H_v are mapped to edges in $Z_a \subseteq G$, the missing edge cannot contain a .

First, $|Z_a| \geq |C_{a,i}|$, because otherwise we could move a to X_i and increase the number of edges across the partition and we chose the partition X_1, \dots, X_r to maximize the number

of cross-edges. Let $H' = \{F - v : F \in H_v\}$ and $Z' = \{F - a : F \in Z_a\}$. Then H' and Z' are $(r - 1)$ -uniform, $(r - 1)$ -partite hypergraphs, and Z' has at least $|C_{a,i}| \geq \epsilon_2 n^{r-1}$ edges. Let $t = |V(H')|$. Then Theorem 16 shows that Z' contains at least δn^t copies of H' , so there are at least $\epsilon_2 n^{r-1} \cdot \delta n^t \cdot \epsilon_1 n^{h-r-t} = \epsilon_1 \epsilon_2 \delta n^{h-1}$ embeddings of $H - E$ which are W -special for some $W \in C_{a,i}$, map v to a , and the edges in H_v are embedded into Z_a .

Now fix $L \in M$, and consider how many of these embeddings have L as their the missing edge. The computation is almost the same as in the previous claim. For each ℓ_1, ℓ_2 , there are $\binom{r}{\ell_1}$ choices for $L \cap W$, there are $\binom{r}{\ell_2}$ choices for $L \cap \phi(H_v)$, there are $n^{r-1-\ell_1}$ choices for W (here we use that L does not contain a), $n^{t-\ell_2}$ choices for $\phi(H_v)$, and $n^{h-2r-t+\ell_1+\ell_2}$ choices for the other vertices of H . Thus each L can kill at most $2^{2r} n^{h-r-1}$ embeddings. Since there are at least $\epsilon_1 \epsilon_2 \delta n^{h-1}$ embeddings, M must have size at least $2^{-2r} \epsilon_1 \epsilon_2 \delta n^r$, contradicting the choice of ϵ . \square

Claims 2 and 3 imply that $|B_a| < 2r\epsilon_2 n^{r-1}$ for each a . Define

$$A = \{a \in X_1 : d_M(a) \geq 2r^2 \epsilon_2 n^{r-1}\}.$$

As in the proofs of the previous two claims, we would like to count embeddings of $H - E$ to obtain a lower bound on $|M|$. Once again, the main difficulty is controlling how the missing edge can intersect W . If there were some W with $W \cap A = \emptyset$, there will be few missing edges intersecting this W , which is how we will overcome this difficulty in this part of the proof.

Claim 4: There exists some $W \in B_1$ with $W \cap A = \emptyset$.

Proof. Assume that every $W \in B_1$ contains an element of A . Then $\sum_{a \in A} |B_a| \geq |B_1|$. Since $|B_a| < 2r\epsilon_2 n^{r-1}$ for every a , we have the following contradiction.

$$2r\epsilon_2 n^{r-1} |A| > \sum_{a \in A} |B_a| \geq |B_1| \geq \frac{1}{r} |B| \geq \frac{1}{r} |M| \geq \frac{1}{r} \sum_{a \in A} d_M(a) \geq \frac{2r^2 \epsilon_2}{r} n^{r-1} |A|.$$

\square

We now finish the proof by counting the W -special embeddings whose missing edge does not intersect W . There are at least $\epsilon_1 n^{h-r}$ embeddings which are W -special by Claim 1. If at least half of these have missing edge intersecting W , then W would contain a vertex in A . Thus there are at least $\frac{\epsilon_1}{2} n^{h-r}$ W -special embeddings where the missing edge does not intersect W . Each $L \in M$ can kill at most n^{h-2r} such embeddings, so M has at least $\frac{\epsilon_1}{2} n^r$ elements, contradicting the choice of ϵ . \square

7 Chromatic threshold of F_5 -free hypergraphs

7.1 An upper bound on the chromatic threshold of F_5 -free graphs

In this section, we prove the upper bound in Theorem 7. As in Section 4, we will give an upper bound on the chromatic threshold by first proving that large dimension forces a copy of F_5 , and then by applying Theorem 8. Let (B, γ, F) be an (r_b, r_γ) -uniform fiber bundle, and make the following definition. A **cut** in (B, γ, F) is a pair (X, S) such that $X \subseteq V(B)$,

$S \subseteq \binom{F}{r_\gamma}$, and if $\gamma(x) \cap S \neq \emptyset$, then $x \in X$. In other words, the fibers that intersect S come exclusively from X . A **k -cut** is a cut (X, S) with $|X| \leq k$. The size of a k -cut is the size of $|S|$.

We now sketch the proof of the upper bound in Theorem 7. Let G be an n -vertex, 3-uniform, F_5 -free hypergraph with minimum degree at least $c \binom{n}{2}$. Let (G, γ, F) be the neighborhood bundle of G , let $H = K_{q,q}$, and assume $\dim_H(G, \gamma, F)$ is large. We would like to find a copy of F_5 in G . We first use the fact that $\dim_H(G, \gamma, F)$ is large to find a set U of vertices of G such that $G[U]$ has small strong independence number. We then argue that because the minimum degree is large, there must be some vertices x, y such that $N(x, y) = \{z : xyz \in G\}$ has large intersection with U . Next, we show that since $N(x, y)$ has large intersection with U and $G[U]$ has small strong independence number, there must be an edge E with at least two vertices in $N(x, y) \cap U$, which gives a copy of F_5 .

The best upper bound on the chromatic threshold will come from the lowest required minimum degree needed in the above proof. The minimum degree is used above to prove that there exists some x, y with $N(x, y) \cap U$ large. If we can find a large cut (X, S) in (G, γ, F) and we make U large enough, we could remove X from U while still maintaining all the useful properties of U . Then for all $\{x, y\} \in S$, we know that $N(x, y) \cap (U - X) = \emptyset$. Since there are now fewer pairs $\{x, y\}$ in $\binom{F}{2}$ with $N(x, y) \cap (U - X) \neq \emptyset$, we can require a weaker lower bound on the minimum degree of G to find $\{x, y\}$ with $N(x, y) \cap U$ large. In other words, the larger the cut of (G, γ, S) we can find, the better upper bound on the chromatic threshold we can prove. This is encoded in the following theorem, which computes the relationship between the minimum degree and the maximum size of a k -cut.

Theorem 21. *Let $0 < c, s < 1$ satisfy $5c \geq 1 - s$, and fix an integer k and a constant $c' > c$. Then there exists a constant $L = L(c, c', k, s)$ such that the following holds. Let G be an n -vertex, F_5 -free hypergraph with $\delta(G) \geq c' \binom{n}{2}$ and let (G, γ, F) be the neighborhood bundle of G . Assume (G, γ, F) contains a k -cut of size at least $s \binom{n}{2}$. Then $\chi(G) \leq L$.*

Note that if $s = 0$ and $c = 1/5$, then $5c \geq 1 - s$ and so this theorem directly proves an upper bound of $1/5$ on the chromatic threshold of F_5 -free hypergraphs. The first part of the proof of Theorem 21 is to find a set U with small strong independence number.

Lemma 22. *Let $\epsilon > 0$ be fixed. Then there exists constants $d = d(\epsilon)$ and $q = q(\epsilon)$ such that the following holds. Let G be an n -vertex, 3-uniform hypergraph and let (G, γ, F) be the neighborhood bundle of G . Let $H = K_{q,q}$ and assume $\dim_H(G, \gamma, F) \geq d$. Then there exists a vertex set $U \subseteq V(G)$ such that $|U| = 5d$ and the strong independence number of $G[U]$ is at most $(1 + \epsilon)d$.*

Proof. Let $d = 100 + 100/\epsilon^2$ and $q = 3d + 2 \cdot 3^d$. Since $\dim_H(G, \gamma, F) \geq d$, there exists a matching E_1, \dots, E_d such that for each $x_1 \in E_1, \dots, x_d \in E_d$ the section of $\{x_1, \dots, x_d\}$ contains a copy of $K_{q,q}$. (See Figure 2 in Section 4 for a picture of this structure.) Since $q = 3d + 2 \cdot 3^d$, from each of these 3^d copies of $K_{q,q}$ we can pick a copy of K_2 such that each K_2 is vertex disjoint from $E_1 \cup \dots \cup E_d$ and all these 3^d copies of K_2 are vertex disjoint. Now choose randomly d copies of K_2 , $\{y_1, z_1\}, \dots, \{y_d, z_d\}$ (with repetition) from these 3^d vertex disjoint copies of K_2 , and define $Z = \{y_1, \dots, y_d, z_1, \dots, z_d\}$, $U = Z \cup E_1 \cup \dots \cup E_d$. Since Z is disjoint from E_1, \dots, E_d , $|U| = 5d$. We need only show that with positive probability the strong independence number of $G[U]$ is at most $(1 + \epsilon)d$.

Notice that any strong independent set in $G[U]$ contains at most d vertices from $E_1 \cup \dots \cup E_d$ and at most d vertices from Z . Thus any strong independent set in $G[U]$ with at least $(1 + \epsilon)d$ vertices must have at least ϵd vertices in $E_1 \cup \dots \cup E_d$ and at least ϵd vertices in Z . We need to prove with positive probability that this does not occur.

For every $P \subseteq Z$ with $|P| = \epsilon d$ and every $S \subseteq E_1 \cup \dots \cup E_d$ with $|S| = \epsilon d$, let $A_{P,S}$ be the event “ $P \cup S$ is a strong independent set in G ” and $X_{P,S}$ be its indicator random variable. Let X be the sum of all indicator random variables. We claim that X is small.

For any $u \in Z$, any $1 \leq i \leq d$, and any $w \in E_i$, let $\{u, v\}$ be the copy of K_2 containing u . Then, because $\{u, v\}$ is contained in some section of $x_1 \in E_1, \dots, x_d \in E_d$, it is the case that $\{u, v, w'\} \in G$ for some $w' \in E_i$. Therefore, $\Pr[\{u, v, w\} \notin G] \leq 2/3$ so that

$$E[X] = \sum X_{P,S} \leq \binom{2d}{\epsilon d} \binom{3d}{\epsilon d} (2/3)^{\epsilon^2 d^2} \leq \left(\binom{2ed}{\epsilon d} \binom{3ed}{\epsilon d} \left(\frac{2}{3} \right)^{\epsilon d} \right)^{\epsilon d} < 1.$$

Also, since we have 3^d copies of K_2 and select only d of them, the probability some K_2 is selected twice is exponentially small. Thus with positive probability, every event $A_{P,S}$ fails which implies the strong independence number of $G[U]$ is at most $(1 + \epsilon)d$. \square

We can now prove Theorem 21.

Proof of Theorem 21. Pick ϵ so that $c' = (1 + 2\epsilon)c$ and let $d = d(\epsilon)$ and $q = q(\epsilon)$ be given by Lemma 22, and also assume that d is large enough so that $5d\epsilon > k(1 + 2\epsilon)$. Suppose that if $H = K_{q,q}$ then $\dim_H(G, \gamma, F) \leq d$. Then by Theorem 8, there exists constants $K_1 = K_1(\epsilon, d, H)$ and $K_2 = K_2(\epsilon, d, H)$ (note that K_1 and K_2 depend only on c, c', k, s) such that either $|F| < K_1$ or $\chi(G) < K_2$. Since $|F| = |V(G)|$, this implies that $\chi(G) < \max\{K_1, K_2\}$.

We can therefore assume that $\dim_H(G, \gamma, F) \geq d$. By Lemma 22, there exists a set $U \subseteq V(G)$ such that $|U| = 5d$ and the strong independence number of $G[U]$ is at most $(1 + \epsilon)d$. Let (X, S) be a k -cut of size $s \binom{n}{2}$. Let G' be the bipartite graph with partite sets $A = U \setminus X$ and $B = \binom{V(G)}{2} \setminus S$ where $\{u, \{v, w\}\}$ is an edge in G' if and only if $\{u, v, w\}$ is an edge in G . $|A| \geq 5d - |X|$, so G' contains at least $(5d - |X|)\delta(G)$ edges. $|B| = \binom{n}{2} - |S|$, so there is some $x \neq y$ such that $d_{G'}(\{x, y\})$ is at least

$$\frac{(5d - |X|)\delta(G)}{\binom{n}{2} - |S|} \geq \frac{(5d - k)(1 + 2\epsilon)c \binom{n}{2}}{(1 - s)\binom{n}{2}} \geq \frac{(5d - k)(1 + 2\epsilon)c}{5c} > (1 + \epsilon)d.$$

This implies that there is some x, y with $|N(x, y) \cap U| > (1 + \epsilon)d$. Since the strong independence number of $G[U]$ is at most $(1 + \epsilon)d$, there exists some edge E with two vertices in $N(x, y)$. Then x, y together with E form a copy of F_5 in G . This contradiction completes the proof. \square

7.2 Finding a large cut in a F_5 -free hypergraph

In order to prove the upper bound in Theorem 7, we now need to show the existence of a large cut. Note that in Theorem 21 the bound on the chromatic number depends on k

but there are no other restrictions on k . Thus to prove an upper bound on the chromatic threshold of F_5 -free graphs, one can pick any fixed integer k and ask what is the size of the largest k -cut. In the following lemma, we set $k = 5$ and prove that there exist a 5-cut of size approximately $4c^2 \binom{n}{2}$. Solving $5c = 1 - s = 1 - 4c^2$ gives $c = (\sqrt{41} - 5)/8$, the bound in Theorem 7.

We suspect that the bound on the chromatic threshold of F_5 -free hypergraphs can be improved by finding a larger cut, perhaps by increasing k . In order to achieve a bound of $c = 6/49$, we would need to find a cut of size $s \binom{n}{2}$ with $s = 1 - 5c = 539/36c^2 \approx 15c^2$.

Lemma 23. *Let $0 < c < c'$ be fixed. There exists a constant $n_0 = n_0(c, c')$ such that for all $n > n_0$ the following holds. Let G be an n -vertex, 3-uniform, F_5 -free hypergraph with $\delta(G) \geq c' \binom{n}{2}$. Let (G, γ, F) be the neighborhood bundle of G . Then (G, γ, F) has a 5-cut of size at least $4 \binom{c(n-1)}{2}$.*

Combining Theorem 21 with Lemma 23, we can prove Theorem 7.

Proof of Theorem 7. Let $c = (\sqrt{41} - 5)/8$, let $c' > c$ be fixed, and let G be any n -vertex, 3-uniform, F_5 -free graph with minimum degree at least $c' \binom{n}{2}$. Let (G, γ, F) be the neighborhood bundle of G . Let $b = (c' + c)/2$ so that $c' > b > c$. Then by Lemma 23, either $|V(G)|$ is bounded or (G, γ, F) contains a 5-cut of size at least $4 \binom{b(n-1)}{2}$. Since $b > c$, if n is large enough this is at least $4c^2 \binom{n}{2}$. Let $s = 4c^2$ and notice that $5c = 1 - s$, so that Theorem 21 implies that the chromatic number of G is bounded. \square

The first step in the proof of Lemma 23 is the following lemma.

Lemma 24. *In a graph G , we call a non-edge $uv \notin E(G)$ **good** if $N(u) \cap N(v) \neq \emptyset$. If G is a triangle-free graph with n vertices and m edges, then G has at least $m - n/2$ good non-edges.*

Proof. We prove this by induction on n . It is obviously true for $n = 1$ and $n = 2$. Now assume $n > 2$. If some component of G is not regular, then there exist vertices u, v in that component such that $u \in N(v)$ and $d(u) < d(v)$. Then $G - u$ has $n - 1$ vertices and $m - d(u)$ edges. By induction, $G - u$ has at least $m - d(u) - \frac{n-1}{2}$ good non-edges. For any vertex $w \in N(v) - u$, uw is a good non-edge, so G has at least $m - d(u) - \frac{n-1}{2} + d(v) - 1 \geq m - n/2$ good non-edges. If all components of G are regular, then pick one component K . Assume K is r -regular, and pick a vertex v in K and let $N_2(v) = \{u : \text{there exists a } P_3 \text{ connecting } u \text{ and } v\}$. If $|N_2(v)| \geq r$, then by the induction hypothesis $G - v$ has at least $m - r - \frac{n-1}{2}$ good non-edges, and since for any vertex $u \in N_2(v)$ it is the case that uv is a good non-edge, G has at least $m - r - \frac{n-1}{2} + |N_2(v)| \geq m - n/2$ good non-edges. If $|N_2(v)| < r$, then since K is triangle-free and r -regular, $K = K_{r,r}$ which has r^2 edges and $r^2 - r$ good non-edges. Now $G - K$ has $n - 2r$ vertices and $m - r^2$ edges, so by induction it has $m - r^2 - (n - 2r)/2$ good non-edges. Then G has $m - r^2 - (n - 2r)/2 + r^2 - r = m - n/2$ good non-edges. \square

Proof of Lemma 23. We examine the copies of F_4 in G where F_4 is the hypergraph with vertex set $\{1, 2, 3, 4\}$ and edges $\{1, 2, 3\}$, $\{1, 2, 4\}$, and $\{2, 3, 4\}$.

Case 1 There exists a vertex v of G such that v is not contained in any copy of F_4 . Consider $L = \gamma(v)[V(G) - v]$, which is a triangle-free graph with $n - 1$ vertices and at least $c \binom{n}{2}$ edges.

By Lemma 24, L has at least $c\binom{n}{2} - \frac{n-1}{2}$ good non-edges. Let $X = \emptyset$ and S be the set of these good non-edges. We claim that (X, S) is a cut in (G, γ, F) . Suppose for contradiction that there exists some $x \in V(G)$ and $\{u, w\} \in S$ such that $\{u, w, x\} \in G$. Pick a vertex y from $N_L(u) \cap N_L(w)$. Then u, v, w, x, y form a copy of F_5 in G , which is a contradiction.

Case 2 Every vertex of G is contained in some copy of F_4 . Pick some $U \subseteq V(G)$ such that $G[U] = F_4$, let $U = \{u_1, u_2, u_3, u_4\}$, and let $G' = \cup_{i=1}^4 \gamma(u_i)$. Consider $\gamma(u_i) \cap \gamma(u_j)$ for $i \neq j$. If $\gamma(u_i) \cap \gamma(u_j)$ contains a matching of size two, then G contains a copy of F_5 . Say $ab, cd \in \gamma(u_i) \cap \gamma(u_j)$ with a, b, c, d distinct. Then since $G[U] = F_4$, there is some edge $E = \{u_i, u_j, w\} \in G$. If $w \neq a$ and $w \neq b$, then a, b, u_i, u_j, w form a copy of F_5 and if $w = a$ or $w = b$, then c, d, u_i, u_j, w form a copy of F_5 . Thus $\gamma(u_i) \cap \gamma(u_j)$ is a star so has at most n elements. Since each $\gamma(x)$ has size at least $c'\binom{n}{2}$, G' has at least $4c'\binom{n}{2} - \binom{4}{2}n > 4c\binom{n}{2}$ edges if n is large enough.

Then G' has n vertices and at least $4c\binom{n}{2}$ edges, so there exist a vertex v whose degree in G' is at least $4c(n-1)$. Let N denote the neighborhood of v in G' and let N_1, \dots, N_4 be a partition of N such that for every $1 \leq i \leq 4$ and every vertex $w \in N_i, vw \in \gamma(u_i)$. Let $X = U \cup \{v\}$ and $S = \bigcup_{i=1}^4 \binom{N_i}{2}$, so that $|X| = 5$ and $|S| \geq 4\binom{|N|/4}{2} = 4\binom{c(n-1)}{2}$. We claim that (X, S) is a cut in (G, γ, F) . Suppose for contradiction that there exists some $z \notin X$ such that $\gamma(z) \cap S \neq \emptyset$. Pick $\{x, y\} \in \gamma(z) \cap S$, then $\{x, y\} \subseteq N_i$ for some $1 \leq i \leq 4$. Now v, u_i, x, y, z form a copy of F_5 , which is a contradiction.

From these two cases we can see that (G, γ, F) has a 5-cut of size at least $\min \left\{ c\binom{n}{2} - \frac{n-1}{2}, 4\binom{c(n-1)}{2} \right\}$. G is F_5 -free, so $c \leq 2/9$ and therefore $\min \left\{ c\binom{n}{2} - \frac{n-1}{2}, 4\binom{c(n-1)}{2} \right\} = 4\binom{c(n-1)}{2}$. \square

7.3 A construction for the lower bound

To prove a lower bound on the chromatic threshold of the family of F_5 -free hypergraphs, we need to construct an infinite sequence of F_5 -free hypergraphs with large chromatic number and large minimum degree. Our construction is inspired by a construction by Hajnal [6] of a dense triangle-free graph with high chromatic number. Hajnal's key idea was to use the Kneser graph to obtain large chromatic number. The Kneser graph $\text{KN}(n, k)$ has vertex set $\binom{[n]}{k}$, and two vertices F_1, F_2 form an edge if and only if $F_1 \cap F_2 = \emptyset$. We use an extension of Kneser graphs to hypergraphs. Alon, Frankl, and Lovász [2] considered the Kneser hypergraph $\text{KN}^r(n, k)$, which is the r -uniform hypergraph with vertex set $\binom{[n]}{k}$, and r vertices F_1, \dots, F_r form an edge if and only if $F_i \cap F_j = \emptyset$ for $i \neq j$. They gave a lower bound on the chromatic number of $\text{KN}^r(n, k)$ as follows.

Theorem 25. *If $n \geq (t-1)(r-1) + rk$, then $\chi(\text{KN}^r(n, k)) \geq t$.*

We first show that $\text{KN}^r(n, k)$ is F_5 -free for $n < 4k$.

Lemma 26. *If $n < 4k$, then $\text{KN}^3(n, k)$ is F_5 -free.*

Proof. Say $\{a, b, c\}$, $\{a, b, d\}$ and $\{c, d, e\}$ are edges in $\text{KN}^3(n, k)$. Then by definition a, b, c , and d are four disjoint k -sets in $[n]$, which is impossible because $n < 4k$. \square

Proof of the lower bound in Theorem 7. Fix $t \geq 2$ and $\epsilon > 0$. Pick $k \geq 2t$ and $n = 3k + 2(t - 1)$ and note that $n < 4k$. By Theorem 25, $\text{KN}^3(n, k)$ has chromatic number at least t and by Lemma 26 is F_5 -free. For integers u, v , and w where n divides u , let U, V and W be disjoint vertex sets of size u, v , and w respectively. Partition U into U_1, \dots, U_n such that $|U_i| = \frac{u}{n}$ for each i . Let H be the hypergraph with vertex set $V(\text{KN}^3(n, k)) \cup U \cup V \cup W$ and the following edges.

- For $\{S_1, S_2, S_3\} \in \text{KN}^3(n, k)$, make $\{S_1, S_2, S_3\}$ an edge of H .
- For $S \in V(\text{KN}^3(n, k))$, $x \in U_i$ with $1 \leq i \leq n$, and $y \in V$, make $\{S, x, y\}$ an edge of H if $i \in S$.
- For $x \in U, y \in V$, and $z \in W$, make $\{x, y, z\}$ an edge of H .

Notice that H has chromatic number at least t because $\text{KN}^3(n, k)$ is a subhypergraph.

Claim 1: H contains no subgraph isomorphic to F_5 .

Proof. Suppose $\{a, b, c\}, \{a, b, d\}$ and $\{c, d, e\}$ are the hyperedges of a copy of F_5 in H . Notice that the hypergraph induced by $U, V, V(\text{KN}^3(n, k)) \cup W$ is 3-partite, apart from those edges within $\text{KN}^3(n, k)$. Note that a 3-uniform, 3-partite hypergraph is F_5 -free, therefore any copy of F_5 must contain an edge from $\text{KN}^3(n, k)$. If that edge is $\{a, b, c\}$ then d must also be contained in $V(\text{KN}^3(n, k))$. But then c and d are both in $V(\text{KN}^3(n, k))$, which means e must be as well. Because $\text{KN}^3(n, k)$ is F_5 -free, this is a contradiction. Similarly, $\{a, b, d\} \not\subseteq V(\text{KN}^3(n, k))$. Therefore, $\{c, d, e\} \subseteq V(\text{KN}^3(n, k))$, and without loss of generality $b \in U$ and $a \in V$. Because $\{a, b, c\}$ and $\{a, b, d\}$ are edges, b must be in both c and d , which contradicts the fact that $\{c, d, e\}$ is an edge of $\text{KN}^3(n, k)$. \square

Claim 2: The minimum degree of H is at least $(1 - \epsilon) \frac{6}{49} \binom{|V(H)|}{2}$ if $|V(H)|$ is large enough.

Proof. Vertices in $\text{KN}^3(n, k)$ have degree at least $k \frac{u}{n} v = \frac{kw}{3k+2(t-1)}$. Since t is fixed, we can choose k large enough that vertices in $\text{KN}^3(n, k)$ have degree at least $(1 - \epsilon/2)uv/3$. Vertices in A have degree at least vw , vertices in B have degree at least uw , and vertices in C have degree at least uv . Thus the minimum degree of H is at least $\min\{(1 - \epsilon/2)\frac{uv}{3}, uw, vw\}$. Choose u, v , and w so that $\frac{uv}{3} = uw = vw$, we obtain that $u = v$ and $w = v/3$ and the minimum degree is at least $(1 - \epsilon/2)u^2/3$. The number of vertices is $u + v + w + \binom{n}{k} = \frac{7}{3}u + \binom{n}{k}$. Since $u^2/3 \approx 6/49 \binom{7u/3}{2}$, we can choose u large enough so that the minimum degree of H is at least $(1 - \epsilon) \frac{6}{49} \binom{|V(H)|}{2}$. \square

We have proved that for every fixed $t \geq 2$ and every $\epsilon > 0$, there is a constant N_0 such that for $N > N_0$ there exists an N -vertex, 3-uniform, F_5 -free hypergraph with chromatic number at least t and minimum degree at least $(1 - \epsilon) \frac{6}{49} \binom{|V(H)|}{2}$. By the definition of chromatic threshold, this implies that the chromatic threshold of the family of F_5 -free hypergraphs is at least $\frac{6}{49}$. \square

8 Generalized Kneser hypergraphs

In Section 7.3, we used a generalization of the Kneser graph to hypergraphs to give a lower bound on the chromatic threshold of the family of F_5 -free hypergraphs. In Section 9, we will use similar constructions to give lower bounds on the chromatic threshold of the family of A -free hypergraphs, for several other hypergraphs A . For some of these constructions, we will need a more general variant of the Kneser hypergraph, which we explore in this section.

Sarkaria [31] considered the generalized Kneser hypergraph $\text{KN}_s^r(n, k)$, which is the r -uniform hypergraph with vertex set $\binom{[n]}{k}$, in which r vertices F_1, \dots, F_r form an edge if and only if no element of $[n]$ is contained in more than s of them. Note that the Kneser hypergraph $\text{KN}^r(n, k)$ is $\text{KN}_1^r(n, k)$. Sarkaria [31] and Ziegler [38] gave lower bounds on the chromatic number of $\text{KN}_s^r(n, k)$, but Lange and Ziegler [21] showed that the lower bounds obtained by Sarkaria and Ziegler apply only if one allow the edges of $\text{KN}_s^r(n, k)$ to have repeated vertices. We conjecture that for $\text{KN}_s^r(n, k)$, a statement similar to Theorem 25 is true.

Conjecture 27. *There exists $T(r, s, t)$ such that if $n \geq T(r, s, t) + rk/s$, then $\chi(\text{KN}_s^r(n, k)) \geq t$.*

The following much weaker statement is sufficient for our purposes. The proof is similar to an argument of Szemerédi which appears in a paper of Erdős and Simonovits [6], and the proof of Claim 1 is motivated by an argument of Kleitman [20].

Theorem 28. *Let $c > 0$; then for any integers r, t , there exists $K_0 = K_0(c, r, t)$ such that if $k \geq K_0$, $s = r - 1$, and $n = (r/s + c)k$, then $\chi(\text{KN}_s^r(n, k)) > t$.*

Before we prove this theorem, we need two definitions. A family \mathcal{F} of subsets of $[n]$ is **monotone decreasing** if $F \in \mathcal{F}$ and $F' \subseteq F$ imply $F' \in \mathcal{F}$. Similarly, it is **monotone increasing** if $F \in \mathcal{F}$ and $F \subseteq F'$ imply $F' \in \mathcal{F}$.

Proof of Lemma 28. Fix an integer t . We would like to prove that if k is large enough then it is impossible to t -color $\text{KN}_s^r(n, k)$. So let k be some integer and assume $\text{KN}_s^r(n, k)$ can be t -colored. Then the k -subsets of $[n]$ can be divided into t families, $\mathcal{F}_1, \dots, \mathcal{F}_t$, such that $F_1 \cap \dots \cap F_r \neq \emptyset$ for all distinct $F_1, \dots, F_r \in \mathcal{F}_i, 1 \leq i \leq t$. For $1 \leq i \leq t$, let $\mathcal{F}_i^* = \{A : A \subseteq [n], \exists F \in \mathcal{F}_i \text{ such that } F \subseteq A\}$. Then $\mathcal{F}_1^*, \dots, \mathcal{F}_t^*$ are monotone increasing families of subsets of $[n]$. Let $w = s/r$; since $s = r - 1$, $w = 1 - 1/r$. For a family \mathcal{F} of subsets of $[n]$, define the weighted size $W[\mathcal{F}]$ of \mathcal{F} by

$$W[\mathcal{F}] = \sum_{F \in \mathcal{F}} w^{|F|} (1 - w)^{n - |F|}.$$

Claim 1: For $1 \leq \ell \leq t$, $W[\cup_{i=1}^{\ell} \mathcal{F}_i^*] \leq 1 - 1/r^{\ell}$.

Proof. We prove this by induction on ℓ . For $\ell = 1$, Frankl and Tokushige [11] showed that for a family \mathcal{F} of subsets of $[n]$, if $F_1 \cap \dots \cap F_r \neq \emptyset$ for all distinct $F_1, \dots, F_r \in \mathcal{F}$, then $W[\mathcal{F}] \leq w = 1 - 1/r$. Now assume that the statement is true for ℓ . Let $U = \cup_{i=1}^{\ell} \mathcal{F}_i^*$ and $L = \overline{\mathcal{F}_{\ell+1}^*}$. Then $W[U] \leq 1 - 1/r^{\ell}$, U is a monotone increasing family of subsets of $[n]$, and L is a monotone decreasing family of subsets of $[n]$. By the FKG Inequality,

$$W[U \cap L] \leq W[U]W[L].$$

Then

$$\begin{aligned} W[\cup_{i=1}^{\ell+1} \mathcal{F}_i^*] &= W[U \cap L] + W[\mathcal{F}_{\ell+1}^*] \leq W[U]W[L] + W[\mathcal{F}_{\ell+1}^*] \\ &\leq (1 - 1/r^\ell)W[L] + W[\mathcal{F}_{\ell+1}^*] = 1 - (1 - W[\mathcal{F}_{\ell+1}^*])/r^\ell. \end{aligned}$$

Since $W[\mathcal{F}_{\ell+1}^*] \leq w = 1 - 1/r$, we have $1 - (1 - W[\mathcal{F}_{\ell+1}^*])/r^\ell \leq 1 - 1/r^{\ell+1}$, so $W[\cup_{i=1}^{\ell+1} \mathcal{F}_i^*] \leq 1 - 1/r^{\ell+1}$. \square

Now we know that $W[\cup_{i=1}^t \mathcal{F}_i^*] \leq 1 - 1/r^t$, so $W[\overline{\cup_{i=1}^t \mathcal{F}_i^*}] \geq 1/r^t$. We also know that $\overline{\cup_{i=1}^t \mathcal{F}_i^*}$ is the family of subsets of $[n]$ whose size is less than $k = n/(r/s + c)$, so

$$W[\overline{\cup_{i=1}^t \mathcal{F}_i^*}] = \sum_{i < \frac{n}{r/s+c}} \binom{n}{i} w^i (1-w)^{n-i}.$$

Since $wn = \frac{n}{r/s} > \frac{n}{r/s+c}$, by Chernoff's inequality we have

$$\sum_{i < \frac{n}{r/s+c}} \binom{n}{i} w^i (1-w)^{n-i} \leq e^{-\left(\frac{c}{r/s+c}\right)^2 \frac{sn}{2r}} = e^{-\frac{c^2 s}{2(r/s+c)r} k}.$$

Then if k is large and t is fixed, $W[\overline{\cup_{i=1}^t \mathcal{F}_i^*}] \leq e^{-\frac{c^2 s}{2(r/s+c)r} k} < 1/r^t$ which contradicts Claim 1. This contradiction implies that for any fixed t , there is no choice of K_0 such that for all $k > K_0$ it is possible to t -color $\text{KN}_s^r(n, k)$. This completes the proof. \square

For an r -uniform hypergraph A , we want to construct an infinite sequence of A -free hypergraphs with $\text{KN}^r(n, k)$ or $\text{KN}_{r-1}^r(n, k)$ as a subhypergraph. This will imply that these A -free hypergraphs have large chromatic number, but we must first show that for any integer k and for some choice of $n = n(k)$ one of $\text{KN}^r(n, k)$, $\text{KN}_{r-1}^r(n, k)$ is A -free. We now show that $\text{KN}_2^3(n, k)$ is T_5 -free and $S(7)$ -free under some conditions on n and k . Here T_5 is a 3-uniform hypergraph with vertices v_1, v_2, v_3, v_4, v_5 and edges $\{v_1, v_2, v_3\}$, $\{v_1, v_4, v_5\}$, $\{v_2, v_4, v_5\}$, $\{v_3, v_4, v_5\}$, and $S(7)$ denotes the Fano plane (the S stands for Steiner Triple System.)

Lemma 29. *If $n < (3/2 + 1/4)k$, then $\text{KN}_2^3(n, k)$ is T_5 -free.*

Proof. If $n < 3k/2$, then $\text{KN}_2^3(n, k)$ has no edge and of course is T_5 -free. Assume $n = (3/2 + \epsilon)k$ with $0 \leq \epsilon < 1/4$, and suppose T_5 is a subhypergraph of $\text{KN}_2^3(n, k)$. Since $\{v_1, v_4, v_5\}$, $\{v_2, v_4, v_5\}$, $\{v_3, v_4, v_5\}$ are edges of T_5 , the vertices v_1, v_2 , and v_3 all lie in $\overline{v_4 \cap v_5}$. Because $|\overline{v_4 \cap v_5}| \leq 2n - 2k = (1 + 2\epsilon)k < 3k/2$, by the pigeonhole principle, $v_1 \cap v_2 \cap v_3 \neq \emptyset$, which means $\{v_1, v_2, v_3\}$ is not an edge, a contradiction. \square

Lemma 30. *If $n < (3/2 + 1/10)k$, then $\text{KN}_2^3(n, k)$ is $S(7)$ -free.*

Proof. Just as in the proof of Lemma 29, assume $n = (3/2 + \epsilon)k$ with $0 \leq \epsilon < 1/10$ and suppose $S(7)$ is a subhypergraph of $\text{KN}_2^3(n, k)$. Let A be a vertex in a copy of $S(7)$ in $\text{KN}_2^3(n, k)$ and let $\{A, B, C\}$, $\{A, D, E\}$, $\{A, F, G\}$ be its incident edges in the copy of $S(7)$. Then $B \cap C, D \cap E, F \cap G \subseteq \overline{A}$. Since $|\overline{A}| = (1/2 + \epsilon)k$, $|B \cap C|, |D \cap E|, |F \cap G| \geq (1/2 - \epsilon)k$. Then since $3(1/2 - \epsilon) > 2(1/2 + \epsilon)$, the pigeonhole principle implies that $B \cap C \cap D \cap E \cap F \cap G \neq \emptyset$. Now the copy of $S(7)$ cannot have an edge not containing A , a contradiction. \square




Figure 3: Assorted Hypergraphs

9 Open Problems and Partial Results

Many open problems remain; for most 3-uniform hypergraphs A the chromatic threshold for the family of A -free hypergraphs is unknown. Interesting hypergraphs to study are those for which we know the extremal number, $ex(n, A)$, and we will examine a few of those here along with partial results and conjectures. We conjecture that most of the lower bounds given by the constructions in this section are tight.

9.1 $\mathcal{TK}^r(s)$ -free hypergraphs

For $s > r$, recall that $\mathcal{TK}^r(s)$ is the family of r -uniform hypergraphs such that there exists a set S of s vertices where each pair of vertices from S are contained together in some edge. The set S is called the set of **core vertices** of the hypergraph. Recall also that $T_{r,s}(n)$ is the complete n -vertex, r -uniform, s -partite hypergraph with part sizes as equal as possible.

The last author [25] showed that if $s > r$ then $ex(n, \mathcal{TK}^r(s)) = |T_{r,s-1}(n)|$ and $ex(n, \mathcal{TK}^r(s)) = (1 + o(1)) |T_{r,s}(n)|$. Recently, Pikhurko [27] has shown that for large n and $s > r$, $ex(n, \mathcal{TK}^r(s)) = |T_{r,s-1}(n)|$ and that $T_{r,s-1}(n)$ is the unique extremal example. Because F_5 is a member of $\mathcal{TK}^3(4)$ it follows that the chromatic threshold of $\mathcal{TK}^3(4)$ -free hypergraphs is at most $(\sqrt{41} - 5)/8$. The following simple variation on the construction from Section 7.3 provides a lower bound of $18/361$ for both $\mathcal{TK}^3(4)$ -free and $\mathcal{TK}^3(4)$ -free hypergraphs.

Proposition 31. *The chromatic threshold of $\mathcal{TK}^3(4)$ -free hypergraphs is at least $\frac{18}{361}$.*

Proof. The proof is very similar to the proof in Section 7.3, we only sketch it here. Choose k, n, u, v, w, U, V, W as in the proof of the lower bound of Theorem 7 in Section 7.3; that is k, n, u, v, w are integers and U, V, W are disjoint sets of vertices of size u, v, w respectively. Divide U into U_1, \dots, U_n so that $|U_i| = u/n$ and divide V into V_1, \dots, V_n such that $|V_i| = v/n$. Let H be the hypergraph formed by taking $\text{KN}^3(n, k)$ and adding the complete 3-partite hypergraph on U, V, W and the following edges. For $S \in V(\text{KN}^3(n, k))$ and $x \in U_i$ and $y \in V_j$, make $\{S, x, y\}$ an edge if $i, j \in S$. The minimum degree is maximized when $a = b$ and $c = a/9$, which gives minimum degree approximately $a^2 N^2 / 9 \approx \frac{18}{361} \cdot \binom{19a/9}{2} N^2$, where $N = u + v + w + \binom{n}{k}$ is the number of vertices in the hypergraphs.

Let F be any hypergraph in $\mathcal{TK}^3(4)$ and assume that F is a subhypergraph of H in which c_1, c_2, c_3, c_4 are the four core vertices. Because any 3-partite hypergraph is $\mathcal{TK}^3(4)$ -free, it is easy to see that some edge of F must lie in $\text{KN}^3(n, k)$, and so there must be at least two core

vertices in $\text{KN}^3(n, k)$. If $c_1, c_2 \in \text{KN}^3(n, k)$ and $c_3 \in U \cup V$ then c_3 is in either U_i or V_i for some i . But then $i \in c_1 \cap c_2$ (recall that vertices in $\text{KN}^3(n, k)$ are k -sets) which contradicts the fact that c_1 and c_2 are contained together in some edge of $\text{KN}^3(n, k)$. Thus all four core vertices must be in $\text{KN}^3(n, k)$, which is not possible because $n < 4k$. \square

This gives lower bounds on the chromatic thresholds of $\text{TK}^3(4)$ -free and $\mathcal{TK}^3(4)$ -free hypergraphs and leads to the following questions.

Question 32. *What is the chromatic threshold for $\text{TK}^3(4)$ -free hypergraphs? It is between $18/361$ and $2/9$. What is the chromatic threshold for $\mathcal{TK}^3(4)$ -free hypergraphs? It has the same lower bound as for $\text{TK}^3(4)$ -free hypergraphs, and because $F_5 \in \mathcal{TK}^3(4)$ the upper bound is $(\sqrt{41} - 5)/8$.*

A similar construction provides a $\mathcal{TK}^3(s)$ -free hypergraph for any $s \geq 5$. We have not optimized the values.

Lemma 33. *When $s \geq 5$, the chromatic threshold of $\mathcal{TK}^3(s)$ -free hypergraphs is at least $\frac{(s-2)(s-3)(s-4)^2}{(s^2-13)^2} = 1 - \frac{13}{s} + O(\frac{1}{s^2})$.*

Proof. Fix $t \geq 2$, $k \geq 2t$, and let $n = 3k + 2(t - 1)$. Notice that $n < 4k$. By Theorem 25, the chromatic number of $\text{KN}^3(n, k)$ is therefore at least t . Fix $N \gg \binom{n}{k}$.

Partition N vertices into one part of size u and $s - 2$ parts of size x , for some u that is divisible by n . Include as an edge each triple that has at most one vertex in each part. Further partition the part of size u into n sets, U_1, \dots, U_n , each of size u/n . From the remaining $s - 2$ parts of size x , choose two and designate them W_1, W_2 ; label the remaining $s - 4$ parts V_1, \dots, V_{s-4} . Let H be the 3-uniform hypergraph formed by taking the disjoint union of $\text{KN}^3(n, k)$ and the above complete $(s - 1)$ -partite hypergraph, and adding the following edges. If $S \in V(\text{KN}^3(n, k))$, $v \in V_i$, and $v' \in V_j$ for $i \neq j$, add the edge $\{S, v, v'\}$. If $S \in V(\text{KN}^3(n, k))$ and $u \in U_i$ and $v \in V_j$ then add the edge $\{S, u, v\}$ if and only if $i \in S$. Notice that H has chromatic number at least t , and that $V(H) = N + \binom{n}{k}$.

Claim 1: H contains no element of $\mathcal{TK}^3(s)$ as a subgraph.

Proof. Suppose there is such a subgraph; then at least one core vertex must be contained in $V(\text{KN}^3(n, k))$, because an $(s - 1)$ -partite graph is $\mathcal{TK}^s(3)$ -free. In that case, no core vertex can be in $W_1 \cup W_2$ because there is no edge that contains a vertex from $W_1 \cup W_2$ as well as a vertex from $V(\text{KN}^3(n, k))$. There must therefore be at least 3 core vertices in $V(\text{KN}^3(n, k))$, which means that two of them must appear in an edge contained within $V(\text{KN}^3(n, k))$. Suppose they are S_1, S_2 . If another core vertex is in U , say $u \in U_i$, then there must be an edge of H containing u and S_1 , and there must be an edge containing u and S_2 . This implies that $i \in S_1 \cap S_2$, which contradicts the fact that S_1 and S_2 appear together in an edge of $\text{KN}^3(n, k)$.

All core vertices must therefore be in $V(\text{KN}^3(n, k)) \cup V$, which means that there must be at least four of them in $V(\text{KN}^3(n, k))$. Because each pair of those four core vertices must appear together in an edge, and that edge must be in $\text{KN}^3(n, k)$, those four sets must be pairwise disjoint. This is impossible because $n < 4k$. \square

The minimum degree of this graph is approximately

$$\min \left\{ \frac{1}{3}(s-4)ax + \binom{s-4}{2}x^2, \binom{s-2}{2}x^2, (s-3)ax + \binom{s-3}{2}x^2 \right\}$$

Notice that a vertex in $W_1 \cup W_2$ has degree strictly less than a vertex in $\text{KN}^3(n, k)$, and so they do not enter into the above computation. This minimum is largest when $u = \frac{3(2s-7)x}{s-4}$, which implies that $x = \left(\frac{s-4}{s^2-13}\right)N$. The minimum degree of H is then

$$\frac{(s-2)(s-3)}{2} \cdot \frac{(s-4)^2}{(s^2-13)^2} N^2 = \left(1 - \frac{13}{x} + O\left(\frac{1}{s^2}\right)\right) \frac{N^2}{2}.$$

□

The construction in Lemma 33 has one part of “type” U (which is partitioned into n sets), $s-4$ parts of “type” V (which are not partitioned, and whose vertices appear in edges that intersect K), and two parts of “type” W (which are not partitioned and have no vertices that appear in edges intersecting K). Using this strategy, one can generate similar constructions for $\text{TK}^r(s)$; the above proof applies whenever there are x parts of type U , $s-(r+1)$ parts of type V , and y parts of type W , where $x+y=r$ and $s-(r+1)+x \geq r-1$. This last condition is needed for the edges intersecting K .

Question 34. *What is the chromatic threshold for $\text{TK}^3(s)$ -free hypergraphs for $s > 3$? It is between $\frac{(s-2)(s-3)(s-4)^2}{(s^2-13)^2} = 1 - \frac{13}{s} + O\left(\frac{1}{s^2}\right)$ and $\left(1 - \frac{1}{s-1}\right)\left(1 - \frac{2}{s-1}\right) = 1 - \frac{3}{s-1} + \frac{2}{(s-1)^2}$. The upper bound comes from $T_{r,s-1}(n)$.*

9.2 $S(7)$ -free hypergraphs

Next, consider the Fano plane $S(7)$. de Caen and Füredi [5] showed that $\text{ex}(n, S(7)) = \left(\frac{3}{4} + o(1)\right)\binom{n}{3}$. The extremal hypergraph for $S(7)$, proven to be extremal by Füredi and Simonovits [13] and also by Keevash and Sudakov [19], is the hypergraph formed by taking two almost equal vertex sets U and V and taking all edges which have at least one vertex in each of U and V . We can modify the hypergraph from Section 7.3 to obtain a lower bound on the chromatic threshold of $S(7)$ -free hypergraphs.

Proposition 35. *The chromatic threshold of $S(7)$ -free hypergraphs is at least $9/17$.*

Proof. Fix $t \geq 2$ and $0 < \epsilon \ll 1$. Then by Lemma 25 there exists k large enough that if $n = (3 + \epsilon)k$ then $\text{KN}^3(n, k)$ has chromatic number at least t . Fix some such k , and fix $N \gg \binom{n}{k}$.

Partition N vertices into two sets, U and V , with $|U| = 9N/17$ and $|V| = 8N/17$. Further partition U into n parts, U_1, \dots, U_n , each of size $|U|/n$. Include as an edge each triple that has at least one vertex in each of U, V . Let H be the hypergraph formed by taking the disjoint union of this hypergraph and $\text{KN}^3(n, k)$ and adding the following edges. For $u \in U_i$, $u' \in U_j$, and $X \in V(\text{KN}^3(n, k))$ include $\{X, u, u'\}$ as an edge if $i, j \in X$ (recall that vertices in $\text{KN}^3(n, k)$ are subsets of $[n]$). Let $K = V(\text{KN}^3(n, k))$. Notice that H has chromatic number at least t , and that $V(H) = N + \binom{n}{k}$.

Claim 1: H contains no subhypergraph isomorphic to $S(7)$.

Proof. First notice that $\text{KN}^3(n, k)$ is $S(7)$ -free because every pair of vertices in $S(7)$ are in an edge, which would require there to be 7 pairwise-disjoint k -subsets of $[n]$. Because $n = (3 + \epsilon)k$, this would be a contradiction. It is easy to see, by considering the partition $U, (K \cup V)$, that if H contains a copy of $S(7)$ then it must involve an edge from $H[K]$ (otherwise the extremal $S(7)$ -free hypergraph also contains a copy of $S(7)$). Call this edge $\{A, B, C\}$.

There are four vertices in $S(7) \setminus \{A, B, C\}$, and at least one must be outside K . No more than one can be in V because there is no edge with one vertex in K and two in V . No more than one can be in U otherwise one of $A \cap B, A \cap C, B \cap C$ is non-empty, which contradicts the assumption that $\{A, B, C\}$ is an edge of $H[K]$. Therefore, there must be either 5 or 6 vertices of $S(7)$ in K . Suppose v is a vertex of $S(7)$ that is outside of K . Then v appears in three edges that overlap only at v , say $\{v, S_1, S_2\}$, $\{v, S_3, S_4\}$, and $\{v, S_5, S_6\}$. At least one of these edges must contain two vertices from K , but there is no such edge in H . \square

The minimum degree of H is at least

$$\min \left\{ |U||V| + \binom{|U|/3}{2}, |U||V| + \binom{|U|}{2}, |U||V| + \binom{|V|}{2} \right\} = \frac{9}{34}N^2 - \frac{3}{34}N.$$

\square

Question 36. *What is the chromatic threshold of $S(7)$ -free hypergraphs? It is at least $9/17$ and at most $3/4$, where the upper bound is from the extremal hypergraph of $S(7)$.*

9.3 T_5 -free hypergraphs

Recall that the 3-uniform hypergraph T_5 has vertices A, B, C, D, E and edges $\{A, B, C\}$, $\{A, D, E\}$, $\{B, D, E\}$, and $\{C, D, E\}$.

Let $B^3(n)$ be the 3-uniform hypergraph with the most edges among all n -vertex 3-graphs whose vertex set can be partitioned into X_1, X_2 such that each edge contains exactly one vertex from X_2 . Füredi, Pikhurko, and Simonovits [12] proved that for n sufficiently large the extremal T_5 -free hypergraph is $B^3(n)$. It follows that the chromatic threshold for the family of T_5 -free hypergraphs is at most $4/9$.

Proposition 37. *The chromatic threshold of T_5 -free hypergraphs is at least $16/49$.*

Proof. Fix $t \geq 2$ and $0 < \epsilon \ll 1$. Then by Lemma 25 there exists k large enough that if $n = (3/2 + \epsilon)k$ then $\text{KN}_2^3(n, k)$ has chromatic number at least t . Fix some such k , and fix $N \gg \binom{n}{k}$.

Partition N vertices into two parts, U and V , with $|U| = 4N/7$ and $|V| = 3N/7$. Further partition U into n parts, U_1, \dots, U_n , each of size $|U|/n$. Include as an edge any triple with two vertices in U and one in V . Let H be the hypergraph formed by taking the disjoint union of this graph and $\text{KN}_2^3(n, k)$ and including the following edges. If $X \in V(\text{KN}_2^3(n, k))$ and $u \in U_i$ and $v \in V$ then let $\{u, v, X\}$ be an edge if $i \in X$ (recall that vertices of $\text{KN}_2^3(n, k)$ are subsets of $[n]$). Let $K = V(\text{KN}_2^3(n, k))$. Notice that H has chromatic number at least t , and that $V(H) = N + \binom{n}{k}$.

Claim 1: T_5 is not a subhypergraph of H .

Proof. Let H' be the hypergraph obtained from H by deleting all edges contained in K , and let $X_1 = K \cup U$ and $X_2 = V$. It is now easy to see that H' is a subhypergraph of the extremal T_5 -free hypergraph; if H contains a copy of T_5 it must therefore involve an edge from K . If that edge is $\{A, D, E\}$ (see the labelling of T_5 above) then because $\{B, D, E\}$ and $\{C, D, E\}$ are edges of T_5 it must be the case that both of B, C are in K , but by Lemma 29 K does not span a copy of T_5 . Similarly, neither $\{B, D, E\}$ nor $\{C, D, E\}$ can be contained in K .

We may therefore assume that $\{A, B, C\}$ is contained in K . Because $\{A, D, E\}$ is an edge, and by Lemma 29, at least one of D, E is in U . Suppose that $D \in U_i$; then because $\{A, D, E\}$, $\{B, D, E\}$, and $\{C, D, E\}$ are all edges of T_5 it must be the case that $i \in A \cap B \cap C$. This contradicts the assumption that $\{A, B, C\}$ is an edge. \square

The minimum degree of H is at least

$$\min \left\{ \frac{2|U||V|}{3}, |U||V|, \binom{|U|}{2} \right\} = \frac{8}{49}N^2 - \frac{2}{7}N.$$

\square

9.4 Co-chromatic thresholds

There is another possibility when generalizing the definition of chromatic threshold from graphs to hypergraphs: we can use the co-degree instead of the degree. Recall that if H is an r -uniform hypergraph and $\{x_1, \dots, x_{r-1}\} \subseteq V(H)$, then the **co-degree** $d(x_1, \dots, x_{r-1})$ of x_1, \dots, x_{r-1} is $|\{z : \{x_1, \dots, x_{r-1}, z\} \in H\}|$. Let F be a family of r -uniform hypergraphs. The **co-chromatic threshold** of F is the infimum of the values $c \geq 0$ such that the subfamily of F consisting of hypergraphs H with minimum co-degree at least $c|V(H)|$ has bounded chromatic number. More generally, the **k -degree** $d(x_1, \dots, x_k)$ of x_1, \dots, x_k is $|\{\{z_{k+1}, \dots, z_r\} : \{x_1, \dots, x_k, z_{k+1}, \dots, z_r\} \in H\}|$ and we can define the k -chromatic threshold similarly. Given a hypergraph H and subsets U, V, W of $V(H)$, we say that an edge $\{u, v, w\}$ is of type UVW if $u \in U, v \in V$ and $w \in W$.

The co-chromatic thresholds of F_5 -free hypergraphs and $\text{TK}^3(4)$ -free hypergraphs are trivially zero because if the minimum co-degree of H is at least 10 then H contains a copy of $\text{TK}^3(4)$ and a copy of F_5 . For the Fano plane, the last author proved [24] that for every $\epsilon > 0$ there exists n_0 such that any 3-uniform hypergraph with $n > n_0$ vertices and minimum co-degree greater than $(1/2 + \epsilon)n$ contains a copy of $S(7)$. In 2009, Keevash [17] improved this by proving that any 3-uniform hypergraph with minimum co-degree greater than $n/2$ contains a copy of $S(7)$ for n sufficiently large. Notice that the lower bound construction for the chromatic threshold described above has non-zero minimum co-degree but the co-degree depends on the parameter t . We can modify the construction to prove a better lower bound on the co-chromatic threshold of $S(7)$ -free hypergraphs.

Proposition 38. *The co-chromatic threshold of $S(7)$ -free hypergraphs is at least $2/5$.*

Proof. Fix $t \geq 2$ and $0 < \epsilon \ll 1$. Then by Lemma 28 there exists k large enough that if $n = (3/2 + \epsilon)k$ then $\text{KN}_2^3(n, k)$ has chromatic number at least t . Fix $N \gg \binom{n}{k}$.

Partition N vertices into two parts, U and V , of size $\frac{3N}{5}$ and $\frac{2N}{5}$ respectively. Include as an edge any triple with at least one vertex in each part. Further partition U into n sets,

U_1, \dots, U_n , each of size $|U|/n$. Let H be the hypergraph formed by taking the disjoint union of this hypergraph with $\text{KN}_2^3(n, k)$ and including the following edges. Include any edge of type KUV , where $K = V(\text{KN}_2^3(n, k))$. For any $X, Y \in K$, if $|X \cap Y| < k - 4\epsilon k$ then include every edge of the form $\{X, Y, u\}$ where $u \in U_i$ for some $i \in X \cup Y$. If $|X \cap Y| \geq k - 4\epsilon k$ then include every edge of the form $\{X, Y, u\}$ where $u \in U_i$ for some $i \in X \cap Y$. Notice that H has chromatic number at least t and that $V(H) = N + \binom{n}{k}$.

Claim 1: The above hypergraph contains no subgraph isomorphic to $S(7)$.

Proof. First notice that the complete bipartite 3-uniform hypergraph contains no copy of $S(7)$. Therefore, by considering the partition $U, V \cup K$, we can see that any copy of $S(7)$ must contain an edge induced by K . Call this edge $\{A, B, C\}$. It also follows from Lemma 30 that there is no copy of $S(7)$ completely contained in K .

Claim 1a: Any copy of $S(7)$ intersects U (or V) in at most one vertex.

Proof. Notice that for any edge e in $S(7)$, every other edge intersects e in at exactly one vertex; therefore for any copy of $S(7)$ in H every edge contains one of A, B, C . If there were two vertices of $S(7)$ in U (or in V) then the edge of $S(7)$ joining them would be unable to intersect A, B , or C . \square

Claim 1b: Any copy of $S(7)$ contains no vertex from V .

Proof. Suppose for contradiction a copy of $S(7)$ contains some vertex from V ; then by Claim 1a it intersects V in exactly one vertex. Every vertex of $S(7)$ is contained in three edges, but because there is at most one vertex from U involved in the copy of $S(7)$ there can be only one edge that contains the vertex from V . \square

Any copy of $S(7)$ must therefore have exactly six vertices in K and exactly one vertex in U . Suppose they are $A, B, C, D, E, F \in K$ and $G \in U_i$. Suppose also that the edges of $S(7)$ induced by K are

$$\{A, B, C\}, \{A, E, F\}, \{C, D, E\}, \{B, D, F\}.$$

Claim 1c: If $\{S_1, S_2, S_3\}$ is an edge in K then $|S_i \cap S_j| \leq k/2 + \epsilon k$ for all $i \neq j$.

Proof. This follows from the definition of the hypergraph on K :

$$k = |S_1| \leq n - |S_2 \cap S_3| = (3/2 + \epsilon)k - |S_2 \cap S_3|, \text{ so } |S_2 \cap S_3| \leq k/2 + \epsilon k,$$

and the claim follows through symmetry. \square

Claim 1d: The following intersections all have size at least $2k - 4\epsilon k$: $A \cap D, B \cap E, C \cap F$.

Proof. We will prove that $|A \cap D| \geq 2k - 4\epsilon k$; the rest follow through symmetry. Because $\{B, D, F\}$ is an edge, $D \subseteq (\overline{B} \cap F) \cup (B \cap \overline{F}) \cup (\overline{B} \cap \overline{F})$. Also, because $\{A, B, C\}$ is an edge, $|\overline{A} \cap \overline{B}| = |\overline{A}| - |\overline{A} \cap B| \leq (k/2 + \epsilon k) - (k/2 - \epsilon k) = 2\epsilon k$. Similarly, because $\{A, E, F\}$ is an edge, $|\overline{A} \cap \overline{F}| \leq 2\epsilon k$. Therefore,

$$|D \cap \overline{A}| \leq |\overline{A} \cap \overline{B} \cap F| + |\overline{A} \cap B \cap \overline{F}| + |\overline{A} \cap \overline{B} \cap \overline{F}| \leq |\overline{A} \cap \overline{B}| + |\overline{A} \cap \overline{F}| \leq 4\epsilon k,$$

and so $|D \cap A| \geq |D| - 4\epsilon k = k - 4\epsilon k$. \square

It follows from Claim 1d that $S(7)$ cannot be a subgraph of H . Otherwise, the edges $\{A, D, u\}, \{B, E, u\}, \{C, F, u\}$ would all appear, and by the definition of H , because the intersections mentioned in Claim 1d are large, it follows that $i \in (A \cap D) \cap (B \cap E) \cap (C \cap F)$. In that case, however, $A \cap B \cap C$ is not empty and so $\{A, B, C\}$ is not an edge. \square

It remains only to compute the minimum degree of H . Vertices $S_1, S_2 \in K$ have co-degree at least $\frac{k-4\epsilon k}{n}|U|$ if $|S_1 \cap S_2| \geq k - 4\epsilon k$ and at least $\frac{k+4\epsilon k}{n}|U|$ otherwise. Vertices $u_1, u_2 \in U$ have co-degree at least $|V|$ and vertices $v_1, v_2 \in V$ have co-degree at least $|U|$. All other pairs of vertices have co-degree at least $|U|$ or $|V|$. The minimum co-degree is therefore at least

$$\min \left\{ \frac{k(1-4\epsilon)}{k(3/2+\epsilon)}|U|, |U|, |V| \right\} = \left\{ \frac{2-8\epsilon}{3+2\epsilon} \cdot \frac{3}{5}N, \frac{3}{5}N, \frac{2}{5}N \right\}.$$

For some choice of ϵ , this is approximately $\frac{2}{5}|V(H)|$. \square

Question 39. *What is the co-chromatic threshold of the Fano-free hypergraphs? It is between $2/5$ and $1/2$.*

In [1] it was proved that if a family \mathcal{F} of graphs has positive chromatic threshold then the chromatic threshold of \mathcal{F} is in fact at least $1/3$. We think that a similar statement holds for hypergraphs. For 3-uniform hypergraphs, we believe that the least positive chromatic threshold is achieved by the family of $\text{TK}^3(4)$ -free hypergraphs (see Section 9.1).

Conjecture 40. *If a family \mathcal{F} of 3-uniform hypergraphs has positive chromatic threshold then the chromatic threshold of \mathcal{F} is at least $18/361$.*

References

- [1] P. Allen, J. Böttcher, S. Griffiths, Y. Kohayakawa, and R. Morris. Chromatic thresholds of graphs. Manuscript.
- [2] N. Alon, P. Frankl, and L. Lovász. The chromatic number of Kneser hypergraphs. *Trans. Amer. Math. Soc.*, 298:359–370, 1986.
- [3] B. Andrásfai, P. Erdős, and V. T. Sós. On the connection between chromatic number, maximal clique and minimal degree of a graph. *Discrete Math.*, 8:205–218, 1974.
- [4] S. Brandt and S. Thomassé. Dense triangle-free graphs are four colorable: A solution to the Erdős-Simonovits problem. to appear in *J. Combin. Theory Ser. B*.
- [5] D. De Caen and Z. Füredi. The maximum size of 3-uniform hypergraphs not containing a Fano plane. *J. Combin. Theory Ser. B*, 78:274–276, 2000.
- [6] P. Erdős and M. Simonovits. On a valence problem in extremal graph theory. *Discrete Math.*, 5:323–334, 1973.
- [7] P. Erdős and M. Simonovits. Supersaturated graphs and hypergraphs. *Combinatorica*, 3:181–192, 1983.

- [8] P. Erdős and A. H. Stone. On the structure of linear graphs. *Bull. Amer. Math. Soc.*, 52:1087–1091, 1946.
- [9] P. Frankl and Z. Füredi. A new generalization of the Erdős-Ko-Rado theorem. *Combinatorica*, 3(3-4):341–349, 1983.
- [10] P. Frankl and Z. Füredi. Extremal problems whose solutions are the blowups of the small Witt-designs. *J. Combin. Theory Ser. A*, 52(1):129–147, 1989.
- [11] P. Frankl and N. Tokushige. Weighted multiply intersecting families. *Studia Sci. Math. Hungar.*, 40(3):287–291, 2003.
- [12] Z. Füredi, O. Pikhurko, and M. Simonovits. On triple systems with independent neighbourhoods. *Combin. Probab. Comput.*, 14(5-6):795–813, 2005.
- [13] Z. Füredi and M. Simonovits. Triple systems not containing a Fano configuration. *Combin. Probab. Comput.*, 14:467–484, 2005.
- [14] W. Goddard and J. Lyle. Dense graphs with small clique number. *Journal of Graph Theory*, 2010.
- [15] W. T. Gowers. Hypergraph regularity and the multidimensional Szemerédi theorem. *Ann. of Math. (2)*, 166:897–946, 2007.
- [16] P. Keevash. Hypergraph Turán problems. to appear in *Surveys in Combinatorics 2011*.
- [17] P. Keevash. A hypergraph regularity method for generalized Turán problems. *Random Structures Algorithms*, 34(1):123–164, 2009.
- [18] P. Keevash and D. Mubayi. Stability theorems for cancellative hypergraphs. *J. Combin. Theory Ser. B*, 92(1):163–175, 2004.
- [19] P. Keevash and B. Sudakov. The Turán number of the Fano plane. *Combinatorica*, 25:561–574, 2005.
- [20] D. J. Kleitman. Families of non-disjoint subsets. *J. Combinatorial Theory*, 1:153–155, 1966.
- [21] C. E. M. C. Lange and G. M. Ziegler. On generalized Kneser hypergraph colorings. *J. Combin. Theory Ser. A*, 114(1):159–166, 2007.
- [22] T. Łuczak. On the structure of triangle-free graphs of large minimum degree. *Combinatorica*, 26:489–493, 2006.
- [23] T. Łuczak and S. Thomassé. Coloring dense graphs via VC-dimension. submitted.
- [24] D. Mubayi. The co-degree density of the Fano plane. *J. Combin. Theory Ser. B*, 95(2):333–337, 2005.
- [25] D. Mubayi. A hypergraph extension of Turán’s theorem. *J. Combin. Theory Ser. B*, 96:122–134, 2006.

- [26] B. Nagle, V. Rödl, and M. Schacht. The counting lemma for regular k -uniform hypergraphs. *Random Structures Algorithms*, 28:113–179, 2006.
- [27] O. Pikhurko. Exact computation of the hypergraph Turán function for expanded complete 2-graphs. accepted in *Journal of Combinatorial Theory. Series B*.
- [28] O. Pikhurko. An exact Turán result for the generalized triangle. *Combinatorica*, 28(2):187–208, 2008.
- [29] V. Rödl and J. Skokan. Regularity lemma for k -uniform hypergraphs. *Random Structures Algorithms*, 25:1–42, 2004.
- [30] V. Rödl and J. Skokan. Applications of the regularity lemma for uniform hypergraphs. *Random Structures Algorithms*, 28(2):180–194, 2006.
- [31] K. S. Sarkaria. A generalized Kneser conjecture. *J. Combin. Theory Ser. B*, 49(2):236–240, 1990.
- [32] N. Sauer. On the density of families of sets. *J. Combinatorial Theory Ser. A*, 13:145–147, 1972.
- [33] M. Simonovits. Extremal graph problems with symmetrical extremal graphs, additional chromatic conditions. *Discrete Math*, 7:349–376, 1974.
- [34] T. Tao. A variant of the hypergraph removal lemma. *J. Combin. Theory Ser. A*, 113:1257–1280, 2006.
- [35] C. Thomassen. On the chromatic number of triangle-free graphs of large minimum degree. *Combinatorica*, 22:591–596, 2002.
- [36] C. Thomassen. On the chromatic number of pentagon-free graphs of large minimum degree. *Combinatorica*, 27:241–243, 2007.
- [37] V. N. Vapnik and A. Y. Chervonenkis. Theory of uniform convergence of frequencies of events to their probabilities and problems of search for an optimal solution from empirical data. *Avtomat. i Telemekh.*, pages 42–53, 1971.
- [38] G. M. Ziegler. Generalized Kneser coloring theorems with combinatorial proofs. *Invent. Math.*, 147(3):671–691, 2002.