

ON KREIN-ŠMULIAN THEOREM FOR WEAKER TOPOLOGIES

B. CASCALES AND R. SHVYDKOY

ABSTRACT. We investigate possible extensions of the classical Krein-Šmulian theorem to various weak topologies. In particular, if X is a WCG Banach space and τ is any locally convex topology weaker than the norm-topology, then for every τ -compact norm-bounded set H , $\overline{\text{conv}}^\tau H$ is τ -compact. In arbitrary Banach spaces, the norm-fragmentability assumption on H is shown to be sufficient for the last property to hold.

A new proof to the following result is given: if a Banach space does not contain a copy of $\ell_1[0, 1]$, then the Krein-Šmulian theorem holds for every topology τ induced by a norming set of functionals. We conclude that in such spaces a norm-bounded set is weakly compact if it is merely compact in topology induced by a boundary. On the other hand, the same statement is obtained for all $C(K)$ and $\ell_1(\Gamma)$ spaces.

1. INTRODUCTION

A well-known result that goes back to M. Krein and V. Šmulian [23] says the following: the closed convex hull of a weakly compact subset of a Banach space X is weakly compact. It is known that his result also holds when the weak topology in X is replaced by any locally convex topology compatible with the dual pair $\langle X, X^* \rangle$, [17, Corollary 9.9.6]. For what other topologies does this statement remain true?

Recent attention to this question is motivated by its connection with the Boundary Problem posed by G. Godefroy in [14].

Let X be a Banach space and B a *boundary* in the unit ball of X^* , i.e such that $\|x\| = \max_{b \in B} b(x)$ holds for all $x \in X$. Denote by $\sigma(X, B)$ the topology in X of pointwise convergence on B . Is a norm-bounded subset H of X weakly compact if it is merely compact with respect to $\sigma(X, B)$?

Date: May 1, 2002.

1991 Mathematics Subject Classification. Primary: 46A50; Secondary: 46B50.

Key words and phrases. Krein-Šmulian theorem, Riemann-Lebesgue integral sum, limit set, boundary, fragmentability.

The first named author was partially supported by the research grant DGES PB 98-0381.

The work of the second named author was partially supported by the student part of NSF grant DMS-9800027.

In [30] S. Simons gives a partial positive answer to this question in the case in which H is a convex set. This establishes the equivalence between the Krein-Šmulian-type theorem for topologies $\sigma(X, B)$ generated by boundaries and the Boundary Problem itself. In other words, if one can prove that in a certain Banach space the $\sigma(X, B)$ -closed convex hull of every norm-bounded $\sigma(X, B)$ -compact subset is again $\sigma(X, B)$ -compact, *i.e.* the analogue of the classical Krein-Šmulian Theorem, then following Simons' result for $\sigma(X, B)$, the Boundary Problem is solved positively in that given Banach space.

Even though the problem remains still open, to the best of our knowledge, a considerable progress has been made by B. Cascales, G. Godefroy, G. Vera and others in a series of papers [3, 4, 5, 6, 8, 9]. In particular, the Boundary Problem has been positively solved for all boundaries in spaces of continuous functions defined on a compact space, [4], and for the particular boundary given by the set of extreme points (in the dual unit ball) for general Banach spaces, [3]. The positive solution was also found for all Banach spaces not containing a copy of $\ell_1[0, 1]$ in [5]. In fact a more general statement was proved.

Theorem 1.1 ([5]). *Suppose X does not contain a copy of $\ell_1[0, 1]$ and B is a norming subset of the unit ball of X^* . Then the $\sigma(X, B)$ -closed convex hull of every $\sigma(X, B)$ -compact norm-bounded set in X is $\sigma(X, B)$ -compact.*

Here and further on, by norming set (also known as 1-norming set) for the Banach space $(X, \|\cdot\|)$ we mean a set $B \subset B_{X^*}$ such that $\|x\| = \sup_{b \in B} b(x)$ for all $x \in X$. For example, any boundary is a norming set.

So, Theorem 1.1 combined with the aforementioned Simons' result solves the Boundary Problem, in particular, for all separable, reflexive and, more generally, all weakly compactly generated (WCG for short) or weakly Lindelöf spaces Banach spaces [16].

In the first part of our paper we recall that in order for a compact set H to have compact closed convex hull, every Radon measure on H must possess a barycenter, and vice versa. This last condition is proved to follow from so-called Riemann-Lebesgue integrability of the identity mapping on H . Based on recent results by V. Kadets, et. al. (see [19, 20, 29]) we immediately obtain the Krein-Šmulian theorem for all topologies weaker than the norm topology of a given Banach space X , provided X is either WCG or X has an unconditional basis (possibly not countable) and fails to contain a copy of $\ell_1(\Gamma)$ over any uncountable set Γ . Furthermore, in Theorem 2.4 we obtain the same result for all compact sets fragmentable by the norm. This, in particular, generalizes an earlier result in [8].

In Section 3 we continue the discussion of the Krein-Šmulian theorem and give an alternative geometrical proof of quoted Theorem 1.1. Our approach is based upon a straightforward construction of a sequence of independent functions (much in the spirit of [28]) whenever the conclusion of the theorem

is violated. This subsequently allows us to embed a copy of $\ell_1[0, 1]$ into the space. Our argument is self-contained and does not employ the non-trivial results used in the original proof in [5].

We further observe that in spite of Theorem 1.1, the Boundary Problem itself has positive solution in any $\ell_1(\Gamma)$. This phenomenon is treated in Section 4. We will find that all $\ell_1(\Gamma)$ and all $C(K)$ -spaces are *angelic* in any topology generated by a boundary. This condition is shown to imply a positive solution the Boundary Problem in Proposition 4.3.

Our notation and terminology are standard. We borrow some standard topological results from books [13, 17, 21, 22, 27]. Our vector spaces are all real. If X is a Banach space, $B(X)$ denotes its closed unit ball, and X^* its topological dual space. For a locally convex space (X, τ) endowed with topology τ its dual is denoted, as usual, by $(X, \tau)^*$. Whenever B is a subset of $(X, \tau)^*$, we write $\sigma(X, B)$ to denote the locally convex topology of convergence on functionals from B . Also we adopt the following short notation for weak topologies: $\sigma(X, X^*)$ in the usual Banach space sense is denoted by 'w' or 'w(τ)' for a general locally convex space with topology τ . Analogously, $\sigma(X^*, X) = w^*$ or $w^*(\tau)$.

The authors are very grateful to the referee who made numerous remarks and suggestions, which substantially improved the text.

2. THE KREIN-ŠMULIAN THEOREM AND BARYCENTERS

The study of compact convex sets is closely related to the existence of barycenters, see for example [10, 11, 26]. If H is a compact subset of the locally convex space (X, τ) we denote by $\mathcal{P}(H)$ the set of all Radon probabilities μ defined on the σ -algebra $\mathcal{B}(H)$ of τ -Borel subsets of H . A barycenter of μ is said to be a vector $x \in X$ such that the equality

$$(1) \quad x^*(x) = \int_H x^*(h) d\mu(h),$$

holds for every $x^* \in (X, \tau)^*$. Observe that the right hand side of equation (1) is well-defined, because $x^*|_H$ is τ -continuous and bounded, hence μ -integrable.

In general, a barycenter may not exist, however its uniqueness follows immediately from the fact that $(X, \tau)^*$ separates the points of X . Let us denote by x_μ the barycenter of $\mu \in \mathcal{P}(H)$ whenever it exists. It is well known that

$$(2) \quad \overline{\text{conv}}^\tau H = \{x_\mu : \mu \in \mathcal{P}(H), \mu \text{ has a barycenter}\},$$

see [26, Proposition 1.2] or [11, Theorem 2, p. 149].

The following lemma exhibits the classical link between barycenters and the Krein-Šmulian theorem.

Lemma 2.1. *Let H be a τ -compact set in a locally convex space (X, τ) . Then $\overline{\text{conv}}^\tau H$ is τ -compact if and only if every measure $\mu \in \mathcal{P}(H)$ has a barycenter in X .*

Proof. If $\overline{\text{conv}}^\tau H$ is τ -compact, then every $\mu \in \mathcal{P}(H)$ has a barycenter after [11, Theorem 1, p. 148].

Conversely, let us suppose that every measure $\mu \in \mathcal{P}(H)$ has a barycenter and let us see that $\overline{\text{conv}}^\tau H$ is τ -compact. Since the mapping $\varphi : \mathcal{P}(H) \rightarrow X$ defined by $\varphi(\mu) = x_\mu$ is weak*-w(τ)-continuous, we obtain that $\varphi(\mathcal{P}(H))$ is w(τ)-compact. According to (2), $\overline{\text{conv}}^\tau H$ is also w(τ)-compact. The τ -compactness of $\overline{\text{conv}}^\tau H$ (that clearly follows from the classical Krein-Smulian's theorem, [17, Corollary 9.9.6]) is recalled below for sake of completeness: since H is τ -compact, the closed convex hull $\overline{\text{conv}}^\tau H$ is precompact (τ -totally bounded). To prove the τ -compactness of $\overline{\text{conv}}^\tau H$ we show that every net in this set has a converging subnet. So, let us fix a net $\{y_\alpha\}$ in $\overline{\text{conv}}^\tau H$. The w(τ)-compactness implies the existence of a subnet $\{y_\beta\}$ of $\{y_\alpha\}$ converging to some $y \in \overline{\text{conv}}^\tau H$ in topology w(τ). In addition, the τ -total boundedness of $\overline{\text{conv}}^\tau H$ implies that there exists a further subnet $\{y_\gamma\}$ of $\{y_\beta\}$ which is τ -Cauchy. Since τ has a basis of neighborhoods of the origin consisting of w(τ)-closed sets, we conclude that actually $y = \tau - \lim_\gamma y_\gamma$ see [17, Theorem 3.2.4] and the proof is over. \square

As we will see in a moment, barycenters are related to the concept of so-called Riemann-Lebesgue integral introduced in [20]. Let us briefly outline the definition.

Suppose that X is a Banach space, (Ω, Σ, μ) is a probability space and $f: \Omega \rightarrow X$ is a norm-bounded function not necessarily measurable in any sense. Given a partition $\Pi = \{A_i\}_{i=1}^n$ of Ω into measurable sets and a collection $T = \{t_i\}_{i=1}^n$ of *sampling points*, i.e. $t_i \in A_i$, $i = \overline{1, n}$, we define the associated *Riemann-Lebesgue integral sum* as follows:

$$S(f, \Pi, T) = \sum_{i=1}^n f(t_i)\mu(A_i).$$

We endow $\{S(f, \Pi, T)\}_{\Pi, T}$ with a net structure by defining a partial order by the rule: $\Pi_1 \succ \Pi_2$ if and only if every element of Π_1 is contained in some element of Π_2 . If this net converges to some element x in the norm topology, then f is called Riemann-Lebesgue integrable, and x is then its Riemann-Lebesgue integral. We refer the reader to [2, 7, 19, 20, 29] for detailed treatment of this and related notions.

Notice that if f is strongly measurable then its Bochner integrability is equivalent to convergence of the entire net of its Riemann-Lebesgue integral sums (see [20]).

Assume now that the Banach space X is also endowed with another locally convex topology τ weaker than the norm topology. If H is a τ -compact set in X and the identity map $id: H \rightarrow X$ is Riemann-Lebesgue integrable with respect to a measure $\mu \in \mathcal{P}(H)$ then its integral is the barycenter of μ . More generally, if the net of the Riemann-Lebesgue integral sums of $id: H \rightarrow X$

has a cluster point, then this point is the barycenter of μ . Indeed, if $x = \lim_{\alpha} S(id, \Pi_{\alpha}, T_{\alpha})$ for some subnet, then for every $x^* \in (X, \tau)^*$ we have

$$x^*(x) = \lim_{\alpha} x^*(S(id, \Pi_{\alpha}, T_{\alpha})) = \lim_{\alpha} S(x^*|_H, \Pi_{\alpha}, T_{\alpha}) = \int_H x^*(h) d\mu(h),$$

since the last integral converges in the conventional Lebesgue sense.

Certain geometric conditions on the Banach space are shown to guarantee existence of a cluster point for any measure μ . Based on [19, Theorem 4.1] and [29, Theorem 2.1.2] where such conditions are formulated we immediately obtain the following result.

Theorem 2.2. *Let X be a Banach space satisfying either of the two conditions below:*

- i) X is a WCG-space;
- ii) X has an unconditional basis (possibly not countable) and fails to contain a copy of $\ell_1(\Gamma)$ over uncountable Γ .

Let also τ be a locally convex topology on X weaker than the norm-topology. Then the τ -closed convex hull of any τ -compact norm-bounded subset H of X is τ -compact.

Although the geometric assumptions on the space X in this theorem are obviously more restrictive than in Theorem 1.1, the conclusion holds for more general topologies.

Next, using Lemma 2.1 and the ideas above we isolate a class of compact sets (for topologies weaker than the weak topology) in a Banach space for which the Krein-Šmulian theorem holds. We will use the notion of fragmentability, originally introduced by Jayne and Rogers in [18], that is stated below:

Definition 2.3. Let (Z, τ) be a topological space and ρ a metric on Z . We say that (Z, τ) is *fragmented by ρ* (or *ρ -fragmented*) if for each non-empty subset C of Z and for each $\varepsilon > 0$ there exists a τ -open subset U of Z such that $U \cap C \neq \emptyset$ and $\rho - \text{diam}(U \cap C) \leq \varepsilon$.

A great variety of sufficient conditions for norm-fragmentability of a subset in a Banach space can be found in the literature: weakly compact sets of Banach spaces are norm-fragmented, [24]; more generally, sets which are Lindelöf for the weak topology and compact with respect to the topology generated by a norming set of functionals are fragmented too, [6, 8].

Theorem 2.4. *Let X be a Banach space and τ any locally convex topology in X weaker than the norm-topology. If $H \subset X$ is a τ -compact norm-bounded set fragmented by the norm, then $\overline{\text{conv}}^{\tau} H$ is τ -compact. Furthermore,*

$$\overline{\text{conv}}^{\tau} H = \overline{\text{conv}}^{\|\cdot\|} H.$$

First we show that a fragmentable set can be essentially split into subsets of small diameter.

Lemma 2.5. *Let (H, τ) be a compact space fragmented by a metric ρ and $\mu \in \mathcal{P}(H)$. Then for every $\varepsilon > 0$ there is a finite partition A_1, A_2, \dots, A_m of H in $\mathcal{B}(H)$ such that:*

- i) $\rho - \text{diam}(A_i) < \varepsilon$, $i = 1, 2, \dots, m - 1$;
- ii) $\mu(A_m) < \varepsilon$.

Proof. Let $\mathcal{A} = \{A \in \mathcal{B}(H) : \rho - \text{diam}(A) < \varepsilon\}$ and let \mathcal{F} be the family made of finite unions of elements in \mathcal{A} . The ρ -fragmentability of (H, τ) implies that \mathcal{A} is not empty; thus \mathcal{F} is not empty either. Let us define $\alpha = \sup\{\mu(B) : B \in \mathcal{F}\}$ and pick a sequence (B_n) in \mathcal{F} such that $\alpha = \lim_n \mu(B_n)$. If $E_n = \bigcup_{k=1}^n B_k$ we still have $\alpha = \lim_n \mu(E_n) = \mu(\bigcup_{n=1}^{\infty} E_n)$. We claim that

$$(3) \quad \mu \left(H \setminus \bigcup_{n=1}^{\infty} E_n \right) = 0.$$

If this is not the case, then there is a compact set $K \subset H \setminus (\bigcup_{n=1}^{\infty} E_n)$ such that $\mu(K) > 0$. The restriction $\mu|_K$ of μ to the Borel sets of K is a Radon measure that has a non empty support $F \subset K$. The ρ -fragmentability of (H, τ) applied to F implies that there is an open set $O \subset H$ such that $O \cap F \neq \emptyset$ and $\rho - \text{diam}(O \cap F) < \varepsilon$. We also have $\mu(O \cap F) > 0$ because F is the support of $\mu|_K$. Consequently,

$$\alpha \geq \lim_n \mu(E_n \cup (O \cap F)) = \lim_n \mu(E_n) + \mu(O \cap F) = \alpha + \mu(O \cap F) > \alpha$$

and we reach the contradiction that establishes the validity of (3). Since

$$\lim_n \mu(H \setminus E_n) = \mu \left(H \setminus \bigcup_{n=1}^{\infty} E_n \right) = 0$$

we can find a $m \in \mathbf{N}$ such that $\mu(H \setminus E_m) < \varepsilon$. Put $A_m = H \setminus E_m$. Then A_m satisfies ii) and clearly E_m can be split as required in i). \square

Let us point out that our lemma above is very much like an argument used in the proof of Theorem 2.3 in [25]: if we assume that ρ is lower semi-continuous with respect to τ in our lemma, then we can take A_1, A_2, \dots, A_{m-1} being compact (just adapt the first part of the proof of Theorem 2.3 in [25] to this situation).

Proof of Theorem 2.4. We show that the identity mapping $id : H \rightarrow X$ is Riemann-Lebesgue integrable with respect to any measure $\mu \in \mathcal{P}(H)$. According to Lemma 2.1 and the preceding discussion, this implies the first part of the theorem. Moreover, from (2) we conclude that $\overline{\text{con}}^\tau H$ lies in the closure of all possible Riemann-Lebesgue integral sums of id , which is obviously a subset of $\overline{\text{con}}^{\|\cdot\|} H$. This implies the second part.

So, let us fix a $\mu \in \mathcal{P}(H)$. Without loss of generality we can assume that H lies in the unit ball of X . For any given $k \in \mathbf{N}$, using Lemma 2.5 we can find a finite partition $V_1^k, V_2^k, \dots, V_{n_k}^k$ of H in $\mathcal{B}(H)$ such that

$$(4) \quad \|\cdot\| - \text{diam} V_i^k < \frac{1}{2^{k+1}}, \quad i = \overline{1, n_k - 1},$$

and

$$(5) \quad \mu(V_{n_k}^k) < \frac{1}{2^{k+1}}.$$

Let us now denote $A_{i_1 i_2 \dots i_k} = \bigcap_{j=1}^k V_{i_j}^j$, where $1 \leq i_j \leq n_j$, $1 \leq j \leq k$, and define a sequence of partitions of H as follows:

$$\Pi_k = \{A_{i_1 \dots i_k} : 1 \leq i_j \leq n_j, 1 \leq j \leq k\}.$$

For each $k \in \mathbf{N}$ we also fix an arbitrary set of sampling points $T_k = \{t_{i_1 \dots i_k} : t_{i_1 \dots i_k} \in A_{i_1 \dots i_k}\}$. We claim that the limit $\lim_{k \rightarrow \infty} S(id, \Pi_k, T_k)$ exists in the norm-topology, and is a limit point of the integral sums, even though the sequence $\{\Pi_k, T_k\}$ is not a proper subnet.

Indeed, in view of (4) and (5), we have

$$\begin{aligned} & \|S(id, \Pi_k, T_k) - S(id, \Pi_{k+1}, T_{k+1})\| \\ = & \left\| \sum_{\substack{1 \leq j \leq k \\ 1 \leq i_j \leq n_j}} t_{i_1 \dots i_k} \mu(A_{i_1 \dots i_k}) - \sum_{\substack{1 \leq j \leq k+1 \\ 1 \leq i_j \leq n_j}} t_{i_1 \dots i_k i_{k+1}} \mu(A_{i_1 \dots i_k i_{k+1}}) \right\| \\ = & \left\| \sum_{\substack{1 \leq j \leq k \\ 1 \leq i_j \leq n_j}} t_{i_1 \dots i_k} \sum_{i_{k+1}=1}^{n_{k+1}} \mu(A_{i_1 \dots i_k i_{k+1}}) - \sum_{\substack{1 \leq j \leq k+1 \\ 1 \leq i_j \leq n_j}} t_{i_1 \dots i_k i_{k+1}} \mu(A_{i_1 \dots i_k i_{k+1}}) \right\| \\ \leq & \sum_{\substack{1 \leq j \leq k+1 \\ 1 \leq i_j \leq n_j}} \|t_{i_1 \dots i_k} - t_{i_1 \dots i_k i_{k+1}}\| \mu(A_{i_1 \dots i_k i_{k+1}}) \leq \frac{3}{2^{k+1}}. \end{aligned}$$

So, the sequence $\{S(id, \Pi_k, T_k)\}$ converges to some vector $x \in X$. An easy computation also gives the estimate

$$\|S(id, \Pi_k, T_k) - x\| \leq \frac{3}{2^k}, \quad k = 1, 2, \dots$$

Given $\varepsilon > 0$ take $k \in \mathbf{N}$ so that $\frac{9}{2^{k+1}} < \varepsilon$. If $\Pi \succ \Pi_k$ and T is any collection of sampling points in Π , the same calculations as above show that

$$\|S(id, \Pi, T) - S(id, \Pi_k, T_k)\| \leq \frac{3}{2^{k+1}},$$

and hence

$$\|S(id, \Pi, T) - x\| \leq \frac{3}{2^{k+1}} + \frac{3}{2^k} = \frac{9}{2^{k+1}} < \varepsilon.$$

This proves the desired result and finishes the argument. \square

Let us note that Theorem 2.4 applied to the spaces of Bochner integrable functions considered in [8, Example E] yields the main results of [1] as a consequence. Furthermore, if X is an Asplund space (i.e. X^* has the Radon-Nikodým property, or equivalently the w^* -compact subsets of X^* are norm-fragmented), then according to our theorem, for every w^* -compact subset H of X^* we have the equality $\overline{\text{conv}}^{w^*} H = \overline{\text{conv}}^{\|\cdot\|} H$, which gives an alternative proof of [24, Theorem 2.3].

We conclude this section with several remarks.

First we comment on the fact that Lemma 2.5 implies strong μ -measurability of id . Hence, id is Bochner integrable and its Riemann-Lebesgue integral x that we found at the end of the proof of Theorem 2.4 is in fact its Bochner integral too.

We also remark that τ -compact sets as in Theorem 2.4 is not automatically norm-bounded even if τ is generated by a norming set of functionals. Indeed, consider $X = \ell_1$ and τ induced by the coordinate-axis vectors $\{e_n\}_{n \in \mathbb{N}} \subset \ell_\infty$. Set $H = \{ne_n\}_{n \in \mathbb{N}} \cup \{0\} \subset \ell_1$. Then H is unbounded, yet τ -compact.

3. A NEW PROOF OF THEOREM 1.1

In this section we give an alternative proof of Theorem 1.1 stated in the introduction. Our approach is based on a geometric construction of a independent sequence of functions on a τ -compact ($\tau = \sigma(X, B)$) set with non-compact convex hull. After a short argument, presented in the original proof in [5], this implies existence of a copy of $\ell_1[0, 1]$ in X .

So, for the rest of this section we assume that there exists a norm-bounded τ -compact set H in X such that $\overline{\text{conv}}^\tau H$ is not τ -compact, and we show that X then contains a copy of $\ell_1[0, 1]$. For purely technical reasons we also assume without loss of generality that H is contained in the unit ball of X and that the norming set B inducing τ is absolutely convex.

In view of Lemma 2.1, there is a measure $\mu \in \mathcal{P}(H)$ without a barycenter. We can decompose μ into the sum of its purely atomic part μ_a and its atomless part. The purely atomic part always has a barycenter. Indeed, in order to see it, we recall that the Radon probability μ has at most countably many disjoint atoms that are singletons $(h_i)_i$. Hence, $\mu_a = \sum_i \lambda_i \delta_{h_i}$, with $\lambda_i \geq 0$ and $\sum_i \lambda_i \leq 1$, and thus $x_{\mu_a} = \sum_i \lambda_i h_i$ is the barycenter for μ_a . This observation implies that only the atomless part of μ does not have a barycenter.

So, from now on we assume that μ has no atoms. Besides, we can identify H with the support of μ , so every open set in H has positive measure with respect to μ .

Our plan is to pick a sequence of functionals $(f_n)_{n \in \mathbf{N}}$ in B so that

$$(6) \quad H \cap \left(\bigcap_{m \in M} \{f_m > r + \delta\} \right) \cap \left(\bigcap_{n \in N} \{f_n < r\} \right) \neq \emptyset$$

holds for every two disjoint sets of natural numbers M and N , and some fixed two real numbers r and δ , $\delta > 0$. Such a sequence is called independent over H (see [28]). Every Banach space, which contains an independent sequence over a compact set, also contains a copy of $\ell_1[0, 1]$ (see Lemma B in [5]).

Our construction is based on the following lemmas.

Lemma 3.1. *There exists an $\varepsilon > 0$ and a Borel set A in H with $\mu(A) > 0$, such that for every Borel subset B in A with $\mu(B) > 0$, and every $h \in \overline{\text{conv}}^\tau H$ there is an $f \in B$ satisfying the following inequality:*

$$(7) \quad f(h) > \varepsilon + \frac{1}{\mu(B)} \int_B f(s) d\mu(s).$$

Proof. Suppose, on the contrary, that for any $\varepsilon > 0$ and measurable $A \subset H$ there is a $B \subset A$ and $h \in \overline{\text{conv}}^\tau H$ such that

$$f(h) \leq \varepsilon + \frac{1}{\mu(B)} \int_B f(s) d\mu(s),$$

whenever $f \in B$.

Let $\varepsilon_n = \frac{1}{2^n}$, $n \in \mathbf{N}$. By the exhaustion argument, using the previous inequality for $\varepsilon_1 = \frac{1}{2}$, we can find a sequence $(h_n^1)_{n \in \mathbf{N}} \subset \overline{\text{conv}}^\tau H$ and a pairwise disjoint sequence $(A_n^1)_{n \in \mathbf{N}}$ in $\mathcal{B}(H)$ such that $\mu(H \setminus \bigcup_{n=1}^\infty A_n^1) = 0$ and

$$f(h_n^1) \leq \varepsilon_1 + \frac{1}{\mu(A_n^1)} \int_{A_n^1} f(s) d\mu(s),$$

for all $f \in B$ and $n \in \mathbf{N}$. Hence, as B is absolutely convex, we have

$$\left| f(h_n^1) - \frac{1}{\mu(A_n^1)} \int_{A_n^1} f(s) d\mu(s) \right| \leq \varepsilon_1,$$

$f \in B$, $n \in \mathbf{N}$. Letting $h^1 = \sum_{n=1}^\infty \mu(A_n^1) h_n^1$ and adding up the previous inequalities we get

$$\left| f(h^1) - \int_H f(s) d\mu(s) \right| \leq \varepsilon_1.$$

In the same manner, for every $n \in \mathbf{N}$, we can construct an $h^n \in \overline{\text{conv}}^\tau H$ so that

$$\left| f(h^n) - \int_H f(s) d\mu(s) \right| \leq \varepsilon_n,$$

for all $f \in B$. Since B is norming, it follows that $\|h^n - h^{n+1}\| \leq \varepsilon_n + \varepsilon_{n+1}$ and hence, the limit $h = \|\cdot\| - \lim_{n \rightarrow \infty} h_n$ exists. Passing to limits in the previous inequality we see that h is the barycenter of μ , which contradicts our assumption. \square

Remark that since μ is a regular measure, A can be chosen closed. Furthermore, restricting μ on A we can and do assume that A is in fact the whole H .

We say that a set $K \subset X$ has a finite ε -net if there is a finite subset F of K such that $K \subset \bigcup_{x \in F} \{y \in X : \|y - x\| \leq \varepsilon\}$. It is a basic fact, that every norm-compact set has a finite ε -net for all $\varepsilon > 0$.

From now on, we fix the $\varepsilon > 0$ found in Lemma 3.1.

Lemma 3.2. *For any norm-compact set $K \subset \overline{\text{conv}}^\tau H$, any collection of open sets $(U_i)_{i=1}^n$ in H and positive numbers $(\lambda_i)_{i=1}^n$, $\sum_{i=1}^n \lambda_i = 1$, there are open sets $(V_i)_{i=1}^n$ satisfying the following conditions:*

- i) $V_i \subset U_i$, $i = \overline{1, n}$;
- ii) $\text{dist}(K, \sum_{i=1}^n \lambda_i v_i) > \frac{\varepsilon}{2}$, whenever $v_i \in V_i$, $i = \overline{1, n}$.

Proof. First we find a Borel subset W_i in every U_i so that $\mu(W_i) = \lambda_i \mu(W) > 0$, where $W = \bigcup_{i=1}^n W_i$ and $W_i \cap W_j = \emptyset$, $i \neq j$. Indeed, since μ is atomless we can pick disjoint Borel sets $A_i \subset U_i$, $i = \overline{1, n}$, such that $\mu(A_i) = \mu(A_j) > 0$ whenever $i \neq j$. By the same token, there are sets $W_i \subset A_i$ such that $\mu(W_i) = \lambda_i \mu(A_i)$, $i = \overline{1, n}$. Clearly, they fulfill our requirement.

Let us fix any finite $\frac{\varepsilon}{2}$ -net $(h_k)_{k=1}^N$ in K . In view of Lemma 3.1 there is an $f \in B$ verifying

$$\begin{aligned} f(h_1) &> \varepsilon + \frac{1}{\mu(W)} \int_W f(s) d\mu(s) \\ &= \varepsilon + \sum_{i=1}^n \frac{\lambda_i}{\mu(W_i)} \int_{W_i} f(s) d\mu(s). \end{aligned}$$

Then for every $i = \overline{1, n}$ one can find $(w_{ij})_{j=1}^M \subset W_i$ such that

$$\begin{aligned} f(h_1) &> \varepsilon + \sum_{i=1}^n \lambda_i \sum_{j=1}^M \frac{1}{M} f(w_{ij}) \\ &= \varepsilon + \sum_{j=1}^M \frac{1}{M} \sum_{i=1}^n \lambda_i f(w_{ij}). \end{aligned}$$

Thus,

$$\sum_{j=1}^M \frac{1}{M} \left| f(h_1) - \sum_{i=1}^n \lambda_i f(w_{ij}) \right| > \varepsilon.$$

So, for some j_0 we have

$$\left| f(h_1) - \sum_{i=1}^n \lambda_i f(w_{ij_0}) \right| > \varepsilon.$$

Since $w_{ij_0} \in W_i \subset U_i$, there are open subsets $W_i^1 \subset U_i$ such that the inequality

$$\left| f(h_1) - \sum_{i=1}^n \lambda_i f(w_i) \right| > \varepsilon$$

holds for all w_i in W_i^1 , $i = \overline{1, n}$. As a consequence, we have

$$\left\| h_1 - \sum_{i=1}^n \lambda_i w_i \right\| > \varepsilon,$$

whenever $w_i \in W_i^1$, $i = \overline{1, n}$.

Doing the same for $(W_i^1)_{i=1}^n$ instead of $(U_i)_{i=1}^n$, and h_2 instead of h_1 we find open sets $W_i^2 \subset W_i^1$ with

$$\left\| h_2 - \sum_{i=1}^n \lambda_i w_i \right\| > \varepsilon,$$

whenever $w_i \in W_i^2$, $i = \overline{1, n}$.

Continuing the process we end up with open sets $V_i = W_i^N$. It is clear from our construction that

$$\left\| h - \sum_{i=1}^n \lambda_i v_i \right\| > \frac{\varepsilon}{2},$$

for all $h \in K$ and $v_i \in V_i$. So, conditions i) and ii) are satisfied. \square

Lemma 3.3. *For any norm-compact set $K \subset \overline{\text{conv}}^\tau H$ and any collection of open sets $(U_i)_{i=1}^n$ in H there are open sets $(V_i)_{i=1}^n$ satisfying the following conditions:*

- i) $V_i \subset U_i$, $i = \overline{1, n}$;
- ii) $\text{dist}(K, \sum_{i=1}^n \lambda_i v_i) > \frac{\varepsilon}{4}$, whenever $v_i \in V_i$, $i = \overline{1, n}$, and $\lambda_i \geq 0$ with $\sum_{i=1}^n \lambda_i = 1$.

Proof. To prove this lemma we fix a finite $\frac{\varepsilon}{4}$ -net in the set

$$\{(\lambda_1, \lambda_2, \dots, \lambda_n) : \sum_{i=1}^n \lambda_i = 1, \lambda_i \geq 0\}$$

equipped with the metric $\rho((\lambda_i), (\nu_i)) = \sum_{i=1}^n |\lambda_i - \nu_i|$. Then we apply Lemma 3.2 successively to all the elements of the net. \square

Lemma 3.4. *For any collection of open sets $(U_i)_{i=1}^n$ in H there exist $f \in B$ and two constants a and b with $b - a \geq \frac{\varepsilon}{8}$ such that*

$$\begin{aligned} \{f > b\} \cap U_i &\neq \emptyset, \\ \{f < a\} \cap U_i &\neq \emptyset, \end{aligned}$$

for all $i = \overline{1, n}$.

Proof. Let us fix arbitrary $u_i \in U_i$, $i = \overline{1, n}$ and denote $K = \text{conv}(u_i)_{i=1}^n$. By Lemma 3.3, there are vectors $v_i \in U_i$ such that if $L = \text{conv}(v_i)_{i=1}^n$, then $\text{dist}(K, L) > \frac{\varepsilon}{4}$.

By the geometric version of the Hahn-Banach Theorem, there exists a $g \in B(X^*)$, $\|g\| = 1$, separating $K - L$ from the ball $\frac{\varepsilon}{4}B(X)$, i.e.

$$g(k - l) > \frac{\varepsilon}{4},$$

for all $k \in K$, $l \in L$. Since the w^* -closure of B coincides with the entire $B(X^*)$, we can find an $f \in B$, for which the inequality

$$f(k - l) > \frac{\varepsilon}{4}$$

holds, whenever $k \in K$ and $l \in L$.

Now it is easy to see that the constants $a = \sup_{l \in L} f(l) + \frac{\varepsilon}{16}$ and $b = \inf_{k \in K} f(k) - \frac{\varepsilon}{16}$ meet the desired conditions. \square

Construction of the independent sequence.

First, applying Lemma 3.4 to $U_1 = U_2 = \dots = U_n = H$ we find $f_1 \in B$ and constants a_1, b_1 with $b_1 - a_1 \geq \frac{\varepsilon}{8}$ such that

$$\begin{aligned} U_1 &= \{f_1 > b_1\} \cap H \neq \emptyset, \\ U_2 &= \{f_1 < a_1\} \cap H \neq \emptyset. \end{aligned}$$

Then we apply Lemma 3.4 to U_1, U_2 and get $f_2 \in B$, a_2, b_2 with $b_2 - a_2 \geq \frac{\varepsilon}{8}$ such that

$$\begin{aligned} \{f_2 > b_2\} \cap U_i &\neq \emptyset, \\ \{f_2 < a_2\} \cap U_i &\neq \emptyset, \quad i = 1, 2. \end{aligned}$$

It is clear how to continue the process to obtain sequences $(f_n)_{n \in \mathbf{N}} \subset B$ and $(b_n, a_n)_{n \in \mathbf{N}}$, $b_n - a_n \geq \frac{\varepsilon}{8}$ such that for all finite disjoint sets M and N in \mathbf{N} we have

$$H \cap \left(\bigcap_{m \in M} \{f_m > b_m\} \right) \cap \left(\bigcap_{n \in N} \{f_n < a_n\} \right) \neq \emptyset.$$

Of course we can assume that $|a_n - a| < \frac{\varepsilon}{32}$, for some constant a and every $n \in \mathbf{N}$. Then letting $\delta = \frac{\varepsilon}{32}$, $r = a + \frac{\varepsilon}{32}$ we finally get

$$H \cap \left(\bigcap_{m \in M} \{f_m > r + \delta\} \right) \cap \left(\bigcap_{n \in N} \{f_n < r\} \right) \neq \emptyset,$$

whenever M and N are finite disjoint subsets of \mathbf{N} . The proof is finished. \square

As explained in the introduction, as a consequence of Theorem 1.1 and Simons' result ([30]) we obtain the positive solution to the Boundary Problem in spaces not containing $\ell_1[0, 1]$. Surprisingly, this is also true for any $\ell_1(\Gamma)$ in the canonical norm. In the next section we discuss the Boundary Problem

in the classical $\ell_1(\Gamma)$ and $C(K)$ -spaces in more detail and prove even stronger results for them.

4. ANGELIC SPACES AND THE BOUNDARY PROBLEM

To motivate the results in this section we start with the following easy fact, which, in particular, yields the positive solution to the Boundary Problem under certain restrictions on the boundary.

Proposition 4.1. *Let X be a Banach space, D a norming subset of $B(X^*)$ and H a norm bounded $\sigma(X, D)$ -compact subset of X . If D is dense in $B(X^*)$ in the topology of uniform convergence on countable subsets of H , then H is weakly compact.*

Proof. It suffices to prove that H is weakly countably compact, which implies that H is weakly compact due to the Eberlein-Šmulian Theorem. Take any sequence (x_n) in H and let $x_0 \in H$ be a $\sigma(X, D)$ -cluster point of (x_n) . For any $x^* \in B_{X^*}$ and $\varepsilon > 0$ iii) implies that there is a point $d^* \in D$ such that

$$|x^*(x_n) - d^*(x_n)| < \varepsilon, \text{ for } n = 0, 1, 2, \dots$$

From this we deduce that x_0 is also a $\sigma(X, X^*)$ -cluster point of X^* and the proof is finished. \square

We stress that when D is moreover absolutely convex in the previous proposition then the fact H being weakly compact implies that D is dense in $B(X^*)$ in the topology of uniform convergence on countable subsets of H (in fact it is dense in the topology of uniform convergence on H) — bear in mind that the closures of D in the Mackey topology $\mu(X^*, X)$ and the weak* topology $\sigma(X^*, X)$ coincide and that $\overline{D}^{\sigma(X^*, X)} = B(X^*)$, see [27].

It is interesting to highlight that the thesis of Proposition 4.1 also holds if we assume that there is a boundary $B' \subset B_{X^*}$ such that:

- α) each $x^* \in B'$ is in the closure of D for the topology of uniform convergence on countable subsets of H ;
- β) norm bounded and $\sigma(X, B')$ -relatively countably compact subsets of X are weakly relatively compact.

This idea was used in [4] for $X = C(K)$ and $B' = K \cup \{-K\} \subset B(C(K)^*)$ to solve the Boundary Problem for $C(K)$ -spaces. We now establish a pure topological statement giving a new proof of this result not only for $C(K)$ but also for all $\ell^1(\Gamma)$ in their canonical norms. In fact, we prove that those spaces are angelic (see the definition below) in the topology induced by a boundary.

Definition 4.2 (Fremlin). A regular topological space E is angelic if every relatively countably compact subset A of E is relatively compact and its closure \overline{A} is made up of the limits of sequences from A .

In angelic spaces the different concepts of compactness and relative compactness coincide: the (relatively) countably compact, (relatively) compact and (relatively) sequentially compact subsets are the same, [13]. Examples of angelic spaces include $C(K)$ endowed with the topology $t_p(K)$ of pointwise convergence on a countably compact space K ([15, 22]) and all Banach spaces with their weak topologies.

The relation between angelicity and the Boundary Problem is seen from the following proposition.

Proposition 4.3. *Let X be a Banach space and let $B \subset B(X^*)$ be a boundary for X such that $(X, \sigma(X, B))$ is angelic. Then a subset H of X is weakly compact if (and only if) H is norm bounded and $\sigma(X, B)$ -countably compact.*

Proof. In view of the Eberlein-Šmulian Theorem, we only have to prove that if H is norm bounded and $\sigma(X, B)$ -compact, then H is $\sigma(X, X^*)$ -sequentially compact. Since the space $(X, \sigma(X, B))$ is angelic, for each sequence (x_n) in H there is a subsequence (x_{n_k}) and a point $x_0 \in H$ such that $x_0 = \sigma(X, B) - \lim_k x_{n_k}$. Now, Corollary 11 in [30] (see alternatively, [31, Theorem on p. 70]) straightforwardly applies to ensure that $x_0 = \sigma(X, X^*) - \lim_k x_{n_k}$. The proof is over. \square

It is not difficult to prove that if X is a separable Banach space then, for any boundary $B \subset B(X^*)$ the space $(X, \sigma(X, B))$ is angelic. Although there are boundaries in the nonseparable case that also provide angelic topologies. For instance, the one with $C(K)$ we mentioned above.

Another example of this phenomenon is given by our next proposition.

Proposition 4.4. *Let Γ be any set and $D = \{-1, 1\}^\Gamma$ the set of the extreme points of $B(\ell_\infty(\Gamma))$. Then,*

- i) $(\ell_1(\Gamma), \sigma(\ell_1(\Gamma), D))$ is angelic;
- ii) If $H \subset \ell_1(\Gamma)$ is $\|\cdot\|_1$ -bounded and $\sigma(\ell_1(\Gamma), D)$ -compact then H is weakly compact.

Proof. To prove i) observe first that $D \subset B(\ell_\infty(\Gamma))$, $(D, \sigma(\ell_\infty(\Gamma), \ell_1(\Gamma)))$ is compact and $(C(D), t_p(D))$ is angelic, [13]. The natural embedding

$$(\ell_1(\Gamma), \sigma(\ell_1(\Gamma), D)) \rightarrow (C(D), t_p(D))$$

is a homeomorphism onto its image. Then the angelicity of the space $\ell_1(\Gamma)$ in the topology $\sigma(\ell_1(\Gamma), D)$ follows from the angelicity of $(C(D), t_p(D))$. Statement ii) is a straightforward consequence of Proposition 4.3. \square

In Theorem 4.9 we will prove that statements as in Proposition 4.4 hold for all boundaries of $B(\ell_\infty(\Gamma))$. Still let us remark that ii) was alternatively obtained in [17, Theorem 10.5.2] using Schur's Lemma for $\ell_1(\Gamma)$.

The next lemma will allow us to transfer the angelic property from one topology to another. We will use it later in the proofs of Theorems 4.8 and 4.9.

Lemma 4.5. *Let X be a non-empty set and τ, \mathfrak{T} two Hausdorff topologies on X such that (X, τ) is regular and (X, \mathfrak{T}) is angelic. Assume that for every sequence (x_n) in X with a τ -cluster point $x \in X$, x is \mathfrak{T} -cluster point of (x_n) . Then, the following assertions hold:*

- i) *If $L \subset X$ is τ -relatively countably compact, then L is \mathfrak{T} -relatively compact;*
- ii) *If $L \subset X$ is τ -compact, then L is \mathfrak{T} -compact;*
- iii) *(X, τ) is an angelic space.*

Proof. Statement i) is a straightforward consequence of the assumptions on τ -cluster points of sequences in X and the fact that (X, \mathfrak{T}) is angelic.

Let us prove ii). If $L \subset X$ is τ -compact, then L is \mathfrak{T} -relatively compact by i). To finish the proof of ii) it will be enough to show that L is \mathfrak{T} -closed. Pick $x \in \overline{L}^{\mathfrak{T}}$. Using that (X, \mathfrak{T}) is angelic, there is a sequence (x_n) in L with

$$(8) \quad x = \mathfrak{T} - \lim_{n \rightarrow \infty} x_n$$

By τ -compactness, there is $y \in L$ which is a τ -cluster point of (x_n) . Our assumption implies that y is a \mathfrak{T} -cluster point of (x_n) , hence by (8) $y = x$ and thus $x \in L$. The proof of ii) is concluded.

The proof of iii) relies upon the following

Claim 4.6. *If L is a τ -relatively countably compact and countable subset of X , then*

$$(9) \quad \overline{L}^{\tau} = \overline{L}^{\mathfrak{T}}$$

and the topologies τ and \mathfrak{T} coincide on $\overline{L}^{\mathfrak{T}}$.

Suppose for a moment that the claim is true and let us prove that (X, τ) is angelic: to this end we will show that if $A \subset X$ is τ -relatively countably compact then $\overline{A}^{\tau} = \overline{A}^{\mathfrak{T}}$ is τ -compact and τ and \mathfrak{T} coincide on \overline{A}^{τ} . We already know that $\overline{A}^{\mathfrak{T}}$ is \mathfrak{T} -compact by i). Now we will prove that the identity map

$$id : (\overline{A}^{\mathfrak{T}}, \mathfrak{T}) \longrightarrow (\overline{A}^{\mathfrak{T}}, \tau)$$

is continuous, that is, we will show that any τ -closed subset of $\overline{A}^{\mathfrak{T}}$ is \mathfrak{T} -closed. Indeed, take a τ -closed subset F of $\overline{A}^{\mathfrak{T}}$. Pick any $x \in \overline{F}^{\mathfrak{T}}$. The angelicity of (X, \mathfrak{T}) provides us with a sequence (x_n) in F such that

$$x = \mathfrak{T} - \lim_{n \rightarrow \infty} x_n$$

Now for every $n \in \mathbf{N}$ we can also take (x_{mn}) in A such that

$$x_n = \mathfrak{T} - \lim_{m \rightarrow \infty} x_{mn}.$$

If we define $L = \{x_{mn} : m, n \in \mathbf{N}\}$ then the claim tells us that τ and \mathfrak{T} coincide on $\overline{L}^{\mathfrak{T}}$ and so

$$x = \tau - \lim_{n \rightarrow \infty} x_n$$

what implies that $x \in F$. So $\overline{A}^{\mathfrak{T}}$ is τ -compact and τ and \mathfrak{T} coincide on $\overline{A}^{\mathfrak{T}}$. So, being $\overline{A}^{\mathfrak{T}}$ τ -closed we obtain $\overline{A}^{\tau} \subset \overline{A}^{\mathfrak{T}}$. On the other hand, as $A \subset \overline{A}^{\mathfrak{T}}$, A is τ -relatively compact and so \overline{A}^{τ} is \mathfrak{T} -compact after ii). Therefore $\overline{A}^{\mathfrak{T}} \subset \overline{A}^{\tau}$, thus $\overline{A}^{\mathfrak{T}} = \overline{A}^{\tau}$ and the proof is complete.

Let us now prove Claim 4.6.

From the assumptions we have $\overline{L}^{\tau} \subset \overline{L}^{\mathfrak{T}}$. Conversely, if we pick x in the \mathfrak{T} -compact subset $\overline{L}^{\mathfrak{T}}$, then the angelicity of (X, \mathfrak{T}) ensures us of the existence of a sequence (x_n) in L such that

$$(10) \quad x = \mathfrak{T} - \lim_{n \rightarrow \infty} x_n$$

By the τ -relatively countably compactness of L , there is $y \in \overline{L}^{\tau}$ which is a τ -cluster point of (x_n) . Therefore y is a \mathfrak{T} -cluster point of (x_n) , hence by (10) $y = x$ and thus $x \in \overline{L}^{\tau}$ what implies the equality (9).

To prove that the topologies τ and \mathfrak{T} coincide on $H := \overline{L}^{\mathfrak{T}}$ it suffice to show, by compactness, that the identity

$$id : (H, \mathfrak{T}) \longrightarrow (H, \tau)$$

is continuous. To this end we will establish that any τ -closed subset F of H is \mathfrak{T} -closed. Indeed, as (H, τ) is a regular topological space we have

$$F = \cap \{\overline{U}^{\tau} : F \subset U \subset H, U \text{ is } \tau\text{-open in } H\}$$

On the other hand for any such a U we have that $\overline{U}^{\tau} = \overline{U \cap L}^{\tau}$ and we can apply the equality (9) to $U \cap L$ to conclude that

$$\overline{U \cap L}^{\tau} = \overline{U \cap L}^{\mathfrak{T}}$$

This implies

$$F = \cap \{\overline{U \cap L}^{\mathfrak{T}} : F \subset U \subset H, U \text{ is } \tau\text{-open in } H\}$$

and so F is \mathfrak{T} -closed. \square

Note that our hypothesis in Claim 4.6 about L , namely, L countable and relatively countably compact in X do not imply (in general) that L has to be relatively compact in X . Indeed, take $\beta\mathbf{N}$ the Stone-Ćech compactification of the natural numbers \mathbf{N} and pick a point $p \in \beta\mathbf{N} \setminus \mathbf{N}$. Take $X := \beta\mathbf{N} \setminus \{p\}$ and $L = \mathbf{N}$. It is well known that an infinite set in L cannot have a unique

cluster point in $\beta\mathbf{N} \setminus \mathbf{N}$. This proves that L is relatively countably compact in X but its closure in X , $\overline{L}^X = X$, is not compact.

The lemma below can be found in [4]. Here we include a slightly different proof that does not use the Uryshon Lemma.

Lemma 4.7. *Let K be a compact space and $B \subset B(C(K)^*)$ a boundary for $(C(K), \|\cdot\|_\infty)$. If (f_n) is an arbitrary sequence in $C(K)$ and $x \in K$, then there is $\mu \in B$ such that*

$$f_n(x) = \int_K f_n d\mu$$

for every $n \in \mathbf{N}$.

Proof. If we define the continuous function $g : K \rightarrow [0, 1]$ by the expression

$$g(t) := 1 - \sum_{n=1}^{\infty} \frac{1}{2^n} \frac{|f_n(t) - f_n(x)|}{1 + |f_n(t) - f_n(x)|}, \quad t \in K,$$

then

$$(11) \quad F = \bigcap_{n=1}^{\infty} \{y \in K; f_n(y) = f_n(x)\} = \{y \in K : g(y) = 1 = \|g\|_\infty\}.$$

Since B is a boundary, there exists $\mu \in B$ such that $\int_K g d\mu = 1$. From here we obtain

$$(12) \quad 1 = \|\mu\| = |\mu|(K) \geq \int_K g d|\mu| \geq \int_K g d\mu = 1.$$

In other words,

$$0 = |\mu|(K) - \int_K g d|\mu| = \int_K (1 - g) d|\mu|.$$

Since $1 - g \geq 0$ we obtain $|\mu|(\{y \in K : 1 - g(y) > 0\}) = 0$, that is $|\mu|(K \setminus F) = 0$. Therefore, for every $n \in \mathbf{N}$ we have

$$\int_K f_n d\mu = \int_F f_n d\mu = \int_F f_n(x) d\mu = f_n(x)$$

because $\mu(F) = \int_F g d\mu = \int_K g d\mu = 1$ by the equalities (11) and (12). \square

We naturally arrive at the following.

Theorem 4.8 ([4]). *Let K be a compact space and $B \subset B(C(K)^*)$ a boundary for $C(K)$. Then $(C(K), \sigma(C(K), B))$ is an angelic space. Consequently, if $H \subset C(K)$ is norm bounded and $\sigma(C(K), B)$ -countably compact, then H is weakly compact.*

Proof. The space $(C(K), t_p(K))$ is angelic, [15, 22] (see also [13]). Bearing this in mind, the first part of the theorem follows from Lemmas 4.7 and 4.5 applied to $X = C(K)$, $\tau = \sigma(C(K), B)$ and $\mathfrak{T} = t_p(K)$.

The second part of the theorem follows from Proposition 4.3. \square

The game we played for spaces $C(K)$ can be played for $\ell_1(\Gamma)$ too.

Theorem 4.9. *Let Γ be any set and $B \subset B(\ell_\infty(\Gamma))$ a boundary for $(\ell_1(\Gamma), \|\cdot\|_1)$. Then,*

- i) $(\ell_1(\Gamma), \sigma(\ell_1(\Gamma), B))$ is angelic;
- ii) If $H \subset \ell_1(\Gamma)$ is $\|\cdot\|_1$ -bounded and $\sigma(X, B)$ -compact then H is weakly compact.

Proof. The fact that B is a boundary implies that for any countable subset $A \subset \Gamma$ and any family of signs $(y_\gamma)_{\gamma \in A} \in \{-1, 1\}^A$, there is $(b_\gamma)_{\gamma \in \Gamma}$ in B such that $b_\gamma = y_\gamma$, for $\gamma \in A$. According to this, if $D = \{-1, 1\}^\Gamma$, $d^* \in D$ and we take a sequence $(z_n)_n \in \ell_1(\Gamma)$ there is $b^* \in B$ such that

$$d^*(z_n) = b^*(z_n)$$

for every $n \in \mathbb{N}$. Due to Proposition 4.4, the space $\ell^1(\Gamma)$ is angelic in the topology $\sigma(\ell_1(\Gamma), D)$. Therefore, statement i) follows from Lemma 4.5 applied to $\tau = \sigma(\ell_1(\Gamma), B)$ and $\mathfrak{T} = \sigma(\ell_1(\Gamma), D)$. Statement ii) is now a consequence of Proposition 4.3. \square

We finish with two questions still open to the best of our knowledge.

A result by Bourgain and Talagrand ([3]) states that if X is a Banach space and H is a norm bounded and $\sigma(X, \text{ext}B(X^*))$ -countably compact subset of X , then H is weakly compact (Rainwater's theorem is a weak version of this). Therefore, a positive solution of the problem below would imply the Boundary Problem.

Problem 4.10. Let X be a Banach space, $B \subset B(X^*)$ a boundary and $D = \text{conv}(B \cup \{-B\})$. Given $e^* \in \text{ext}B(X^*)$, $\varepsilon > 0$ and a sequence $(x_n)_n$, is there $d^* \in D$ such that

$$|d^*(x_n) - e^*(x_n)| < \varepsilon,$$

for every $n \in \mathbb{N}$?

Problem 4.11. Is a Banach space X angelic in the topology $\sigma(X, \text{ext}B(X^*))$?

REFERENCES

- [1] Jürgen Batt and Wolfgang Hiermeyer, *On compactness in $L_p(\mu, X)$ in the weak topology and in the topology $\sigma(L_p(\mu, X), L_q(\mu, X'))$* , Math. Z. **182** (1983), no. 3, 409–423. MR 84m:46039
- [2] G. Birkhoff, *Integration of functions with values in a Banach space*, Trans. Amer. Math. Soc. **38** (1935), no. 2, 357–378. MR 1 501 815
- [3] Jean Bourgain and Michel Talagrand, *Compacité extrême*, Proc. Amer. Math. Soc. **80** (1980), no. 1, 68–70. MR 81h:46011
- [4] B. Cascales and G. Godefroy, *Angelicity and the boundary problem*, Mathematika **45** (1998), no. 1, 105–112. MR 99f:46019
- [5] B. Cascales, G. Manjabacas, and G. Vera, *A Krein-Šmulian type result in Banach spaces*, Quart. J. Math. Oxford Ser. (2) **48** (1997), no. 190, 161–167. MR 99c:46009

- [6] B. Cascales, I. Namioka, and G. Vera, *The Lindelöf property and fragmentability*, Proc. Amer. Math. Soc. **128** (2000), no. 11, 3301–3309.
- [7] B. Cascales and J. Rodriguez, *Birkhoff integral and the property of Bourgain*, Preprint, 2003.
- [8] B. Cascales and G. Vera, *Topologies weaker than the weak topology of a Banach space*, J. Math. Anal. Appl. **182** (1994), no. 1, 41–68. MR 95c:46017
- [9] ———, *Norming sets and compactness*, Rocky Mountain J. Math. **25** (1995), no. 3, 919–925. MR 96i:46011
- [10] Gustave Choquet, *Lectures on analysis. Vol. II: Representation theory*, Edited by J. Marsden, T. Lance and S. Gelbart, W. A. Benjamin, Inc., New York-Amsterdam, 1969. MR 40 #3253
- [11] J. Diestel, *Sequences and series in Banach spaces*, Graduate Texts in Mathematics, vol. 92, Springer-Verlag, New York, 1984. MR 85i:46020
- [12] J. Diestel and J. J. Uhl, Jr., *Vector measures*, American Mathematical Society, Providence, R.I., 1977, With a foreword by B. J. Pettis, Mathematical Surveys, No. 15. MR 56 #12216
- [13] K. Floret, *Weakly compact sets*, Springer, Berlin, 1980, Lectures held at S.U.N.Y., Buffalo, in Spring 1978. MR 82b:46001
- [14] Gilles Godefroy, *Boundaries of a convex set and interpolation sets*, Math. Ann. **277** (1987), no. 2, 173–184. MR 88f:46037
- [15] A. Grothendieck, *Critères de compacité dans les espaces fonctionnels généraux*, Amer. J. Math. **74** (1952), 168–186. MR 13,857e
- [16] P. Habala, P. Hájek, and Zizler V., *Introduction to Banach spaces*, Matfyz Press, Prague, 1996.
- [17] H. Jarchow, *Locally convex spaces*, B. G. Teubner, Stuttgart, 1981, Mathematische Leitfäden. [Mathematical Textbooks].
- [18] J. E. Jayne and C. A. Rogers, *Borel selectors for upper semicontinuous set-valued maps*, Acta Math. **155** (1985), no. 1-2, 41–79. MR 87a:28011
- [19] V. Kadets, B. Shumyatskiy, R. Shvidkoy, L. Tseytlin, and Zheltukhin K., *Some remarks on vector-valued integration.*, Mat. Fiz. Anal. Geom. **9** (2002), no. 1, 48–65.
- [20] V. M. Kadets and L. M. Tseytlin, *On “integration” of non-integrable vector-valued functions*, Mat. Fiz. Anal. Geom. **7** (2000), no. 1, 49–65. MR 2001e:28017
- [21] J. L. Kelley, *General topology*, Springer-Verlag, New York, 1975, Reprint of the 1955 edition [Van Nostrand, Toronto, Ont.], Graduate Texts in Mathematics, No. 27.
- [22] J. L. Kelley and Isaac Namioka, *Linear topological spaces*, D. Van Nostrand Co., Inc., Princeton, N.J., 1963. MR 29 #3851
- [23] M. Krein and V. Šmulian, *On regular convex sets in the space conjugate to a Banach space*, Ann. of Math. (2) **41** (1940), 556–583. MR 1,335e
- [24] I. Namioka, *Separate continuity and joint continuity*, Pacific J. Math. **51** (1974), 515–531. MR 51 #6693
- [25] ———, *Radon-Nikodým compact spaces and fragmentability*, Mathematika **34** (1987), no. 2, 258–281. MR 89i:46021
- [26] Robert R. Phelps, *Lectures on Choquet’s theorem*, second ed., Lecture Notes in Mathematics, vol. 1757, Springer-Verlag, Berlin, 2001. MR 2002k:46001
- [27] A. P. Robertson and W. Robertson, *Topological vector spaces*, second ed., Cambridge University Press, London, 1973, Cambridge Tracts in Mathematics and Mathematical Physics, No. 53. MR 50 #2854
- [28] H. P. Rosenthal, *A characterization of Banach spaces containing l_1* , Proc. Nat. Acad. Sci. U.S.A. **71** (1974), 2411–2413. MR 50 #10773
- [29] R. Shvydkoy, *Operators and integrals in Banach spaces*, Ph.D. thesis, University of Missouri - Columbia, 2001.

- [30] S. Simons, *A convergence theorem with boundary*, Pacific J. Math. **40** (1972), 703–708. MR 47 #755
- [31] S. Simons, *An eigenvector of Fatou's lemma for continuous functions*, The Mathematical Intelligencer **17** (1995), no. 3, 67–70.

DEPARTAMENTO DE MATEMÁTICAS, FACULTAD DE MATEMÁTICAS, UNIVERSIDAD DE MURCIA, 30.100 ESPINARDO, MURCIA, SPAIN

E-mail address: `beca@um.es`

DEPARTMENT OF MATHEMATICS C1200, UNIVERSITY OF TEXAS, AUSTIN, TX 78712, USA

E-mail address: `shvidkoy@math.utexas.edu`