

MATH 504 HOMEWORK 5

Due Friday, April 1.

Problem 1. Show that there is a projection $\pi : \text{Add}(\omega, \lambda) \rightarrow \text{Add}(\omega, 1)$.

Problem 2. Let M be a transitive model of ZFC and $\mathbb{P} \in M$ be a poset. Suppose that $p \in \mathbb{P}$ is such that $p \Vdash \dot{f} : \lambda \rightarrow \tau$ is a function”.

- (1) Show that for every $\alpha < \lambda$, $\{q \mid \exists \gamma \in \tau q \Vdash \dot{f}(\alpha) = \gamma\}$ is dense below p .
- (2) Let $B = \{\gamma < \tau \mid (\exists q \leq p)(\exists \alpha < \lambda)(q \Vdash \dot{f}(\alpha) = \gamma)\}$. Show that if $\sup(B) < \tau$, then $p \Vdash \dot{f}$ is bounded”.

We say that \mathbb{P} preserves cofinalities if for every ordinal α , if in V , $\text{cf}(\alpha) = \tau$, then $1_{\mathbb{P}} \Vdash \text{cf}(\alpha) = \tau$.

Problem 3. Prove (in detail) that if \mathbb{P} preserves cofinalities, then \mathbb{P} preserves cardinals.

Problem 4. Suppose \mathbb{P} and \mathbb{Q} are two posets and $i : \mathbb{P} \rightarrow \mathbb{Q}$ is such that:

- $i(1_{\mathbb{P}}) = 1_{\mathbb{Q}}$;
- if $p' \leq_{\mathbb{P}} p$, then $i(p') \leq_{\mathbb{Q}} i(p)$;
- for all $p_1, p_2 \in \mathbb{P}$, $p_1 \perp p_2$ iff $i(p_1) \perp i(p_2)$;
- If \mathcal{A} is a maximal antichain of \mathbb{P} , then $i''\mathcal{A} := \{i(p) \mid p \in \mathcal{A}\}$ is a maximal antichain in \mathbb{Q} .

Suppose also that H is \mathbb{Q} -generic. Show that $G := \{p \in \mathbb{P} \mid i(p) \in H\}$ is \mathbb{P} -generic and that $V[G] \subset V[H]$, where V is the ground model.