


Density-type theorems for semi-algebraic hypergraphs

Jacob Fox (MIT), Janos Pach (EPFL), Andrew Suk (UIC)

November 13, 2014


An old Ramsey-type result, Kövári, Sós, and Turán and Erdős

Bipartite graph G , edge set E .




An old Ramsey-type result, Kövári, Sós, and Turán and Erdős

Bipartite graph G , edge set E .




An old Ramsey-type result, Kővári, Sós, and Turán and Erdős


Bipartite graph G , edge set E .




k -partite k -uniform hypergraph H , edge set E .




k -partite k -uniform hypergraph H , edge set E .



k -partite k -uniform hypergraph H , edge set E .




k -partite k -uniform hypergraph H , edge set E .




These results are tight.

k -partite k -uniform hypergraph H , edge set E .




In this talk: We can do much better if H is a semi-algebraic k -uniform hypergraph.

k -partite k -uniform hypergraph H , edge set E .




Semi-algebraic hypergraphs: $V = \{\text{simple geometric objects in } \mathbb{R}^d\}$, $E = \{\text{simple relation on } k \text{ tuples of } V\}$.


$V = \{A_1, \dots, A_n\}$, n disks in the plane. $E = \{\text{pairs of disks that intersect}\}$.




$V = \{A_1, \dots, A_n\}$, n disks in the plane. $E = \{\text{pairs of disks that intersect}\}$.



$V = \{A_1, \dots, A_n\}$, n disks in the plane. $E = \{\text{pairs of disks that intersect}\}$.




$V = \{A_1, \dots, A_n\}$, n disks in the plane. $E = \{\text{pairs of disks that intersect}\}$.



$A_i \rightarrow p_i = (x_i, y_i, r_i)$, $A_j \rightarrow p_j = (x_j, y_j, r_j)$. A_i and A_j cross if and only if

$$-x_i^2 + 2x_i x_j - x_j^2 - y_i^2 + 2y_i y_j - y_j^2 + r_i^2 + 2r_i r_j + r_j^2 \geq 0.$$

$V = \{A_1, \dots, A_n\}$, n disks in the plane. $E = \{ \text{pairs of disks that intersect} \}$.



Graph $G = (V, E)$, $V = n$ points in \mathbb{R}^3

E defined by the polynomial



$$f(z_1, \dots, z_6) = -z_1^2 + 2z_1z_4 - z_4^2 - z_2^2 + 2z_2z_5 - z_5^2 + z_3^2 + 2z_3z_6 + z_6^2.$$

$$(p_i, p_j) \in E \Leftrightarrow f(p_i, p_j) \geq 0.$$

More examples of semi-algebraic hypergraphs

Examples



- 1 $V = \{n \text{ circles in } \mathbb{R}^3\}$
 $E = \{\text{pairs that are linked}\}.$
- 2 $V = \{n \text{ hyperplanes in } \mathbb{R}^d \text{ in general position}\},$
 $E = \{d\text{-tuples whose intersection point is above the}$
 $\text{hyperplane } x_d = 0\}.$



More examples of semi-algebraic hypergraphs

Examples

- 1 $V = \{n \text{ circles in } \mathbb{R}^3\} \rightarrow n \text{ points in higher dimensions.}$
 $E = \{\text{pairs that are linked}\} \rightarrow \text{polynomials } f_1, \dots, f_t.$
- 2 $V = \{n \text{ hyperplanes in } \mathbb{R}^d \text{ in general position}\} \rightarrow n \text{ points in higher dimensions,}$
 $E = \{d\text{-tuples whose intersection point is above the hyperplane } x_d = 0\} \rightarrow \text{polynomials } f_1, \dots, f_t.$



We say that $H = (V, E)$ is a **semi-algebraic k -uniform hypergraph in d -space** if

$$V = \{n \text{ points in } \mathbb{R}^d\}$$

E defined by polynomials f_1, \dots, f_t and a Boolean formula Φ such that

$$(p_{i_1}, \dots, p_{i_k}) \in E$$

$$\Leftrightarrow \Phi(f_1(p_{i_1}, \dots, p_{i_k}) \geq 0, \dots, f_t(p_{i_1}, \dots, p_{i_k}) \geq 0) = \text{yes}$$


Example

3-uniform hypergraph $H = (V, E)$, $V = \{p_1, \dots, p_n\}$ points in \mathbb{R}^d .

Relation $E \subset \binom{V}{3}$ depends on f and Φ

$$\phi(f(x_1, x_2, x_3) \geq 0) = \{\text{yes, no}\}$$

$(p_1, p_2, p_3) \in E$ depends on $f(p_1, p_2, p_3) \rightarrow \{+, -, 0\}$.




Example

3-uniform hypergraph $H = (V, E)$, $V = \{p_1, \dots, p_n\}$ points in \mathbb{R}^d .

Relation $E \subset \binom{V}{3}$ depends on f and Φ

$$\phi(f(x_1, x_2, x_3) \geq 0) = \{\text{yes, no}\}$$

$(p_1, p_2, p_3) \in E$ depends on $f(p_1, p_2, p_3) \rightarrow \{+, -, 0\}$.




Example

3-uniform hypergraph $H = (V, E)$, $V = \{p_1, \dots, p_n\}$ points in \mathbb{R}^d .

Relation $E \subset \binom{V}{3}$ depends on f and Φ

$$\phi(f(x_1, x_2, x_3) \geq 0) = \{\text{yes, no}\}$$

Zero set $f(p_1, p_2, x_3) = 0$, surface in \mathbb{R}^d .




Example

3-uniform hypergraph $H = (V, E)$, $V = \{p_1, \dots, p_n\}$ points in \mathbb{R}^d .

Relation $E \subset \binom{V}{3}$ depends on f and Φ

$$\phi(f(x_1, x_2, x_3) \geq 0) = \{\text{yes, no}\}$$

Zero set $f(p_1, p_2, x_3) = 0$, surface in \mathbb{R}^d .




Example

3-uniform hypergraph $H = (V, E)$, $V = \{p_1, \dots, p_n\}$ points in \mathbb{R}^d .

Relation $E \subset \binom{V}{3}$ depends on f and Φ

$$\phi(f(x_1, x_2, x_3) \geq 0) = \{\text{yes, no}\}$$

Zero set $f(p_1, p_2, x_3) = 0$, surface in \mathbb{R}^d .



E has complexity (t, D) , degree of $f(p_1, p_2, x_3) \leq D$.

Complexity of relation E

$$x_i \in \mathbb{R}^d$$

E has complexity (t, D)


- 1 described by polynomials f_1, \dots, f_t ,
- 2 and the degree of ALL kt d -variate polynomials $f_i(x_1, \dots, x_{k-1}, x_k), f_i(x_1, \dots, x_{k-2}, x_{k-1}, x_k), \dots, f_i(x_1, x_2, \dots, x_k)$, for $i = 1 \dots t$, is at most D .

Note. f_i has degree at most Dk .

Motivation: Orientations and order-types


Our motivation, E is related to order-types and orientations.

$f(p_1, p_2, x_3)$ is linear.



E has complexity $(t, D) = (t, 1)$.

Example




$V = \{n \text{ points in the plane}\},$

$E = \{\text{triples having a clockwise orientation}\}.$

$H = (V, E)$ semi-algebraic 3-uniform hypergraph in the plane
($d = 2$)

Example




$V = \{n \text{ points in the plane}\},$

$E = \{\text{triples having a clockwise orientation}\}.$

$H = (V, E)$ semi-algebraic 3-uniform hypergraph in the plane
($d = 2$)

Example




$V = \{n \text{ points in the plane}\},$

$E = \{\text{triples having a clockwise orientation}\}.$

$H = (V, E)$ semi-algebraic 3-uniform hypergraph in the plane

$(d = 2)$

Example




$V = \{n \text{ points in the plane}\},$

$E = \{\text{triples having a clockwise orientation}\}.$

$H = (V, E)$ semi-algebraic 3-uniform hypergraph in the plane

$(d = 2)$


Example



$E = \{\text{triples having a clockwise orientation}\}.$

$$\det \begin{pmatrix} 1 & 1 & 1 \\ x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{pmatrix} > 0.$$


Example



Complexity of E is (t, D) , where $t = 1, D = 1$

$$\det \begin{pmatrix} 1 & 1 & 1 \\ x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{pmatrix} > 0.$$


Example



Complexity of E is (t, D) , where $t = 1, D = 1$

$$\det \begin{pmatrix} 1 & 1 & 1 \\ x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{pmatrix} > 0$$


Example



Complexity of E is (t, D) , where $t = 1, D = 1$

$$\det \begin{pmatrix} 1 & 1 & 1 \\ x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{pmatrix} > 0$$

Example



Complexity of E is (t, D) , where $t = 1, D = 1$

$$\det \begin{pmatrix} 1 & 1 & 1 \\ x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{pmatrix} > 0$$

Example in higher dimensions

$E = (d + 1)$ -tuples with a positive orientation, complexity
 $(t, D) = (1, 1)$.

$$\det \begin{pmatrix} 1 & 1 & \cdots & 1 \\ | & | & \cdots & | \\ p_{i_1} & p_{i_2} & \vdots & p_{i_{d+1}} \\ | & | & \cdots & | \end{pmatrix} > 0.$$

Example in higher dimensions

$E = (d + 1)$ -tuples with a positive orientation, complexity
 $(t, D) = (1, 1)$.

$$\det \begin{pmatrix} 1 & 1 & \cdots & 1 \\ | & | & \cdots & | \\ p_{i_1} & p_{i_2} & \vdots & p_{i_{d+1}} \\ | & | & \cdots & | \end{pmatrix} > 0.$$

Example in higher dimensions


$E = (d + 1)$ -tuples with a positive orientation, complexity
 $(t, D) = (1, 1)$.

$$\det \begin{pmatrix} 1 & \mathbf{1} & \cdots & \mathbf{1} \\ | & | & \cdots & | \\ p_{i_1} & \mathbf{p}_{i_2} & \vdots & \mathbf{p}_{i_{d+1}} \\ | & | & \cdots & | \end{pmatrix} > 0.$$

Previous results

Theorem (Alon, Pach, Pinchasi, Radoicic, Sharir 2005)


Let $H = (V_1, V_2, E)$ be a bipartite semi-algebraic graph ($k = 2$) in d -space, where $|V_1| = |V_2| = n$ and E has complexity (t, D) . Then there are subsets $V'_1, V'_2 \subset V$ such that $|V'_i| \geq \epsilon n$ and either $(V'_1, V'_2) \subset E$ or $(V'_1, V'_2) \subset \bar{E}$, and $\epsilon = \epsilon(d, t, D)$.



Previous results

Theorem (Alon, Pach, Pinchasi, Radoicic, Sharir 2005)


Let $H = (V_1, V_2, E)$ be a bipartite semi-algebraic graph ($k = 2$) in d -space, where $|V_1| = |V_2| = n$ and E has complexity (t, D) . Then there are subsets $V'_1, V'_2 \subset V$ such that $|V'_i| \geq \epsilon n$ and either $(V'_1, V'_2) \subset E$ or $(V'_1, V'_2) \subset \bar{E}$, and $\epsilon = \epsilon(d, t, D)$.



Previous results

Theorem (Alon, Pach, Pinchasi, Radoicic, Sharir 2005)

Let $H = (V_1, V_2, E)$ be a bipartite semi-algebraic graph ($k = 2$) in d -space, where $|V_1| = |V_2| = n$ and E has complexity (t, D) . Then there are subsets $V'_1, V'_2 \subset V$ such that $|V'_i| \geq \epsilon n$ and either $(V'_1, V'_2) \subset E$ or $(V'_1, V'_2) \subset \bar{E}$, and $\epsilon = \epsilon(d, t, D)$.




Stronger density theorem

Including an argument of Komlos:

Theorem (Alon, Pach, Pinchasi, Radoicic, Sharir 2005)

Let $H = (V_1, V_2, E)$ be a bipartite semi-algebraic graph ($k = 2$) in d -space, where $|V_1| = |V_2| = n$ and E has complexity (t, D) . If $|E| \geq \epsilon n^2$, then there are subsets $V'_1, V'_2 \subset V$ such that $|V'_i| \geq \epsilon^C n$ where $C = C(d, t, D)$, and $(V'_1, V'_2) \subset E$.




Stronger density theorem

Including an argument of Komlos:

Theorem (Alon, Pach, Pinchasi, Radoicic, Sharir 2005)


Let $H = (V_1, V_2, E)$ be a bipartite semi-algebraic graph ($k = 2$) in d -space, where $|V_1| = |V_2| = n$ and E has complexity (t, D) . If $|E| \geq \epsilon n^2$, then there are subsets $V'_1, V'_2 \subset V$ such that $|V'_i| \geq \epsilon^C n$ where $C = C(d, t, D)$, and $(V'_1, V'_2) \subset E$.



Generalization

Theorem (Fox, Gromov, Lafforgue, Naor, Pach 2012, Bukh and Hubard 2012)


Let $H = (V_1, \dots, V_k, E)$ be a k -partite semi-algebraic k -uniform hypergraph in d -space, where $|V_1| = \dots = |V_k| = n$ and E has complexity (t, D) . If $|E| \geq \epsilon n^k$, then there are subsets $V'_1, \dots, V'_k \subset V$ such that $|V'_i| \geq \epsilon^C n$ where $C = C(k, d, t, D)$, and $(V'_1, \dots, V'_k) \subset E$.



Generalization

Theorem (Fox, Gromov, Lafforgue, Naor, Pach 2012, Bukh and Hubard 2012)

Let $H = (V_1, \dots, V_k, E)$ be a k -partite semi-algebraic k -uniform hypergraph in d -space, where $|V_1| = \dots = |V_k| = n$ and E has complexity (t, D) . If $|E| \geq \epsilon n^k$, then there are subsets $V'_1, \dots, V'_k \subset V$ such that $|V'_i| \geq \epsilon^C n$ where $C = C(k, d, t, D)$, and $(V'_1, \dots, V'_k) \subset E$.




Generalization

$C(k, d, t, D)$: Dependency on uniformity k is very bad.

Fox, Gromov, Lafforgue, Naor, Pach: $C(k, d, t, D) \sim \underbrace{2^{2^{\dots 2^d}}}_k$

(tower-type)

Bukh-Hubard: $C(k, d, t, D) \sim 2^{2^{k+d}}$, double exponential in k .




Bukh-Hubard: Set sizes decay triple exponentially in k

New results

For simplicity, complexity (t, D) is fixed.


Theorem (Fox, Pach, S., 2014)

Let $H = (V_1, \dots, V_k, E)$ be a k -partite semi-algebraic k -uniform hypergraph in d -space, where $|V_1| = \dots = |V_k| = n$ and E has complexity (t, D) . If $|E| \geq \epsilon n^k$, then there are subsets $V'_1, \dots, V'_k \subset V$ such that $|V'_i| \geq \frac{\epsilon^{d^c}}{2^{ckd^c}} n$, and $(V'_1, \dots, V'_k) \subset E$.




Theorem (Fox, Pach, S., 2014)

Let $H = (V_1, \dots, V_k, E)$ be a k -partite semi-algebraic k -uniform hypergraph in d -space, where $|V_1| = \dots = |V_k| = n$ and E has complexity (t, D) . If $|E| \geq \epsilon n^k$, then there are subsets $V'_1, \dots, V'_k \subset V$ such that $|V'_i| \geq \frac{\epsilon^{d^c}}{2^{ckd^c}} n$, and $(V'_1, \dots, V'_k) \subset E$.



Theorem (Fox, Pach, S., 2014)

Let $H = (V_1, \dots, V_k, E)$ be a k -partite semi-algebraic k -uniform hypergraph in d -space, where $|V_1| = \dots = |V_k| = n$ and E has complexity $(t, 1)$. If $|E| \geq \epsilon n^k$, then there are subsets $V'_1, \dots, V'_k \subset V$ such that $|V'_i| \geq \frac{\epsilon^{d+1}}{2^{ckd \log d}} n$, and $(V'_1, \dots, V'_k) \subset E$.




Applications, Tverberg-type result

Theorem (Pach, 1998)

Let $P_1, P_2, \dots, P_{d+1} \subset \mathbb{R}^d$ be disjoint n -element point sets with $P_1 \cup \dots \cup P_{d+1}$ in general position. Then there is a point $q \in \mathbb{R}^d$ and subsets $P'_1 \subset P_1, \dots, P'_{d+1} \subset P_{d+1}$, with

$$|P'_i| \geq 2^{-2^{2^{O(d)}}} n,$$

such that all closed rainbow simplices contains q .




Applications, Tverberg-type result

Theorem (Pach, 1998)

Let $P_1, P_2, \dots, P_{d+1} \subset \mathbb{R}^d$ be disjoint n -element point sets with $P_1 \cup \dots \cup P_{d+1}$ in general position. Then there is a point $q \in \mathbb{R}^d$ and subsets $P'_1 \subset P_1, \dots, P'_{d+1} \subset P_{d+1}$, with

$$|P'_i| \geq 2^{-2^{2^{O(d)}}} n,$$

such that all closed rainbow simplices contains q .




Applications, Tverberg-type result

Theorem (Pach, 1998)

Let $P_1, P_2, \dots, P_{d+1} \subset \mathbb{R}^d$ be disjoint n -element point sets with $P_1 \cup \dots \cup P_{d+1}$ in general position. Then there is a point $q \in \mathbb{R}^d$ and subsets $P'_1 \subset P_1, \dots, P'_{d+1} \subset P_{d+1}$, with

$$|P'_i| \geq 2^{-2^{2^{O(d)}}} n,$$

such that all closed rainbow simplices contains q .




Applications, Tverberg-type result

Theorem (Pach, 1998)

Let $P_1, P_2, \dots, P_{d+1} \subset \mathbb{R}^d$ be disjoint n -element point sets with $P_1 \cup \dots \cup P_{d+1}$ in general position. Then there is a point $q \in \mathbb{R}^d$ and subsets $P'_1 \subset P_1, \dots, P'_{d+1} \subset P_{d+1}$, with

$$|P'_i| \geq 2^{-2^{2^{O(d)}}} n,$$

such that all closed rainbow simplices contains q .




Applications, Tverberg-type result

Theorem (Pach, 1998)

Let $P_1, P_2, \dots, P_{d+1} \subset \mathbb{R}^d$ be disjoint n -element point sets with $P_1 \cup \dots \cup P_{d+1}$ in general position. Then there is a point $q \in \mathbb{R}^d$ and subsets $P'_1 \subset P_1, \dots, P'_{d+1} \subset P_{d+1}$, with

$$|P'_i| \geq 2^{-2^{2^{O(d)}}} n,$$

such that all closed rainbow simplices contains q .




Applications, Tverberg-type result

Theorem (Pach, 1998)

Let $P_1, P_2, \dots, P_{d+1} \subset \mathbb{R}^d$ be disjoint n -element point sets with $P_1 \cup \dots \cup P_{d+1}$ in general position. Then there is a point $q \in \mathbb{R}^d$ and subsets $P'_1 \subset P_1, \dots, P'_{d+1} \subset P_{d+1}$, with

$$|P'_i| \geq 2^{-2^{2^{O(d)}}} n,$$

such that all closed rainbow simplices contains q .



Theorem (Pach, 1998)

$$|P'_i| \geq 2^{-2^{2^{O(d)}}} n,$$

Theorem (Fox, Pach, S., 2014)

Let $P_1, P_2, \dots, P_{d+1} \subset \mathbb{R}^d$ be disjoint n -element point sets with $P_1 \cup \dots \cup P_{d+1}$ in general position. Then there is a point $q \in \mathbb{R}^d$ and subsets $P'_1 \subset P_1, \dots, P'_{d+1} \subset P_{d+1}$, with

$$|P'_i| \geq 2^{-O(d^2 \log d)} n,$$

such that all closed rainbow simplices contains q .


Applications, Same-type Lemma

Theorem (Bárány and Valtr, 1998)

Let P_1, \dots, P_k be n -element point sets in \mathbb{R}^d such that $P_1 \cup \dots \cup P_k$ is in general position. Then there are subsets $P'_1 \subset P_1, \dots, P'_k \subset P_k$ such that the k -tuple (P'_1, \dots, P'_k) has same-type transversals and

$$|P'_i| \geq 2^{-k^{O(d)}} n,$$

for all i .




Applications, Same-type Lemma

Theorem (Bárány and Valtr, 1998)

Let P_1, \dots, P_k be n -element point sets in \mathbb{R}^d such that $P_1 \cup \dots \cup P_k$ is in general position. Then there are subsets $P'_1 \subset P_1, \dots, P'_k \subset P_k$ such that the k -tuple (P'_1, \dots, P'_k) has same-type transversals and

$$|P'_i| \geq 2^{-k^{O(d)}} n,$$

for all i .



Theorem (Bárány and Valtr, 1998)

Let P_1, \dots, P_k be n -element point sets in \mathbb{R}^d such that $P_1 \cup \dots \cup P_k$ is in general position. Then there are subsets $P'_1 \subset P_1, \dots, P'_k \subset P_k$ such that the k -tuple (P'_1, \dots, P'_k) has same-type transversals and

$$|P'_i| \geq 2^{-k^{O(d)}} n,$$

for all i .

$$2^{-k^{O(d)}} n$$




Applications, Same-type Lemma

Theorem (Bárány and Valtr, 1998)

Let P_1, \dots, P_k be n -element point sets in \mathbb{R}^d such that $P_1 \cup \dots \cup P_k$ is in general position. Then there are subsets $P'_1 \subset P_1, \dots, P'_k \subset P_k$ such that the k -tuple (P'_1, \dots, P'_k) has same-type transversals and

$$|P'_i| \geq 2^{-k^{O(d)}} n,$$

for all i .



Theorem (Bárány and Valtr, 1998)

$$|P'_i| \geq 2^{-k^{O(d)}} n,$$

Theorem (Fox, Pach, S., 2014)

Let P_1, \dots, P_k be n -element point sets in \mathbb{R}^d such that $P_1 \cup \dots \cup P_k$ is in general position. Then there are subsets $P'_1 \subset P_1, \dots, P'_k \subset P_k$ such that the k -tuple (P'_1, \dots, P'_k) has same-type transversals and

$$|P'_i| \geq 2^{-O(d^3 k \log k)} n,$$

for all i .

Sketch proof of Same-type lemma


Theorem (Fox, Pach, S., 2014)

Let P_1, \dots, P_k be n -element point sets in \mathbb{R}^d such that $P_1 \cup \dots \cup P_k$ is in general position. Then there are subsets $P'_1 \subset P_1, \dots, P'_k \subset P_k$ such that the k -tuple (P'_1, \dots, P'_k) has same-type transversals and

$$|P'_i| \geq 2^{-O(d^3 k \log k)} n,$$

for all i .

Sketch proof.



Sketch proof of Same-type lemma


Theorem (Fox, Pach, S., 2014)


Let P_1, \dots, P_k be n -element point sets in \mathbb{R}^d such that $P_1 \cup \dots \cup P_k$ is in general position. Then there are subsets $P'_1 \subset P_1, \dots, P'_k \subset P_k$ such that the k -tuple (P'_1, \dots, P'_k) has same-type transversals and

$$|P'_i| \geq 2^{-O(d^3 k \log k)} n,$$

for all i .


Sketch proof.





Goodman-Pollack: Number of different order-types of k -element point sets in d dimensions is at most $k^{O(d^2k)}$.

There exists an order type π , such that at least $k^{-O(d^2k)} n^k$ (rainbow) k -tuples have order type π .



k -partite k -uniform semi-algebraic hypergraph $H = (P_1, \dots, P_k, E)$
in d -space


$E = \{k\text{-tuples with order type } \pi\}$. $|E| \geq k^{-O(d^2 k)} n^k$.

Complexity of E ?

To check if (p_1, \dots, p_k) has order π , just check the orientation of each $(d+1)$ -tuple. For each $p_{i_1}, \dots, p_{i_{d+1}}$


$$\det \begin{pmatrix} 1 & 1 & \cdots & 1 \\ | & | & \cdots & | \\ p_{i_1} & p_{i_2} & \vdots & p_{i_{d+1}} \\ | & | & \cdots & | \end{pmatrix} \rightarrow \{+, -\}.$$

Hence we need to check $t = \binom{k}{d+1}$ polynomial inequalities.
Complexity of E is $(t, D) = \left(\binom{k}{d+1}, 1\right)$.



k -partite k -uniform semi-algebraic hypergraph $H = (P_1, \dots, P_k, E)$ in d -space.

$$\epsilon = k^{-O(d^2 k)}, t = \binom{k}{d+1}, D = 1$$



k -partite k -uniform semi-algebraic hypergraph $H = (P_1, \dots, P_k, E)$ in d -space.


$$\epsilon = k^{-O(d^2 k)}, t = \binom{k}{d+1}, D = 1$$

$$|P'_i| \geq 2^{-O(d^3 k \log k)} n,$$

Density theorem (restated)

Theorem (Fox, Pach, S., 2014)

Let $H = (V_1, \dots, V_k, E)$ be a k -partite semi-algebraic k -uniform hypergraph in d -space, where $|V_1| = \dots = |V_k| = n$ and E has complexity (t, D) . If $|E| \geq \epsilon n^k$, then there are subsets $V'_1, \dots, V'_k \subset V$ such that $|V'_i| \geq \frac{\epsilon^{d^c}}{2^{ckd^c}} n$, and $(V'_1, \dots, V'_k) \subset E$.



Future work and open problems

- 1 Find more applications.
- 2 Extend results to more complicated relations, i.e., Semi-Pfaffian, o-minimal, etc.

Thank you!