


# Density-type theorems for semi-algebraic hypergraphs

Jacob Fox, Janos Pach, Andrew Suk

March 21, 2014


# An old Ramsey-type result, Kövári, Sós, and Turán and Erdős

Bipartite graph  $G$ , edge set  $E$ .




# An old Ramsey-type result, Kövári, Sós, and Turán and Erdős

Bipartite graph  $G$ , edge set  $E$ .




# An old Ramsey-type result, Kővári, Sós, and Turán and Erdős


Bipartite graph  $G$ , edge set  $E$ .




$k$ -partite  $k$ -uniform hypergraph  $H$ , edge set  $E$ .




$k$ -partite  $k$ -uniform hypergraph  $H$ , edge set  $E$ .



$k$ -partite  $k$ -uniform hypergraph  $H$ , edge set  $E$ .




$k$ -partite  $k$ -uniform hypergraph  $H$ , edge set  $E$ .



These results are tight.




$k$ -partite  $k$ -uniform hypergraph  $H$ , edge set  $E$ .




**In this talk:** We can do much better if  $H$  is a semi-algebraic  $k$ -uniform hypergraph.

$k$ -partite  $k$ -uniform hypergraph  $H$ , edge set  $E$ .




**Semi-algebraic** hypergraphs:  $V = \{\text{simple geometric objects in } \mathbb{R}^d\}$ ,  $E = \{\text{simple relation on } k \text{ tuples of } V\}$ .


$V = \{A_1, \dots, A_n\}$ ,  $n$  disks in the plane.  $E = \{\text{pairs of disks that intersect}\}$ .




$V = \{A_1, \dots, A_n\}$ ,  $n$  disks in the plane.  $E = \{\text{pairs of disks that intersect}\}$ .



$V = \{A_1, \dots, A_n\}$ ,  $n$  disks in the plane.  $E = \{\text{pairs of disks that intersect}\}$ .




$V = \{A_1, \dots, A_n\}$ ,  $n$  disks in the plane.  $E = \{\text{pairs of disks that intersect}\}$ .



$A_i \rightarrow p_i = (x_i, y_i, r_i)$ ,  $A_j \rightarrow p_j = (x_j, y_j, r_j)$ .  $A_i$  and  $A_j$  cross if and only if

$$-x_i^2 + 2x_i x_j - x_j^2 - y_i^2 + 2y_i y_j - y_j^2 + r_i^2 + 2r_i r_j + r_j^2 \geq 0.$$

$V = \{A_1, \dots, A_n\}$ ,  $n$  disks in the plane.  $E = \{ \text{pairs of disks that intersect} \}$ .



Graph  $G = (V, E)$ ,  $V = n$  points in  $\mathbb{R}^3$

$E$  defined by the polynomial



$$f(z_1, \dots, z_6) = -z_1^2 + 2z_1z_4 - z_4^2 - z_2^2 + 2z_2z_5 - z_5^2 + z_3^2 + 2z_3z_6 + z_6^2.$$

$$(p_i, p_j) \in E \Leftrightarrow f(p_i, p_j) \geq 0.$$

# More examples of semi-algebraic hypergraphs

## Examples

- 1  $V = \{n \text{ circles in } \mathbb{R}^3\}$   
 $E = \{\text{pairs that are linked}\}.$
- 2  $V = \{n \text{ hyperplanes in } \mathbb{R}^d \text{ in general position}\},$   
 $E = \{d\text{-tuples whose intersection point is above the}$   
 $\text{hyperplane } x_d = 0\}.$







# More examples of semi-algebraic hypergraphs

## Examples

- 1  $V = \{n \text{ circles in } \mathbb{R}^3\} \rightarrow n \text{ points in higher dimensions.}$   
 $E = \{\text{pairs that are linked}\} \rightarrow \text{polynomials } f_1, \dots, f_t.$
- 2  $V = \{n \text{ hyperplanes in } \mathbb{R}^d \text{ in general position}\} \rightarrow n \text{ points in higher dimensions,}$   
 $E = \{d\text{-tuples whose intersection point is above the hyperplane } x_d = 0\} \rightarrow \text{polynomials } f_1, \dots, f_t.$



We say that  $H = (V, E)$  is a **semi-algebraic  $k$ -uniform hypergraph in  $d$ -space** if

$$V = \{n \text{ points in } \mathbb{R}^d\}$$

$E$  defined by polynomials  $f_1, \dots, f_t$  and a Boolean formula  $\Phi$  such that

$$(p_{i_1}, \dots, p_{i_k}) \in E$$

$$\Leftrightarrow \Phi(f_1(p_{i_1}, \dots, p_{i_k}) \geq 0, \dots, f_t(p_{i_1}, \dots, p_{i_k}) \geq 0) = \text{yes}$$


# Example

3-uniform hypergraph  $H = (V, E)$ ,  $V = \{p_1, \dots, p_n\}$  points in  $\mathbb{R}^d$ .

Relation  $E \subset \binom{V}{3}$  depends on  $f$  and  $\Phi$

$$\phi(f(x_1, x_2, x_3) \geq 0) = \{\text{yes, no}\}$$

$(p_1, p_2, p_3) \in E$  depends on  $f(p_1, p_2, p_3) \rightarrow \{+, -, 0\}$ .




# Example

3-uniform hypergraph  $H = (V, E)$ ,  $V = \{p_1, \dots, p_n\}$  points in  $\mathbb{R}^d$ .

Relation  $E \subset \binom{V}{3}$  depends on  $f$  and  $\Phi$

$$\phi(f(x_1, x_2, x_3) \geq 0) = \{\text{yes, no}\}$$

$(p_1, p_2, p_3) \in E$  depends on  $f(p_1, p_2, p_3) \rightarrow \{+, -, 0\}$ .




# Example

3-uniform hypergraph  $H = (V, E)$ ,  $V = \{p_1, \dots, p_n\}$  points in  $\mathbb{R}^d$ .

Relation  $E \subset \binom{V}{3}$  depends on  $f$  and  $\Phi$

$$\phi(f(x_1, x_2, x_3) \geq 0) = \{\text{yes, no}\}$$

Zero set  $f(p_1, p_2, x_3) = 0$ , surface in  $\mathbb{R}^d$ .




# Example

3-uniform hypergraph  $H = (V, E)$ ,  $V = \{p_1, \dots, p_n\}$  points in  $\mathbb{R}^d$ .

Relation  $E \subset \binom{V}{3}$  depends on  $f$  and  $\Phi$

$$\phi(f(x_1, x_2, x_3) \geq 0) = \{\text{yes, no}\}$$

Zero set  $f(p_1, p_2, x_3) = 0$ , surface in  $\mathbb{R}^d$ .




# Example

3-uniform hypergraph  $H = (V, E)$ ,  $V = \{p_1, \dots, p_n\}$  points in  $\mathbb{R}^d$ .

Relation  $E \subset \binom{V}{3}$  depends on  $f$  and  $\Phi$

$$\phi(f(x_1, x_2, x_3) \geq 0) = \{\text{yes, no}\}$$

Zero set  $f(p_1, p_2, x_3) = 0$ , surface in  $\mathbb{R}^d$ .



$E$  has complexity  $(t, D)$ , degree of  $f(p_1, p_2, x_3) \leq D$ .

# Complexity of relation $E$

$$x_i \in \mathbb{R}^d$$

$E$  has complexity  $(t, D)$

- 1 described by polynomials  $f_1, \dots, f_t$ ,
- 2 and the degree of ALL  $kt$   $d$ -variate polynomials  $f_i(x_1, \dots, x_{k-1}, x_k), f_i(x_1, \dots, x_{k-2}, x_{k-1}, x_k), \dots, f_i(x_1, x_2, \dots, x_k)$ , for  $i = 1 \dots t$ , is at most  $D$ .


Note.  $f_i$  has degree at most  $Dk$ .



# Motivation: Orientations and order-types


Our motivation,  $E$  is related to order-types and orientations.

$f(p_1, p_2, x_3)$  is linear.



$E$  has complexity  $(t, D) = (t, 1)$ .

# Example




$V = \{n \text{ points in the plane}\},$

$E = \{\text{triples having a clockwise orientation}\}.$

$H = (V, E)$  semi-algebraic 3-uniform hypergraph in the plane  
( $d = 2$ )

# Example




$V = \{n \text{ points in the plane}\},$

$E = \{\text{triples having a clockwise orientation}\}.$

$H = (V, E)$  semi-algebraic 3-uniform hypergraph in the plane

$(d = 2)$

# Example




$V = \{n \text{ points in the plane}\},$

$E = \{\text{triples having a clockwise orientation}\}.$

$H = (V, E)$  semi-algebraic 3-uniform hypergraph in the plane  
( $d = 2$ )

# Example




$V = \{n \text{ points in the plane}\},$

$E = \{\text{triples having a clockwise orientation}\}.$

$H = (V, E)$  semi-algebraic 3-uniform hypergraph in the plane

$(d = 2)$


# Example



$E = \{\text{triples having a clockwise orientation}\}.$

$$\det \begin{pmatrix} 1 & 1 & 1 \\ x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{pmatrix} > 0.$$


# Example



Complexity of  $E$  is  $(t, D)$ , where  $t = 1, D = 1$

$$\det \begin{pmatrix} 1 & 1 & 1 \\ x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{pmatrix} > 0.$$

# Example




Complexity of  $E$  is  $(t, D)$ , where  $t = 1, D = 1$

$$\det \begin{pmatrix} 1 & 1 & 1 \\ x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{pmatrix} > 0$$




# Example



Complexity of  $E$  is  $(t, D)$ , where  $t = 1, D = 1$

$$\det \begin{pmatrix} 1 & 1 & 1 \\ x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{pmatrix} > 0$$

# Example



Complexity of  $E$  is  $(t, D)$ , where  $t = 1, D = 1$

$$\det \begin{pmatrix} 1 & 1 & 1 \\ x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{pmatrix} > 0$$

# Example in higher dimensions

$E = (d + 1)$ -tuples with a positive orientation, complexity  
 $(t, D) = (1, 1)$ .

$$\det \begin{pmatrix} 1 & 1 & \cdots & 1 \\ | & | & \cdots & | \\ p_{i_1} & p_{i_2} & \vdots & p_{i_{d+1}} \\ | & | & \cdots & | \end{pmatrix} > 0.$$

# Example in higher dimensions

$E = (d + 1)$ -tuples with a positive orientation, complexity  
 $(t, D) = (1, 1)$ .

$$\det \begin{pmatrix} 1 & 1 & \cdots & 1 \\ | & | & \cdots & | \\ p_{i_1} & p_{i_2} & \vdots & p_{i_{d+1}} \\ | & | & \cdots & | \end{pmatrix} > 0.$$

# Example in higher dimensions


$E = (d + 1)$ -tuples with a positive orientation, complexity  
 $(t, D) = (1, 1)$ .

$$\det \begin{pmatrix} 1 & \mathbf{1} & \cdots & \mathbf{1} \\ | & | & \cdots & | \\ p_{i_1} & \mathbf{p}_{i_2} & \vdots & \mathbf{p}_{i_{d+1}} \\ | & | & \cdots & | \end{pmatrix} > 0.$$

# Previous results

Theorem (Alon, Pach, Pinchasi, Radoicic, Sharir 2005)


Let  $H = (V_1, V_2, E)$  be a bipartite semi-algebraic graph ( $k = 2$ ) in  $d$ -space, where  $|V_1| = |V_2| = n$  and  $E$  has complexity  $(t, D)$ . Then there are subsets  $V'_1, V'_2 \subset V$  such that  $|V'_i| \geq \epsilon n$  and either  $(V'_1, V'_2) \subset E$  or  $(V'_1, V'_2) \subset \bar{E}$ , and  $\epsilon = \epsilon(d, t, D)$ .



# Previous results

Theorem (Alon, Pach, Pinchasi, Radoicic, Sharir 2005)


Let  $H = (V_1, V_2, E)$  be a bipartite semi-algebraic graph ( $k = 2$ ) in  $d$ -space, where  $|V_1| = |V_2| = n$  and  $E$  has complexity  $(t, D)$ . Then there are subsets  $V'_1, V'_2 \subset V$  such that  $|V'_i| \geq \epsilon n$  and either  $(V'_1, V'_2) \subset E$  or  $(V'_1, V'_2) \subset \bar{E}$ , and  $\epsilon = \epsilon(d, t, D)$ .



# Previous results

Theorem (Alon, Pach, Pinchasi, Radoicic, Sharir 2005)

Let  $H = (V_1, V_2, E)$  be a bipartite semi-algebraic graph ( $k = 2$ ) in  $d$ -space, where  $|V_1| = |V_2| = n$  and  $E$  has complexity  $(t, D)$ . Then there are subsets  $V'_1, V'_2 \subset V$  such that  $|V'_i| \geq \epsilon n$  and either  $(V'_1, V'_2) \subset E$  or  $(V'_1, V'_2) \subset \bar{E}$ , and  $\epsilon = \epsilon(d, t, D)$ .






# Stronger density theorem

Including an argument of Komlos:

Theorem (Alon, Pach, Pinchasi, Radoicic, Sharir 2005)

Let  $H = (V_1, V_2, E)$  be a bipartite semi-algebraic graph ( $k = 2$ ) in  $d$ -space, where  $|V_1| = |V_2| = n$  and  $E$  has complexity  $(t, D)$ . If  $|E| \geq \epsilon n^2$ , then there are subsets  $V'_1, V'_2 \subset V$  such that  $|V'_i| \geq \epsilon^C n$  where  $C = C(d, t, D)$ , and  $(V'_1, V'_2) \subset E$ .




# Stronger density theorem

Including an argument of Komlos:

Theorem (Alon, Pach, Pinchasi, Radoicic, Sharir 2005)


Let  $H = (V_1, V_2, E)$  be a bipartite semi-algebraic graph ( $k = 2$ ) in  $d$ -space, where  $|V_1| = |V_2| = n$  and  $E$  has complexity  $(t, D)$ . If  $|E| \geq \epsilon n^2$ , then there are subsets  $V'_1, V'_2 \subset V$  such that  $|V'_i| \geq \epsilon^C n$  where  $C = C(d, t, D)$ , and  $(V'_1, V'_2) \subset E$ .



# Generalization

Theorem (Fox, Gromov, Lafforgue, Naor, Pach 2012, Bukh and Hubard 2012)


Let  $H = (V_1, \dots, V_k, E)$  be a  $k$ -partite semi-algebraic  $k$ -uniform hypergraph in  $d$ -space, where  $|V_1| = \dots = |V_k| = n$  and  $E$  has complexity  $(t, D)$ . If  $|E| \geq \epsilon n^k$ , then there are subsets  $V'_1, \dots, V'_k \subset V$  such that  $|V'_i| \geq \epsilon^C n$  where  $C = C(k, d, t, D)$ , and  $(V'_1, \dots, V'_k) \subset E$ .



# Generalization

Theorem (Fox, Gromov, Lafforgue, Naor, Pach 2012, Bukh and Hubard 2012)

Let  $H = (V_1, \dots, V_k, E)$  be a  $k$ -partite semi-algebraic  $k$ -uniform hypergraph in  $d$ -space, where  $|V_1| = \dots = |V_k| = n$  and  $E$  has complexity  $(t, D)$ . If  $|E| \geq \epsilon n^k$ , then there are subsets  $V'_1, \dots, V'_k \subset V$  such that  $|V'_i| \geq \epsilon^C n$  where  $C = C(k, d, t, D)$ , and  $(V'_1, \dots, V'_k) \subset E$ .




# Generalization

$C(k, d, t, D)$ : Dependency on uniformity  $k$  is very bad.

Fox, Gromov, Lafforgue, Naor, Pach:  $C(k, d, t, D) \sim \underbrace{2^{2^{\dots 2^d}}}_k$

(tower-type)

Bukh-Hubard:  $C(k, d, t, D) \sim 2^{2^{k+d}}$ , double exponential in  $k$ .




Bukh-Hubard: Set sizes decay triple exponentially in  $k$

# New results

For simplicity, complexity  $(t, D)$  is fixed.


Theorem (Fox, Pach, Suk, 2013+)

Let  $H = (V_1, \dots, V_k, E)$  be a  $k$ -partite semi-algebraic  $k$ -uniform hypergraph in  $d$ -space, where  $|V_1| = \dots = |V_k| = n$  and  $E$  has complexity  $(t, D)$ . If  $|E| \geq \epsilon n^k$ , then there are subsets  $V'_1, \dots, V'_k \subset V$  such that  $|V'_i| \geq \frac{\epsilon^{d^c}}{2^{ckd^c}} n$ , and  $(V'_1, \dots, V'_k) \subset E$ .




## Theorem (Fox, Pach, Suk, 2013+)

Let  $H = (V_1, \dots, V_k, E)$  be a  $k$ -partite semi-algebraic  $k$ -uniform hypergraph in  $d$ -space, where  $|V_1| = \dots = |V_k| = n$  and  $E$  has complexity  $(t, D)$ . If  $|E| \geq \epsilon n^k$ , then there are subsets  $V'_1, \dots, V'_k \subset V$  such that  $|V'_i| \geq \frac{\epsilon^{d^c}}{2^{ckd^c}} n$ , and  $(V'_1, \dots, V'_k) \subset E$ .



## Theorem (Fox, Pach, Suk, 2013+)

Let  $H = (V_1, \dots, V_k, E)$  be a  $k$ -partite semi-algebraic  $k$ -uniform hypergraph in  $d$ -space, where  $|V_1| = \dots = |V_k| = n$  and  $E$  has complexity  $(t, 1)$ . If  $|E| \geq \epsilon n^k$ , then there are subsets  $V'_1, \dots, V'_k \subset V$  such that  $|V'_i| \geq \frac{\epsilon^{d+1}}{2^{ckd \log d}} n$ , and  $(V'_1, \dots, V'_k) \subset E$ .






# Applications, Tverberg-type result

## Theorem (Pach, 1998)

Let  $P_1, P_2, \dots, P_{d+1} \subset \mathbb{R}^d$  be disjoint  $n$ -element point sets with  $P_1 \cup \dots \cup P_{d+1}$  in general position. Then there is a point  $q \in \mathbb{R}^d$  and subsets  $P'_1 \subset P_1, \dots, P'_{d+1} \subset P_{d+1}$ , with

$$|P'_i| \geq 2^{-2^{2^{O(d)}}} n,$$

such that all closed rainbow simplices contains  $q$ .




# Applications, Tverberg-type result

## Theorem (Pach, 1998)

Let  $P_1, P_2, \dots, P_{d+1} \subset \mathbb{R}^d$  be disjoint  $n$ -element point sets with  $P_1 \cup \dots \cup P_{d+1}$  in general position. Then there is a point  $q \in \mathbb{R}^d$  and subsets  $P'_1 \subset P_1, \dots, P'_{d+1} \subset P_{d+1}$ , with

$$|P'_i| \geq 2^{-2^{2^{O(d)}}} n,$$

such that all closed rainbow simplices contains  $q$ .




# Applications, Tverberg-type result

## Theorem (Pach, 1998)

Let  $P_1, P_2, \dots, P_{d+1} \subset \mathbb{R}^d$  be disjoint  $n$ -element point sets with  $P_1 \cup \dots \cup P_{d+1}$  in general position. Then there is a point  $q \in \mathbb{R}^d$  and subsets  $P'_1 \subset P_1, \dots, P'_{d+1} \subset P_{d+1}$ , with

$$|P'_i| \geq 2^{-2^{2^{O(d)}}} n,$$

such that all closed rainbow simplices contains  $q$ .




# Applications, Tverberg-type result

## Theorem (Pach, 1998)

Let  $P_1, P_2, \dots, P_{d+1} \subset \mathbb{R}^d$  be disjoint  $n$ -element point sets with  $P_1 \cup \dots \cup P_{d+1}$  in general position. Then there is a point  $q \in \mathbb{R}^d$  and subsets  $P'_1 \subset P_1, \dots, P'_{d+1} \subset P_{d+1}$ , with

$$|P'_i| \geq 2^{-2^{2^{O(d)}}} n,$$

such that all closed rainbow simplices contains  $q$ .




# Applications, Tverberg-type result

## Theorem (Pach, 1998)

Let  $P_1, P_2, \dots, P_{d+1} \subset \mathbb{R}^d$  be disjoint  $n$ -element point sets with  $P_1 \cup \dots \cup P_{d+1}$  in general position. Then there is a point  $q \in \mathbb{R}^d$  and subsets  $P'_1 \subset P_1, \dots, P'_{d+1} \subset P_{d+1}$ , with

$$|P'_i| \geq 2^{-2^{2^{O(d)}}} n,$$

such that all closed rainbow simplices contains  $q$ .




# Applications, Tverberg-type result

## Theorem (Pach, 1998)

Let  $P_1, P_2, \dots, P_{d+1} \subset \mathbb{R}^d$  be disjoint  $n$ -element point sets with  $P_1 \cup \dots \cup P_{d+1}$  in general position. Then there is a point  $q \in \mathbb{R}^d$  and subsets  $P'_1 \subset P_1, \dots, P'_{d+1} \subset P_{d+1}$ , with

$$|P'_i| \geq 2^{-2^{2^{O(d)}}} n,$$

such that all closed rainbow simplices contains  $q$ .



Theorem (Pach, 1998)

$$|P'_i| \geq 2^{-2^{2^{O(d)}}} n,$$

Theorem (Fox, Pach, Suk, 2013+)

Let  $P_1, P_2, \dots, P_{d+1} \subset \mathbb{R}^d$  be disjoint  $n$ -element point sets with  $P_1 \cup \dots \cup P_{d+1}$  in general position. Then there is a point  $q \in \mathbb{R}^d$  and subsets  $P'_1 \subset P_1, \dots, P'_{d+1} \subset P_{d+1}$ , with

$$|P'_i| \geq 2^{-O(d^3 \log d)} n,$$

such that all closed rainbow simplices contains  $q$ .


# Applications, Same-type Lemma

Theorem (Bárány and Valtr, 1998)

Let  $P_1, \dots, P_k$  be  $n$ -element point sets in  $\mathbb{R}^d$  such that  $P_1 \cup \dots \cup P_k$  is in general position. Then there are subsets  $P'_1 \subset P_1, \dots, P'_k \subset P_k$  such that the  $k$ -tuple  $(P'_1, \dots, P'_k)$  has same-type transversals and

$$|P'_i| \geq 2^{-k^{O(d)}} n,$$

for all  $i$ .






# Applications, Same-type Lemma

Theorem (Bárány and Valtr, 1998)

Let  $P_1, \dots, P_k$  be  $n$ -element point sets in  $\mathbb{R}^d$  such that  $P_1 \cup \dots \cup P_k$  is in general position. Then there are subsets  $P'_1 \subset P_1, \dots, P'_k \subset P_k$  such that the  $k$ -tuple  $(P'_1, \dots, P'_k)$  has same-type transversals and

$$|P'_i| \geq 2^{-k^{O(d)}} n,$$

for all  $i$ .



## Theorem (Bárány and Valtr, 1998)

Let  $P_1, \dots, P_k$  be  $n$ -element point sets in  $\mathbb{R}^d$  such that  $P_1 \cup \dots \cup P_k$  is in general position. Then there are subsets  $P'_1 \subset P_1, \dots, P'_k \subset P_k$  such that the  $k$ -tuple  $(P'_1, \dots, P'_k)$  has same-type transversals and

$$|P'_i| \geq 2^{-k^{O(d)}} n,$$

for all  $i$ .

$$2^{-k^{O(d)}} n$$




# Applications, Same-type Lemma

## Theorem (Bárány and Valtr, 1998)

Let  $P_1, \dots, P_k$  be  $n$ -element point sets in  $\mathbb{R}^d$  such that  $P_1 \cup \dots \cup P_k$  is in general position. Then there are subsets  $P'_1 \subset P_1, \dots, P'_k \subset P_k$  such that the  $k$ -tuple  $(P'_1, \dots, P'_k)$  has same-type transversals and

$$|P'_i| \geq 2^{-k^{O(d)}} n,$$

for all  $i$ .



Theorem (Bárány and Valtr, 1998)

$$|P'_i| \geq 2^{-k^{O(d)}} n,$$

Theorem (Fox, Pach, Suk, 2013+)

*Let  $P_1, \dots, P_k$  be  $n$ -element point sets in  $\mathbb{R}^d$  such that  $P_1 \cup \dots \cup P_k$  is in general position. Then there are subsets  $P'_1 \subset P_1, \dots, P'_k \subset P_k$  such that the  $k$ -tuple  $(P'_1, \dots, P'_k)$  has same-type transversals and*

$$|P'_i| \geq 2^{-O(d^3 k \log k)} n,$$

*for all  $i$ .*

# Sketch proof of Same-type lemma


Theorem (Fox, Pach, Suk, 2013+)

Let  $P_1, \dots, P_k$  be  $n$ -element point sets in  $\mathbb{R}^d$  such that  $P_1 \cup \dots \cup P_k$  is in general position. Then there are subsets  $P'_1 \subset P_1, \dots, P'_k \subset P_k$  such that the  $k$ -tuple  $(P'_1, \dots, P'_k)$  has same-type transversals and

$$|P'_i| \geq 2^{-O(d^3 k \log k)} n,$$

for all  $i$ .

**Sketch proof.**



# Sketch proof of Same-type lemma


Theorem (Fox, Pach, Suk, 2013+)


Let  $P_1, \dots, P_k$  be  $n$ -element point sets in  $\mathbb{R}^d$  such that  $P_1 \cup \dots \cup P_k$  is in general position. Then there are subsets  $P'_1 \subset P_1, \dots, P'_k \subset P_k$  such that the  $k$ -tuple  $(P'_1, \dots, P'_k)$  has same-type transversals and

$$|P'_i| \geq 2^{-O(d^3 k \log k)} n,$$

for all  $i$ .


**Sketch proof.**





Goodman-Pollack: Number of different order-types of  $k$ -element point sets in  $d$  dimensions is at most  $k^{O(d^2k)}$ .

There exists an order type  $\pi$ , such that at least  $k^{-O(d^2k)} n^k$  (rainbow)  $k$ -tuples have order type  $\pi$ .



$k$ -partite  $k$ -uniform semi-algebraic hypergraph  $H = (P_1, \dots, P_k, E)$   
in  $d$ -space

$E = \{k\text{-tuples with order type } \pi\}$ .  $|E| \geq k^{-O(d^2 k)} n^k$ .


Complexity of  $E$ ?



To check if  $(p_1, \dots, p_k)$  has order  $\pi$ , just check the orientation of each  $(d+1)$ -tuple. For each  $p_{i_1}, \dots, p_{i_{d+1}}$


$$\det \begin{pmatrix} 1 & 1 & \cdots & 1 \\ | & | & \cdots & | \\ p_{i_1} & p_{i_2} & \vdots & p_{i_{d+1}} \\ | & | & \cdots & | \end{pmatrix} \rightarrow \{+, -\}.$$

Hence we need to check  $t = \binom{k}{d+1}$  polynomial inequalities.  
Complexity of  $E$  is  $(t, D) = (\binom{k}{d+1}, 1)$ .



$k$ -partite  $k$ -uniform semi-algebraic hypergraph  $H = (P_1, \dots, P_k, E)$  in  $d$ -space.

$$\epsilon = k^{-O(d^2 k)}, t = \binom{k}{d+1}, D = 1$$



$k$ -partite  $k$ -uniform semi-algebraic hypergraph  $H = (P_1, \dots, P_k, E)$  in  $d$ -space.


$$\epsilon = k^{-O(d^2 k)}, t = \binom{k}{d+1}, D = 1$$

$$|P'_i| \geq 2^{-O(d^3 k \log k)} n,$$


## Theorem (Fox, Pach, Suk, 2013+)

Let  $H = (V_1, \dots, V_k, E)$  be a  $k$ -partite semi-algebraic  $k$ -uniform hypergraph in  $d$ -space, where  $|V_1| = \dots = |V_k| = n$  and  $E$  has complexity  $(t, D) = (1, 1)$ . If  $|E| \geq \epsilon n^k$ , then there are subsets  $V'_1, \dots, V'_k \subset V$  such that  $|V'_i| \geq \frac{\epsilon^{d+1}}{2^{ckd \log d}} n$ , and  $(V'_1, \dots, V'_k) \subset E$ .


**Proof.** Induction on  $k$ .  $E$  depends on  $f(\mathbb{R}^d, \mathbb{R}^d, \dots, \mathbb{R}^d) \geq 0$ .




$$V_0 = V_2 \times V_3 \times \cdots \times V_k \subset \mathbb{R}^{(k-1)d}. \quad |V_0| = n^{k-1}.$$




Bipartite graph  $G = (V_1, V_0, E)$ ,  $E$  depends on  
 $f(\mathbb{R}^d, \mathbb{R}^{(d-1)k}) \geq 0$ .



$$V_1 \subset \mathbb{R}^d.$$




For each  $p \in V_0 \subset \mathbb{R}^{(d-1)k}$







For each  $p \in V_0 \subset \mathbb{R}^{(d-1)k}$ , hyperplane  $f(\mathbb{R}^d, p) = 0$  in  $\mathbb{R}^d$ .




$p' \in V_0 \subset \mathbb{R}^{(d-1)k}$ , hyperplane  $f(\mathbb{R}^d, p') = 0$  in  $\mathbb{R}^d$ .




$H = \{n^{k-1} \text{ hyperplanes in } \mathbb{R}^d\}.$




**Cutting Lemma (Chazelle, Friedman 1990).** For  $r > 0$ ,  
 Subdivide  $\mathbb{R}^d$  into at most  $2^{10d \log d} r^d$  simplices, such that at most  
 $n^{k-1}/r$  hyperplanes from  $H$  crosses each cell.




At most  $\frac{n^{k-1}}{r}$  hyperplanes crosses  $\Delta$ .  $f(\mathbb{R}^d, p) = 0$




Most hyperplanes  $f(\mathbb{R}^d, p) = 0$  do not cross  $\Delta$ .



If hyperplane  $f(\mathbb{R}^d, p) = 0$  does not cross  $\Delta$ , then sign pattern does not change.




If hyperplane  $f(\mathbb{R}^d, \rho) = 0$  does not cross  $\Delta$ , then sign pattern does not change.







Divided  $\mathbb{R}^d$  into  $2^{10d \log d} r^d$  cells, on average a cell has  $\frac{n}{2^{10d \log d} r^d}$  points.




If a cell has fewer than  $\frac{n}{2^{10d \log d} r^d} (\epsilon/2)$  points, DELETE all edges emanating out of it. Still have a  $(\epsilon/2)n^k$  edges




Cells with edges has at least  $\frac{n}{2^{10d \log d} r^d} (\epsilon/2)$  points. Still have a  $(\epsilon/2)n^k$  edges




Each “big” cell gives rise to a certain number of vertices in  $V_0$  that is adjacent to all points in it.



Each “big” cell gives rise to a certain number of vertices in  $V_0$  that is adjacent to all points in it.




Each “big” cell gives rise to a certain number of vertices in  $V_0$  that is adjacent to all points in it.




Since  $|E'| \geq (\epsilon/2)n^k$  edges,

Set  $r \sim 8/\epsilon$ , at least  $\frac{\epsilon}{8}n^{k-1}$  vertices in  $V_0$  adjacent to all vertices in  $\Delta$ .




For  $r \sim 8/\epsilon$ . At least  $(\epsilon/2) \frac{n}{2^{10d \log d} r^d} = \frac{\epsilon^{d+1}}{2^{cd \log d}} n$  points inside  $\Delta$ .






$H'$   $(k - 1)$ -partite  $(k - 1)$ -uniform hypergraphs with density  $\epsilon/8$ .  
 Apply induction hypothesis.




$H'$   $(k - 1)$ -partite  $(k - 1)$ -uniform hypergraphs with density  $\epsilon/8$ .  
 Apply induction hypothesis.




$E'$  are  $k - 1$  tuples adjacent to all vertices in  $\Delta$ , and that gives rise to a hyperplane  $f(R^d, p) = 0$  that does not cross  $\Delta$ .

$H'$   $(k - 1)$ -partite  $(k - 1)$ -uniform hypergraphs with density  $\epsilon/8$ .  
Apply induction hypothesis.



$E' \rightarrow f(p, \mathbb{R}^{(k-1)d}) \geq 0$ , and if hyperplane  $f(\mathbb{R}^d, p_2, \dots, p_k) = 0$  crosses  $\Delta$ .


$H'$   $(k - 1)$ -partite  $(k - 1)$ -uniform hypergraphs with density  $\epsilon/8$ .  
Apply induction hypothesis.



Find desired parts  $V'_1, \dots, V'_k$

$$|V'_i| \geq \frac{\epsilon^{d+1}}{2^{cd \log d}} n$$

Found a complete  $k$ -partite  $k$ -uniform hypergraph.



Find desired parts  $V'_1, \dots, V'_k$

$$|V'_i| \geq \frac{\epsilon^{d+1}}{2^{ckd \log d}} n$$

Regularity lemma:  $H = (P, E)$  semi-algebraic  $k$ -uniform hypergraph in  $\mathbb{R}^d$ .

Theorem (Fox, Pach, Suk, 2013+)

For any  $\epsilon > 0$ , we can partition  $P$  into at most  $M(\epsilon)$  parts, such that almost all  $k$ -tuples of parts are **complete or empty**. Moreover  $M(\epsilon) < (1/\epsilon)^c$ , where  $c$  depends only on  $k, d, E$ .

Usual regularity: almost all  $k$ -tuples of parts are "**random**".  $M(\epsilon)$  is huge:

- $k = 2$ ,  $M(\epsilon) \leq \text{tower}(1/\epsilon) = 2^{2^{\dots^2}}$
- $k = 3$ ,  $M(\epsilon) \leq \text{wowzer}(1/\epsilon) = \text{tower}(\text{tower}(\dots(\text{tower}(2))))$
- $k = 4$ ,  $M(\epsilon) \leq \text{wowzer}(\text{wowzer}(\dots(\text{wowzer}(2))))$ .

# Future work and open problems

- 1 Find more applications.
- 2 Extend results to more complicated relations, i.e., Semi-Pfaffian, o-minimal, etc.

**Thank you!**