

# 1 Definition

## 1.1 Functor of points

Suppose we have a scheme  $G$  over  $\mathbb{k}$ , then a  $\mathbb{k}$ -point is just a map  $\text{Spec } \mathbb{k} \rightarrow G$ . Thus we can think of the set  $G(\mathbb{k})$  as  $\text{Hom}(\text{Spec } \mathbb{k}, G)$ . Generalizing this, for any scheme  $G$  we can define a functor of points

$$h_G : (\text{Affine schemes}/\mathbb{k})^{\text{op}} \rightarrow \text{Sets}, \quad X \mapsto \text{Mor}_{\mathbb{k}}(X, G)$$

and Yoneda's lemma says that the functor  $G \mapsto h_G$  is fully faithful, i.e., a scheme is determined up to isomorphism by its functor of points. Now, we say that a functor

$$F : (\text{Affine schemes}/\mathbb{k})^{\text{op}} \rightarrow \text{Sets}$$

is **representable** if it is isomorphic to  $h_G$  for some scheme  $G$ .

**Theorem 1.1.** *Such a functor is  $F$  representable if and only if  $F$  admits an open cover by representable functors and  $F$  is a sheaf with respect to the Zariski topology on the category of schemes.*

**Note 1.1.** The motivation for this topology comes from gluing sheaves. We say that a functor  $F : \text{Sch}^{\text{op}} \rightarrow \text{Sets}$  satisfies the sheaf property if for every scheme  $T$  and every open covering  $T = \bigcup_{\alpha} U_{\alpha}$  we have an exact complex:

$$0 \rightarrow F(T) \rightarrow \prod_{\alpha} F(U_{\alpha}) \rightrightarrows \prod_{\alpha, \beta} F(U_{\alpha} \times_T U_{\beta})$$

**Example 1.2.** The functor  $X \mapsto H^0(X, \mathcal{O}_X)$  is represented by  $\mathbb{A}^1$ . The functor  $X \mapsto H^0(X, \mathcal{O}_X)^*$  is represented by  $\mathbb{G}_m$ .

For a group scheme  $G$  we just ask  $h_G$  to factor, i.e.,

$$h_G : \begin{array}{ccc} (\text{Sch}/\mathbb{k})^{\text{op}} & \xrightarrow{\quad\quad\quad} & \text{Sets} \\ & \searrow & \nearrow \\ & \text{Grps} & \end{array}$$

so we can think of a group scheme  $G$  over  $\mathbb{k}$  as a functor  $(\text{Sch}/\mathbb{k})^{\text{op}} \rightarrow \text{Grps}$ .

## 1.2 Jacobian functor

Let  $C$  be a complete nonsingular curve over  $\mathbb{k}$ . Recall that a Weil divisor is just a formal sum of points

$$D = \sum_{j=1}^n n_j P_j, \quad \deg D = \sum_{j=1}^n n_j [\mathbb{k}(P_j) : \mathbb{k}]$$

and we have a correspondence between divisors and line bundles on  $C$ . We defined  $\text{Pic}^0(C)$  to be the group of degree 0 line bundles on  $C$ ; this is not necessarily a scheme.

Let  $T$  be a connected scheme over  $\mathbb{k}$ , look at the fiber product

$$\pi : C \times_{\mathbb{k}} T \rightarrow T, \quad C_t = \pi^{-1}(t)$$

and for  $\mathcal{L} \in \text{Pic}(C \times_{\mathbb{k}} T)$  we define  $\mathcal{L}_t = \mathcal{L}|_{C_t}$ . Then we have that the map  $t \mapsto \chi(C_t, \mathcal{L}_t)$  is locally constant (this is an example of a flat family of curves). By Riemann Roch, this implies that  $\deg(\mathcal{L}_t)$  is independent of  $t \in T$ . This degree is also invariant under base change, so we can define a functor  $J : (\text{Sch}/\mathbb{k})^{\text{op}} \rightarrow \text{Grps}$ ,

$$J(T) = \left\{ \mathcal{L} \in \text{Pic}(C \times_{\mathbb{k}} T) \mid \deg(\mathcal{L}_t) = 0 \forall t \in T \right\} / \pi^* \text{Pic}(T)$$

and we can think of  $h_J(T)$  as the group of degree 0 line bundles on  $C$  parametrized by  $T$ , modulo the trivial family. Notice that  $J(\mathbb{k}) = \text{Pic}^0(C)$ .

**Definition 1.3.** If  $J$  is representable, then we call the representative scheme  $\text{Jac}(C)$ .

### 1.3 Obstruction to representability

Suppose  $J$  is representable by a group scheme  $\text{Jac}(C)$ , and let  $K/\mathbb{k}$  be a Galois extension with group  $\Gamma$ . Then

$$J(K) = \text{Mor}_{\mathbb{k}}(\text{Spec } K, \text{Jac}(C)) \simeq \text{Mor}_K(\text{Spec } K, \text{Jac}(C) \times_{\mathbb{k}} K)$$

**Note 1.2.** Let's convince myself of the affine case, i.e., to show  $\text{Hom}_{\mathbb{k}}(A, K) = \text{Hom}_K(A \otimes_{\mathbb{k}} K, K)$ . This comes from the fact that tensor product is a pushout, i.e., we have a diagram

$$\begin{array}{ccc}
 & & K \\
 & \swarrow & \uparrow \text{Id} \\
 & & K \\
 & \swarrow & \uparrow \\
 A \otimes_{\mathbb{k}} K & \longleftarrow & K \\
 \uparrow & & \uparrow \\
 A & \longleftarrow & \mathbb{k}
 \end{array}$$

Here (on  $K$ ) we have a Galois action by  $\Gamma$ . Since

$$\text{Mor}_K(\text{Spec } K, \text{Jac}(C) \times_{\mathbb{k}} K)^{\Gamma} \simeq \text{Mor}_{\mathbb{k}}(\text{Spec } \mathbb{k}, \text{Jac}(C))$$

**Note 1.3.** Once again, easy to prove for affine case. The Galois action on  $\text{Spec } A \times_{\mathbb{k}} K \simeq \text{Spec}(A \otimes_{\mathbb{k}} K)$  is just  $1 \otimes \sigma$  for  $\sigma \in \Gamma$ .

we have that  $J(K)^{\Gamma} = J(\mathbb{k})$ . In other words, we would expect

$$\text{Pic}^0(C \times_{\mathbb{k}} K)^{\Gamma} = \text{Pic}^0(C)$$

but this is not true in general. In fact, we can measure the failure of this equality by an exact sequence

$$0 \rightarrow \text{Pic}(C) \rightarrow \text{Pic}(C \times_{\mathbb{k}} K)^{\Gamma} \rightarrow \text{Br}(\mathbb{k})$$

where  $\text{Br}(\mathbb{k})$  is the Brauer group of  $\mathbb{k}$ .

**Example 1.4.** Consider  $C = V(x^2 + y^2 + z^2) \in \mathbb{P}_{\mathbb{R}}^2$ , which is empty. Now,  $C \times_{\mathbb{R}} \mathbb{C}$  is a conic in  $\mathbb{P}_{\mathbb{C}}^2$ , hence isomorphic to  $\mathbb{P}_{\mathbb{C}}^1$ . If  $\text{Pic}(C) = \text{Pic}(C \times_{\mathbb{R}} \mathbb{C})^{\mathbb{Z}/2\mathbb{Z}}$  then  $\text{Pic}(C)$  is a subgroup of index at most 2 in  $\text{Pic}(C \times_{\mathbb{R}} \mathbb{C}) = \mathbb{Z}$ , but this is impossible.

The possible issue here is that a line bundle  $\mathcal{L} \in \text{Pic}(C \times_{\mathbb{k}} K)^\Gamma$  has too many automorphisms (coming from  $\Gamma$ ), and they have to satisfy some compatible conditions for  $\mathcal{L}$  to descend to  $\text{Pic}(C)$ . Fortunately, if  $C(\mathbb{k})$  is nonempty then everything works.

**Theorem 1.5.** *Suppose  $C$  has a  $\mathbb{k}$ -point. Then the functor  $J$  can be represented by a group scheme  $\text{Jac}(C)$ , called the Jacobian variety of  $C$ .*

The idea here is that if we include the  $\mathbb{k}$ -point in our data, then we kill all the automorphisms. The forgetful functor getting rid of the extra data is actually an isomorphism, so we are good.

**Example 1.6.**  $\text{Jac}(\mathbb{P}^1) = \text{Spec } \mathbb{k}$ , since there is no nontrivial divisor of degree 0 ( $\text{Pic}(\mathbb{P}^1) = \mathbb{Z}$ , two points are linearly equivalent). The Jacobian of an elliptic curve is isomorphic to the elliptic curve itself.

**Example 1.7.** Let  $C$  be a projective curve over  $\mathbb{F}_p$ , and  $p$  a  $\mathbb{F}_p$ -point of  $C$ . Then  $C \setminus \{p\}$  is affine, and the class group of its coordinate ring is  $J(\mathbb{F}_p)$ . The reason is that  $\text{Pic}(C \setminus \{p\}) = \text{Pic}^0(C)$  by mapping  $D \mapsto D - \deg D \cdot p$ .

## 2 Properties and applications

Clearly,  $J = \text{Jac}(C)$  is a nonsingular abelian variety.

**Proposition 2.1.** *The tangent space  $T_0J$  is canonically isomorphic to  $H^1(C, \mathcal{O}_C)$ . Thus the dimension of  $J$  is equal to the genus of  $C$ .*

**Definition 2.2.** For each point  $p \in C(\mathbb{k})$  we can define a map  $f_p : C \rightarrow \text{Jac}(C)$  such that at the level of  $\mathbb{k}$ -points,

$$f_p : C(\mathbb{k}) \rightarrow \text{Jac}(C)(\mathbb{k}) = \text{Pic}^0(C), \quad x \mapsto [x - p]$$

**Proposition 2.3.** *The map  $f_p^* : H^0(J, \Omega_J) \rightarrow H^0(C, \Omega_C)$  is an isomorphism.*

*Proof.* Essentially we need to show that the following diagram commutes

$$\begin{array}{ccc} H^0(J, \Omega_J) & \xrightarrow{f_p^*} & H^0(C, \Omega_C) \\ \cong \downarrow & & \uparrow \cong \\ (T_0J)^\vee & \xrightarrow{\cong} & H^1(C, \mathcal{O}_C)^\vee \end{array}$$

□

**Note 2.1.** What is this map  $H^0(J, \Omega_J) \simeq (T_0J)^\vee$ ? It's just evaluating the 1-form at 0; the idea is that a group variety is homogeneous, so a vector  $X_0$  in  $T_0J$  extends uniquely to a vector field  $X$  hence we get an isomorphism.

$$H^0(J, \Omega_J) \ni \omega \mapsto (X_0 \mapsto \omega_0(X_0))$$

**Proposition 2.4.** *The map  $f_p$  is a closed embedding.*

*Proof.* Field extensions are faithfully flat, so it suffices prove this for the case  $\mathbb{k} = \overline{\mathbb{k}}$ . Then we just need to show that the map separates points and tangents. For points, suppose  $f_p(x) = f_p(y)$  then  $[x - p] = [y - p]$  which implies  $x, y$  are linearly equivalent, but this is impossible on a curve of genus  $> 0$ .  $\square$

Now consider the map:

$$f_p^r : C^r \rightarrow J, \quad (p_1, \dots, p_r) \mapsto [p_1 + \dots + p_r - r \cdot p]$$

which descends to a map  $f_p^{(r)} : C^{(r)} \rightarrow J$ . The image  $W^r = f_p^{(r)}(C^{(r)})$  is a closed subvariety of  $J$ , and thus  $W^g = J$ .

**Note 2.2.** Abel's theorem says that fibers of  $f_p^{(r)}$  correspond to linear equivalence classes of effective divisors of degree  $r$ .

**Theorem 2.5.** *For all  $r \leq g$ , the map  $f_p^{(r)} : C^{(r)} \rightarrow W^r$  is birational. In particular,  $J$  is the unique abelian variety birational to  $C^{(g)}$ .*

**Example 2.6.** Consider a curve  $C$  of genus 2. We have a double cover (by the canonical divisor)  $\pi : C \rightarrow \mathbb{P}^1$  branched at 6 points. Each fiber  $\pi^{-1}(x) = \{p, q\}$  (not necessarily distinct) defines a degree 2 divisor  $p + q$ . Since any 2 points on  $\mathbb{P}^1$  are linearly equivalent, all these degree 2 divisors are linearly equivalent and get mapped to the same point by  $f^{(2)}$ .

So we have a family of degree 2 divisors (which is itself a divisor in  $C^{(2)}$ ) which gets contracted in  $J(C)$ . In other words,  $f^{(2)}$  is a blow down here.

Now let  $\Theta = W^{g-1}$  then this is a divisor in  $J$ . This does depend on the chosen point  $p$ , but only up to translation. Such a divisor induces a map:

$$\phi_{\mathcal{L}(\Theta)} : J \rightarrow J^\vee, \quad x \mapsto [t_x^* \mathcal{L}(\Theta) \otimes \mathcal{L}(\Theta)^{-1}]$$

which is an isomorphism in this case. Hence  $(A, \Theta)$  is a principally polarized abelian variety.

**Theorem 2.7** (Torelli).  *$C$  is determined, up to isomorphism, by its principally polarized Jacobian variety.*