

OPTIMAL SYNTACTIC DEFINITIONS OF BACK-AND-FORTH TYPES

RUIYUAN CHEN, DAVID GONZALEZ, AND MATTHEW HARRISON-TRAINOR

ABSTRACT. The back-and-forth relations $\mathcal{M} \leq_\alpha \mathcal{N}$ are central to computable structure theory and countable model theory. It is well-known that the relation $\{(\mathcal{M}, \mathcal{N}) : \mathcal{M} \leq_\alpha \mathcal{N}\}$ is (lightface) $\Pi_{2\alpha}^0$. We show that this is optimal as the set is $\Pi_{2\alpha}^0$ -complete. We are also interested in the one-sided relations $\{\mathcal{N} : \mathcal{M} \leq_\alpha \mathcal{N}\}$ and $\{\mathcal{N} : \mathcal{M} \geq_\alpha \mathcal{N}\}$ for a fixed \mathcal{M} , measuring the Π_α and Σ_α types of \mathcal{M} . We show that these sets are always $\Pi_{\alpha+2}^0$ and $\Pi_{\alpha+3}^0$ respectively, and that for most α there are structures \mathcal{M} for which these relations are complete at that level. In particular, there are structures \mathcal{M} such that there is no Π_α (or even $\Pi_{\alpha+1}$) sentence φ such that $\mathcal{N} \models \varphi \iff \mathcal{M} \leq_\alpha \mathcal{N}$.

This is unfortunate as not all $\Pi_{\alpha+2}$ sentences are preserved under \leq_α . We define a new hierarchy of syntactic complexity closely related to the back-and-forth game, which can both define the back-and-forth types as well as be preserved by them. These hierarchies of formulas have already been useful in certain Henkin constructions, one of which we give in this paper, and another previously used by Gonzalez and Harrison-Trainor to show that every Π_α theory of linear orders has a model with Scott rank at most $\alpha + 3$.

1. INTRODUCTION

Originally used to prove that any two countable dense linear orders without endpoints are isomorphic, the back-and-forth method is a classical tool in mathematical logic dating back over one hundred years. A general viewpoint, beginning with Ehrenfeucht [Ehr61] and Fraïssé [Fra50], is that by playing back-and-forth games between two first-order structures \mathcal{M} and \mathcal{N} we can capture which sets of formulas they agree on. Generally two players, Spoiler and Duplicator, take turns playing elements from the two structures, with Duplicator trying to maintain atomic facts between the elements in \mathcal{M} and the elements in \mathcal{N} . In the Ehrenfeucht–Fraïssé game most commonly used in model theory, both players play single elements, and (in a finite language) the second player wins the game of length n if and only if the two structures \mathcal{M} and \mathcal{N} satisfy the same sentences with quantifier depth n . Other versions, such as pebble games, can be used for finite variable logics.

In this paper, we work with the back-and-forth game that corresponds to infinitary logic, and our main goal is to understand the complexity of deciding which player has a winning strategy. The key features of this particular back-and-forth game are that (a) on their turn, players play tuples of arbitrary size with the size chosen by Spoiler, (b) Spoiler must alternate between playing in \mathcal{M} and playing in \mathcal{N} , with Duplicator alternating between playing in \mathcal{N} and playing in \mathcal{M} , and (c) the games are played over a possibly infinite ordinal clock.

This back-and-forth game yields the back-and-forth relations. (For simplicity, we assume that all languages in this paper are relational.)

Definition 1.1. *The standard asymmetric back-and-forth relations \leq_α , for a countable ordinal $\alpha < \omega_1$, are defined by:*

2020 *Mathematics Subject Classification.* 03C15, 03C57, 03D45, 03E15, 03C75.

The first author acknowledges support from the National Science Foundation under Grant No. DMS-2224709. The third author acknowledges support from a Sloan Research Fellowship and from the National Science Foundation under Grant No. DMS-2153823.

- $(\mathcal{M}, \bar{a}) \leq_0 (\mathcal{N}, \bar{b})$ if \bar{a} and \bar{b} satisfy the same quantifier-free formulas from among the first $|\bar{a}|$ -many formulas.
- For $\alpha > 0$, $(\mathcal{M}, \bar{a}) \leq_\alpha (\mathcal{N}, \bar{b})$ if for each $\beta < \alpha$ and $\bar{d} \in \mathcal{N}$ there is $\bar{c} \in \mathcal{M}$ such that $(\mathcal{N}, \bar{b}\bar{d}) \leq_\beta (\mathcal{M}, \bar{a}\bar{c})$.

We define $\bar{a} \equiv_\alpha \bar{b}$ if $\bar{a} \leq_\alpha \bar{b}$ and $\bar{b} \leq_\alpha \bar{a}$.

The interpretation of $(\mathcal{M}, \bar{a}) \leq_\alpha (\mathcal{N}, \bar{b})$ is that in the back-and-forth game between \mathcal{M} and \mathcal{N} , starting with the partial isomorphism $\bar{a} \mapsto \bar{b}$ and with the first player **Spoiler** to play next in \mathcal{N} , the second player **Duplicator** can play without losing along an ordinal clock α . The content of the classical back-and-forth argument is then that if **Duplicator** can continue playing forever, and \mathcal{M} and \mathcal{N} are countable, then $\mathcal{M} \cong \mathcal{N}$.

These back-and-forth relations are related to the infinitary logic $\mathcal{L}_{\omega_1\omega}$ which extends the standard finitary first-order logic by allowing countably infinite disjunctions and conjunctions. The formulas of $\mathcal{L}_{\omega_1\omega}$ are stratified in terms of their complexity as measured by counting the number of alternations between existential and universal quantifiers. For an ordinal $\alpha < \omega_1$:

- A formula is Σ_0 and Π_0 if it is finitary quantifier-free.
- A formula is Σ_α if it is of the form

$$\bigvee_i \exists \bar{x}_i \psi_i(\bar{x}_i)$$

where each ψ_i is Π_β for some $\beta < \alpha$.

- A formula is Π_α if it is of the form

$$\bigwedge_i \forall \bar{x}_i \psi_i(\bar{x}_i)$$

where each ψ_i is Π_β for some $\beta < \alpha$.

The back-and-forth relations \leq_α characterize when two structures \mathcal{M} and \mathcal{N} satisfy the same Σ_α/Π_α sentences as in the following theorem of Karp [Kar65] (see, e.g., [Mon], Theorem II.36).

Theorem 1.2. *For any non-zero ordinal α , structures \mathcal{M} and \mathcal{N} and tuples $\bar{a} \in \mathcal{M}$ and $\bar{b} \in \mathcal{N}$, the following are equivalent:*

- (1) $(\mathcal{M}, \bar{a}) \leq_\alpha (\mathcal{N}, \bar{b})$.
- (2) Every Π_α formula true about \bar{a} in \mathcal{M} is true about \bar{b} in \mathcal{N} .
- (3) Every Σ_α formula true about \bar{b} in \mathcal{N} is true about \bar{a} in \mathcal{M} .

Both these back-and-forth relations and the infinitary logic $\mathcal{L}_{\omega_1\omega}$ have proven to be very useful in countable structure theory, the study of countable structures using tools from model theory, descriptive set theory, and computability theory.

This paper is about the complexity of characterizing when it is that $\mathcal{M} \leq_\alpha \mathcal{N}$. One might hope, because the α -back-and-forth relations are about preservation of Σ_α/Π_α formulas, that there would be a characterization which is also on the level of α -many alternations of quantifiers. While interesting in its own right, this question is motivated by certain appearances of the back-and-forth relations in arguments in countable model theory. Before giving our main results, we will talk about some of the motivation behind this question to set the stage.

Our main motivation has to do with Scott's isomorphism theorem. Given a countable \mathcal{L} -structure \mathcal{M} , the set of \mathcal{L} -structures isomorphic to \mathcal{M} is naively an analytic set in the Polish space $\text{Mod}(\mathcal{L})$ of countably infinite \mathcal{L} -structures. It is a surprising result of Scott [Sco65] that this set is actually always Borel and thus any countable structure admits a characterization up to isomorphism. Scott's argument uses the back-and-forth relations: He shows that for any structure \mathcal{M} there is a countable ordinal α such that for any countable structure \mathcal{N} ,

$$\mathcal{M} \leq_\alpha \mathcal{N} \implies \mathcal{M} \cong \mathcal{N}.$$

From this, one checks that the set $\{\mathcal{N} : \mathcal{M} \leq_\alpha \mathcal{N}\}$ is $\mathbf{\Pi}_{2\alpha}^0$ by writing out the definition of the back-and-forth relations—each step of the inductive definition uses two quantifiers—and concludes that $\{\mathcal{N} : \mathcal{M} \cong \mathcal{N}\}$ is $\mathbf{\Pi}_{2\alpha}^0$ and hence Borel. (The fact that the back-and-forth relations \leq_α have a $\mathbf{\Pi}_{2\alpha}$ definition is well-known and is used in various contexts. It appears as Exercise VIII.6 in [Mon] and underlies, for example, Nadel’s [Nad74] observation that a computable structure has a computable Scott sentence if and only if it has computable Scott rank.)

For a given \mathcal{M} , the Borel complexity of this set $\{\mathcal{N} : \mathcal{M} \cong \mathcal{N}\}$, as measured by Wadge reducibility, is called the *Scott complexity* of \mathcal{M} [AGHTT21]. By the Lopez-Escobar theorem [LE66, Vau75], the Scott complexity of $\{\mathcal{N} : \mathcal{M} \cong \mathcal{N}\}$ is the same as the least complexity of a Scott sentence for \mathcal{M} , that is, a sentence φ of the infinitary logic $\mathcal{L}_{\omega_1\omega}$ that characterizes \mathcal{M} up to isomorphism among countable structures: for any countable structure \mathcal{N} ,

$$\mathcal{M} \models \varphi \implies \mathcal{N} \cong \mathcal{M}.$$

Thus in the context of the mentioned proof of Scott’s isomorphism theorem, $\{\mathcal{N} : \mathcal{M} \cong \mathcal{N}\}$ is $\mathbf{\Pi}_{2\alpha}^0$ and so is defined by a $\mathbf{\Pi}_{2\alpha}$ sentence of $\mathcal{L}_{\omega_1\omega}$. This gives an upper bound of $\mathbf{\Pi}_{2\alpha}$ on the Scott complexity of \mathcal{M} , but it is far from the lower bound of $\mathbf{\Pi}_\alpha$.¹

In this paper, we ask whether this upper bound can be improved, and if so by how much? We work not just with Scott complexity but also with the problem of defining the back-and-forth relations \leq_α in general. Though we generally state our results in terms of structures, considering sets such as $\{\mathcal{N} : \mathcal{M} \leq_\alpha \mathcal{N}\}$, by naming constants our results also apply to tuples, e.g., $\{(\mathcal{N}, \bar{b}) : (\mathcal{M}, \bar{a}) \leq_\alpha (\mathcal{N}, \bar{b})\}$. Such a set represents the $\mathbf{\Pi}_\alpha$ -type of \bar{a} in \mathcal{M} , and these types are important objects in countable structure theory.

We show that the $\mathbf{\Pi}_{2\alpha}^0$ definition is the best definition of \leq_α in terms of *both* of the structures \mathcal{M} and \mathcal{N} , that is, the set $\{(\mathcal{M}, \mathcal{N}) : \mathcal{M} \leq_\alpha \mathcal{N}\}$ is $\mathbf{\Pi}_{2\alpha}^0$ -complete for most languages and α . Fixing one of the structures, this gives a “schematic” $\mathbf{\Pi}_{2\alpha}^0$ definition of the sets $\{\mathcal{N} : \mathcal{M} \leq_\alpha \mathcal{N}\}$ and $\{\mathcal{N} : \mathcal{M} \geq_\alpha \mathcal{N}\}$ where we replace quantifiers over \mathcal{M} by infinitary conjunctions and disjunctions. However, this is not the best possible definition of these sets; we show that there are simpler $\mathbf{\Pi}_{\alpha+2}^0$ and $\mathbf{\Pi}_{\alpha+3}^0$ definitions for $\{\mathcal{N} : \mathcal{M} \leq_\alpha \mathcal{N}\}$ and $\{\mathcal{N} : \mathcal{M} \geq_\alpha \mathcal{N}\}$ respectively. These simpler definitions are not “schematic” in \mathcal{M} and are non-effective. The exact complexities are given by the following theorem and are proved to be best possible. There are exceptional cases when α is too small or too close to a limit ordinal, but for the most part we have the same bounds.

Theorem 1.3. *For any language containing a relation symbol of arity ≥ 2 ,² the set*

$$\{(\mathcal{M}, \mathcal{N}) : \mathcal{M} \leq_\alpha \mathcal{N}\}$$

is $\mathbf{\Pi}_{2\alpha}^0$ -complete. Moreover:

(1) *The sets*

$$\{\mathcal{N} : \mathcal{M} \leq_\alpha \mathcal{N}\}$$

have the following complexities Γ depending on α , and there is a structure \mathcal{M} such that the set is Γ -complete.

(a) *For $\alpha = 1$, $\Gamma = \mathbf{\Pi}_1^0 = \mathbf{\Pi}_\alpha^0$.*

(b) *For $\alpha = \lambda$ a limit ordinal, $\Gamma = \mathbf{\Pi}_\lambda^0 = \mathbf{\Pi}_\alpha^0$.*

¹We also note that this is related to the variations between different notions of Scott rank. There have been many different non-equivalent definitions of Scott rank, some of them using the back-and-forth relations, and some using the complexity of Scott sentences. See [Mon15] for a survey.

²Any language with a binary relation symbol can effectively bi-interpret any theory in any language. In particular, these maps preserve properties regarding back-and-forth relations, so such languages exhibit all properties exhibited in any language. See [Mon21] Section VI.3.2 and [Mon] Theorem XI.7 for more details.

- (c) For $\alpha = \lambda + 1$ the successor of a limit ordinal, $\Gamma = \mathbf{\Pi}_{\lambda+2}^0 = \mathbf{\Pi}_{\alpha+1}^0$.
 (d) For $\alpha = \beta + 2$, $\Gamma = \mathbf{\Pi}_{\alpha+2}^0$.
 (2) The sets

$$\{\mathcal{N} : \mathcal{M} \geq_\alpha \mathcal{N}\}$$

have the following complexities Γ depending on α , and there is a structure \mathcal{M} such that the set is Γ -complete.

- (a) For $\alpha = 1$, $\Gamma = \mathbf{\Pi}_2^0 = \mathbf{\Pi}_{\alpha+1}^0$.
 (b) For $\alpha = 2$, $\Gamma = \mathbf{\Pi}_3^0 = \mathbf{\Pi}_{\alpha+1}^0$.
 (c) For $\alpha = \lambda$ a limit ordinal, $\Gamma = \mathbf{\Pi}_\lambda^0 = \mathbf{\Pi}_\alpha^0$.
 (d) For $\alpha = \lambda + 1$ the successor of a limit ordinal, $\Gamma = \mathbf{\Pi}_{\lambda+2}^0 = \mathbf{\Pi}_{\alpha+1}^0$.
 (e) For $\alpha = \lambda + 2$ the double successor of a limit ordinal, $\Gamma = \mathbf{\Pi}_{\lambda+4}^0 = \mathbf{\Pi}_{\alpha+2}^0$.
 (f) For $\alpha = \beta + 3$, $\Gamma = \mathbf{\Pi}_{\alpha+3}^0$.

The first part of the paper is devoted to proving this theorem. We note that while $\{(\mathcal{M}, \mathcal{N}) : \mathcal{M} \leq_\alpha \mathcal{N}\}$ is (lightface) $\mathbf{\Pi}_{2\alpha}^0$ when α is computable, the other bounds where \mathcal{M} is fixed are not generally effective in \mathcal{M} . That is, for example, if \mathcal{M} is computable, we do not know whether $\{\mathcal{N} : \mathcal{M} \leq_n \mathcal{N}\}$ is (lightface) $\mathbf{\Pi}_{n+2}^0$. In [ACHT] Alvir, Csima, and Harrison-Trainor undertake an analysis of the computability of Scott sentences for computable structures with $\mathbf{\Pi}_2$ Scott sentences; it is shown that a computable structure \mathcal{M} can have a $\mathbf{\Pi}_2$ Scott sentence but no computable Σ_4 Scott sentence. This implies that for this particular \mathcal{M} the set $\{\mathcal{N} : \mathcal{M} \leq_2 \mathcal{N}\}$ is (boldface) $\mathbf{\Pi}_2^0$ but not (lightface) Σ_4^0 .

Question 1.4. *Assuming that \mathcal{M} is computable, what is the lightface complexity of the sets from Theorem 1.3?*

The fact that $2 + 2 = 2 \cdot 2$ means that in the case $\alpha = 2$ the complexities $\mathbf{\Pi}_{2\alpha}^0$ and $\mathbf{\Pi}_{\alpha+2}^0$ are the same, and so we get no evidence about the behavior in general.

In the second part of the paper, we will consider the following question. What class of $\mathcal{L}_{\omega_1\omega}$ formulas corresponds *exactly* with the back-and-forth relations? Namely, we want a class of formulas Γ_α that has the properties:

- (1) for each \mathcal{M} and α there is a sentence $\varphi \in \Gamma_\alpha$ such that

$$\mathcal{N} \models \varphi \implies \mathcal{M} \leq_\alpha \mathcal{N},$$

- (2) if $\mathcal{M} \leq_\alpha \mathcal{N}$, and $\mathcal{M} \models \varphi$, then $\mathcal{N} \models \varphi$.

While $\mathbf{\Pi}_\alpha$ formulas satisfy (2), above we saw that one requires $\mathbf{\Pi}_{\alpha+2}$ formulas for (1). We will introduce a new hierarchy of formulas, which we call the $\mathfrak{A}_\alpha/\mathfrak{E}_\alpha$ hierarchy, which is better suited for characterizing the back-and-forth relations. The definition is somewhat complicated, but the essence is that when counting quantifiers, $\forall \mathbb{W} \mathbb{A} \exists$ counts only as two alternations of quantifiers, whereas in the $\Sigma_\alpha/\mathbf{\Pi}_\alpha$ hierarchy, this would count for four alternations of quantifiers. There are also associated $\overline{\mathfrak{E}}_\alpha$ and $\overline{\mathfrak{A}}_\alpha$ formulas which are the closure, under \mathbb{A} and \mathbb{W} , of the \mathfrak{A}_α and \mathfrak{E}_α formulas. We leave the full definition to Section 6. We prove that these classes of formulas satisfy (1) and (2) above:

Proposition 1.5. *For \mathcal{M} a countable structure, $\bar{a} \in \mathcal{M}$, and $\alpha \geq 1$, there are a $\overline{\mathfrak{E}}_\alpha$ formula $\varphi_{\bar{a}, \mathcal{M}, \alpha}(\bar{x})$ and \mathfrak{A}_α formula $\psi_{\bar{a}, \mathcal{M}, \alpha}(\bar{x})$ such that for \mathcal{N} any structure,*

$$\mathcal{N} \models \varphi_{\bar{a}, \mathcal{M}, \alpha}(\bar{b}) \iff (\mathcal{M}, \bar{a}) \geq_\alpha (\mathcal{N}, \bar{b})$$

and

$$\mathcal{N} \models \psi_{\bar{a}, \mathcal{M}, \alpha}(\bar{b}) \iff (\mathcal{M}, \bar{a}) \leq_\alpha (\mathcal{N}, \bar{b}).$$

Proposition 1.6. *Suppose that $(\mathcal{M}, \bar{a}) \leq_\alpha (\mathcal{N}, \bar{b})$ for $\alpha \geq 1$. Then given a $\bar{\mathfrak{E}}_\alpha$ formula $\varphi(\bar{x})$ and a $\bar{\mathfrak{A}}_\alpha$ formula $\psi(\bar{x})$,*

$$\mathcal{N} \models \varphi(\bar{b}) \implies \mathcal{M} \models \varphi(\bar{a})$$

and

$$\mathcal{M} \models \psi(\bar{a}) \implies \mathcal{N} \models \psi(\bar{b})$$

At first, we thought that these classes of formulas were only a curiosity, but they have proved very useful. For example, they appear naturally in the proof of the upper bounds in 1(d) and 2(d) of Theorem 1.3. More importantly, they have been useful for certain Henkin constructions. The most important example of this is in [GHT], where the proof of the main result is a Henkin construction using formulas of complexity \mathfrak{E}_α . We will discuss briefly in Section 7.1 the role that \mathfrak{E}_α formulas play in that construction. Finally, in Section 7.2, we give a Henkin construction using \mathfrak{E}_α formulas to give an improved version of Montalbán's type omitting theorem for infinitary logic [Mon15]. This type omitting theorem was originally due to Gonzalez but was unpublished; we give a new and easier proof here making use of the $\bar{\mathfrak{A}}_\alpha/\mathfrak{E}_\alpha$ formulas.

Theorem 1.7. *Let \mathcal{M} be a countable structure. Let $(\Gamma_i)_{i \in \omega}$ be a list of Π_α -types which are not Σ_α -supported in \mathcal{M} . Let φ be a $\Pi_{\alpha+1}$ sentence true in \mathcal{M} . Then there is $\mathcal{N} \leq_\alpha \mathcal{M}$ such that $\mathcal{N} \models \varphi$ and \mathcal{N} omits all of the Γ_i .*

This adds the additional condition that $\mathcal{N} \leq_\alpha \mathcal{M}$ to Montalbán's type omitting theorem. In particular, we maintain the entire Σ_α theory of \mathcal{M} . It is still open whether we can have $\mathcal{N} \geq_\alpha \mathcal{M}$ or even stronger $\mathcal{N} \equiv_\alpha \mathcal{M}$.

In Section 7.2, we state several consequences of this type-omitting theorem such as the following.

Corollary 1.8. *Let \mathcal{M} be a countable structure.*

(1) *If, for all countable \mathcal{N} ,*

$$\mathcal{M} \leq_\alpha \mathcal{N} \implies \mathcal{M} \cong \mathcal{N}$$

then \mathcal{M} has a $\Pi_{\alpha+2}$ Scott sentence.

(2) *If, for all countable \mathcal{N} ,*

$$\mathcal{N} \leq_\alpha \mathcal{M} \implies \mathcal{M} \cong \mathcal{N}$$

then \mathcal{M} has a $\Pi_{\alpha+1}$ Scott sentence.

Item (1) simply follows from our general characterization of $\{\mathcal{N} : \mathcal{M} \leq_\alpha \mathcal{N}\}$ as a $\mathbf{\Pi}_{\alpha+2}^0$ set for any α . For (2), such an argument would yield a $\Pi_{\alpha+3}$ Scott sentence. Using the type omitting theorem, we improve this to a $\Pi_{\alpha+1}$ Scott sentence, and this cannot be further improved; for example, take \mathcal{M} to be the infinite-dimensional \mathbb{Q} -vector space. This has a Π_3 Scott sentence and no simpler, but if $\mathcal{V} \leq_2 \mathbb{Q}^{\mathbb{N}}$ then $\mathcal{V} \cong \mathbb{Q}^{\mathbb{N}}$. Whether (1) is optimal remains open.

Question 1.9. *Let \mathcal{M} be a countable structure. Suppose that for all countable \mathcal{N}*

$$\mathcal{M} \leq_\alpha \mathcal{N} \implies \mathcal{M} \cong \mathcal{N}.$$

Must \mathcal{M} have a $\Pi_{\alpha+1}$ Scott sentence? A Π_α Scott sentence?

2. THE DEFINABILITY OF THE BACK-AND-FORTH RELATIONS IN THE Σ AND Π HIERARCHIES

We begin by demonstrating the lower bound of $\mathbf{\Pi}_{2n}^0$ in the symmetrical case of the theorem for finite values. In other words, we demonstrate that the set $\{(\mathcal{M}, \mathcal{N}) : \mathcal{M} \leq_n \mathcal{N}\}$ is sometimes as difficult to define as possible. Note that $\mathbf{\Pi}_{2n}^0$ matches the classical upper bound, as the definition of the back-and-forth game gives a $\mathbf{\Pi}_{2n}^0$ description of the back-and-forth relations. (We expand on this point regarding the classical upper bound in the proof below.)

Lemma 2.1. *For any language \mathcal{L} containing a relation symbol of arity ≥ 2 , the set*

$$\{(\mathcal{M}, \mathcal{N}) : \mathcal{M} \leq_n \mathcal{N}\}$$

is Π_{2n}^0 -complete in $\text{Mod}(\mathcal{L}) \times \text{Mod}(\mathcal{L})$.

Proof. Any language containing a relation symbol of arity ≥ 2 can simulate, via effective bi-interpretation, any other language. Thus, in showing completeness we can work with whichever language is most useful.

The fact that this set is Π_{2n}^0 has been observed previously (see, e.g., Lemma VI.14 in [Mon]). To be a bit more explicit, the Π_{2n}^0 definition follows directly from the definition of the back-and-forth relations. In the product space, $\text{Mod}(\mathcal{L}) \times \text{Mod}(\mathcal{L})$, a universal quantifier over the elements of the first coordinate (or second coordinate) naturally corresponds to a countable intersection of sets. Dually, an existential quantifier naturally corresponds to a countable union. The definition of \leq_n alternates quantifiers (some over the first structure, some over the second structure) $2n$ -times and begins with a universal quantifier, so it gives a Π_{2n}^0 definition.

We focus the rest of the proof on showing that the set is Π_{2n}^0 -hard. We will give an inductive construction but we need a stronger statement to carry out the induction. For $n \geq 2$, given a Π_{2n-2}^0 set $U \subseteq 2^\omega \times \omega$ and $x \in 2^\omega$ we will construct structures $\mathcal{A}^{x,U}$ and $\mathcal{B}_i^{x,U}$ such that

- (1) if $(x, i) \in U$ then $\mathcal{A}^{x,U} \geq_n \mathcal{B}_i^{x,U}$, and
- (2) if $(x, i) \notin U$ then $\mathcal{B}_i^{x,U} \not\leq_{n-1} \mathcal{A}^{x,U}$.

For a given value of n , the language of these structures will consist of equivalence relations $\mathbb{E}_n, \dots, \mathbb{E}_3$ and unary relations $R_{i,m}^{n'}$ for $2 \leq n' \leq n$ and $i, m \in \omega$. The equivalence relation \mathbb{E}_i will refine \mathbb{E}_j if $i < j$. Inductively, in constructing these structures for $n+1$, we may assume that we know how to construct them for n .

Our base case is $n=2$. We assume that we work relative to an oracle that makes a given Π_2^0 set Π_2^0 and note that our proof relativizes; we make a similar assumption as needed throughout this proof. Given a Π_2^0 set $V \subseteq 2^\omega \times \omega$, $x \in 2^\omega$, and $k \in \omega$ we must construct structures $\mathcal{A}^{x,V}$ and $\mathcal{B}_i^{x,V}$ with the desired properties. Write

$$(x, i) \in V \iff \forall m \exists j (x, i, m, j) \in U,$$

where U is computable. We will have unary relations $R_{i,m}^2$. Let $\mathcal{A}^{x,V}$ have infinitely many elements $a_{i',m,j}^s$. The elements $a_{i',m,j}^s$ will behave the same for fixed values of i', m, j . In other words, each element is infinitely replicated by the index s . Put $a_{i',m,j}^s \in R_{i',m}^2$ if $(x, i', m, j) \in U$. Let $\mathcal{B}_i^{x,V}$ have infinitely many elements $b_{i',m,j}^s$ as well as elements b_m^* . Put $b_{i',m,j}^s \in R_{i',m}^2$ if $(x, i', m, j) \in U$ and put $b_m^* \in R_{i,m}^2$. Informally, $\mathcal{B}_i^{x,V}$ is just the same as $\mathcal{A}^{x,V}$ except for the fact that there is an additional element added to $R_{i,m}^2$ for each m . Then:

- If $(x, i) \in V$, then we can check that $\mathcal{A}^{x,V} \cong \mathcal{B}_i^{x,V}$ and hence $\mathcal{A}^{x,V} \geq_2 \mathcal{B}_i^{x,V}$. We see that $\mathcal{A}^{x,V}$ embeds into $\mathcal{B}_i^{x,V}$. The elements not in the image are the b_m^* . For each m there is j_m such that $(x, i, m, j_m) \in U$ and so b_m^* satisfies the same relations as each a_{i,m,j_m}^s (for each s). Thus we can map, say $b_m^* \leftrightarrow a_{i,m,j_m}^0$ and $b_{i,m,j}^s \leftrightarrow a_{i,m,j}^{s+1}$.
- If $(x, i) \notin V$ then there is m such that for all j we have $(x, i, m, j) \notin U$. Note that in $\mathcal{B}_i^{x,V}$ there is always an element b_m^* satisfying $R_{i,m}^2$. In contrast, in $\mathcal{A}^{x,V}$, any $a_{i',m',j}$ satisfying $R_{i',m'}^2$ must have $i' = i$ and $m' = m$ and $(x, i, m, j) \in U$. This cannot happen by the choice of i and m , and so we have an existential formula true in $\mathcal{B}_i^{x,V}$ which is not true in $\mathcal{A}^{x,V}$. Thus $\mathcal{B}_i^{x,V} \not\leq_1 \mathcal{A}^{x,V}$.

This finishes the case of $n=2$.

Now, suppose that we have a construction for n , and we will give a construction for $n + 1$. Let V be a Π_{2n}^0 set, say

$$x \in V \iff \forall m \exists j (x, i, m, j) \in U$$

where U is Π_{2n-2}^0 . For each i, m , let

$$U_{i,m} = \{(x, j) : (x, i, m, j) \in U\}.$$

By the inductive hypothesis, we have structures $\mathcal{A}^{x, U_{i,m}}$ and $\mathcal{B}_j^{x, U_{i,m}}$ such that

- (1) if $(x, j) \in U_{i,m}$ then $\mathcal{A}^{x, U_{i,m}} \geq_n \mathcal{B}_j^{x, U_{i,m}}$, and
- (2) if $(x, j) \notin U_{i,m}$ then $\mathcal{B}_j^{x, U_{i,m}} \not\geq_{n-1} \mathcal{A}^{x, U_{i,m}}$.

To be explicit, the above definitions and hypotheses immediately give that

$$(x, i) \in V \iff \forall m \exists j (x, j) \in U_{i,m}.$$

Therefore,

$$(x, i) \in V \implies \forall m \exists j \mathcal{A}^{x, U_{i,m}} \geq_n \mathcal{B}_j^{x, U_{i,m}} \text{ and } (x, i) \notin V \implies \exists m \forall j \mathcal{A}^{x, U_{i,m}} \not\geq_{n-1} \mathcal{B}_j^{x, U_{i,m}}.$$

Define $\mathcal{A}^{x, V}$ and $\mathcal{B}_i^{x, V}$ as follows:

- $\mathcal{A}^{x, V}$ is split into sorts indexed by i, m by the relations $R_{i,m}^{n+1}$. Each sort is, in turn, split into equivalence classes with respect to the equivalence relation \mathbb{E}_{n+1} with each equivalence class being the domain of one of the structures from the previous inductive case. The (i, m) sort has infinitely many equivalence classes: For each $j \in \omega$ there are infinitely many classes containing a copy of $\mathcal{B}_j^{x, U_{i,m}}$ for each $j \in \omega$, and there are no other equivalence classes.
- $\mathcal{B}_i^{x, V}$ is similarly split into sorts indexed by i, m by the relations $R_{i,m}^{n+1}$. Again, each sort is split into equivalence classes with respect to the equivalence relation \mathbb{E}_{n+1} , and each equivalence class is the domain of a structure. The (i', m) sort has infinitely many equivalence classes of the following types: For each $j \in \omega$ there are infinitely many equivalence classes containing a copy of $\mathcal{B}_j^{x, U_{i',m}}$, and if $i' = i$, there are also infinitely many equivalence classes containing a copy of $\mathcal{A}^{x, U_{i,m}}$; there are no other equivalence classes.

Informally, $\mathcal{B}_i^{x, V}$ is just the same as $\mathcal{A}^{x, V}$ except for the fact that there are infinitely many additional copies of $\mathcal{A}^{x, U_{i,m}}$ added to $R_{i,m}^{n+1}$ for each m . We now demonstrate that the $\mathcal{A}^{x, V}$ and $\mathcal{B}_i^{x, V}$ have Properties (1) and (2) required to continue the induction.

Sublemma 2.1.1. *Let \mathcal{A} and \mathcal{B} be structures in the relational language $\mathcal{L} \cup \{\mathbb{E}\}$ which consist of an equivalence relation \mathbb{E} with infinitely many equivalence classes, such that each equivalence class is an \mathcal{L} -structure (and no relations from \mathcal{L} hold between different equivalence classes). Write $\mathcal{A} = \sqcup_i \mathcal{A}_i$ and $\mathcal{B} = \sqcup_i \mathcal{B}_i$ as disjoint unions of equivalence classes, with the \mathcal{A}_i and \mathcal{B}_i being \mathcal{L} -structures. Suppose furthermore that each isomorphism type that appears does so infinitely many times. If the following conditions hold, then $(\mathcal{A}, \bar{a}) \geq_n (\mathcal{B}, \bar{b})$:*

- (A) For each k, ℓ , $a_k \mathbb{E} a_\ell$ if and only if $b_k \mathbb{E} b_\ell$.
- (B) Given (A), we may reorder \bar{a} and \bar{b} into subtuples $\bar{a}_1, \dots, \bar{a}_\ell$ and $\bar{b}_1, \dots, \bar{b}_\ell$ such that there are i_1, \dots, i_ℓ and j_1, \dots, j_ℓ with $\bar{a}_k \in \mathcal{A}_{i_k}$ and $\bar{b}_k \in \mathcal{B}_{j_k}$. Then for $k = 1, \dots, \ell$,

$$(\mathcal{A}_{i_k}, \bar{a}_k) \geq_n (\mathcal{B}_{j_k}, \bar{b}_k).$$

- (C) For each i there is j such that $\mathcal{A}_i \geq_n \mathcal{B}_j$.
- (D) For each j there is i such that $\mathcal{B}_j \geq_{n-1} \mathcal{A}_i$.

Proof. We argue inductively on n , starting with the base case of $n = 0$. This case is immediate. Supposing that (A), (B), and (C) hold, we see that these determine for each type of atomic formula that it is true of \bar{a} in \mathcal{A} if and only if it is true of \bar{b} in \mathcal{B} .

Now we inductively assume we know the sublemma for n , and prove it for $n + 1$. We argue that if (A), (B), (C), and (D) hold for $n + 1$, then $(\mathcal{A}, \bar{a}) \geq_{n+1} (\mathcal{B}, \bar{b})$. Suppose that we are given $\bar{a}' \in \mathcal{A}$. First, since (A) holds, we may break up \bar{a} and \bar{b} as in (B) into tuples $\bar{a}_1, \dots, \bar{a}_\ell$ and $\bar{b}_1, \dots, \bar{b}_\ell$. We may break \bar{a}' up into tuples $\bar{a}'_1, \dots, \bar{a}'_\ell$ such that $\bar{a}'_k \in \mathcal{A}_{i_k}$, and further elements $\bar{a}_1, \dots, \bar{a}_s^*$ such that $\bar{a}_t^* \in \mathcal{A}_{i_t^*}$. By (B), for each $k = 1, \dots, \ell$ we have $(\mathcal{A}_{i_k}, \bar{a}_k) \geq_{n+1} (\mathcal{B}_{i_k}, \bar{b}_k)$ and so we may pick $\bar{b}'_k \in \mathcal{B}_{i_k}$ such that $(\mathcal{B}_{i_k}, \bar{b}_k \bar{b}'_k) \geq_n (\mathcal{A}_{i_k}, \bar{a}_k \bar{a}'_k)$. By (C), for each $t = 1, \dots, s$ we may choose j_t^* and $\bar{b} \in \mathcal{B}_{j_t^*}$ such that $(\mathcal{B}_{j_t^*}, \bar{b}_t^*) \geq_n (\mathcal{A}_{i_t^*}, \bar{a}_t^*)$. Furthermore, as each isomorphism type appears infinitely many times, we may choose each of the j_t^* to be an index that we have not yet chosen. Our goal is to argue that

$$(\mathcal{B}, \bar{b}_1 \bar{b}'_1, \dots, \bar{b}_k \bar{b}'_k, \bar{b}_1^*, \dots, \bar{b}_s^*) \geq_n (\mathcal{A}, \bar{a}_1 \bar{a}'_1, \dots, \bar{a}_k \bar{a}'_k, \bar{a}_1^*, \dots, \bar{a}_s^*)$$

from which we can conclude that

$$(\mathcal{A}, \bar{a}) = (\mathcal{A}, \bar{a}_1, \dots, \bar{a}_k) \geq_{n+1} (\mathcal{B}, \bar{b}_1, \dots, \bar{b}_k) = (\mathcal{B}, \bar{b}).$$

To show the former, we will use the inductive hypothesis. (A) and (B) follow by choice of the \bar{b}'_i and \bar{b}_i^* . (C) for n is exactly (D) for $n + 1$, and (D) follows from (C) as $\mathcal{A}_i \geq_{n+1} \mathcal{B}_j$ implies $\mathcal{A}_i \leq_n \mathcal{B}_j$. \square

Claim 2.1.2. *If $(x, i) \in V$ then $\mathcal{A}^{x, V} \geq_{n+1} \mathcal{B}_i^{x, V}$.*

Proof. $\mathcal{A}^{x, V}$ and $\mathcal{B}_i^{x, V}$ are of the form specified by the previous claim. Furthermore, each sort indexed by i', m is of the specified form, so we may check that the conditions of the previous claim are satisfied sort by sort. If $i' \neq i$, the (i', m) th sort of $\mathcal{A}^{x, V}$ and $\mathcal{B}_i^{x, V}$ are isomorphic by construction, so there is nothing non-trivial to check. Consider instead when $i' = i$. (A) and (B) are vacuously true since there are no tuples. (C) is due to the fact that each structure $\mathcal{B}_j^{x, U_{i, m}}$ appearing as an equivalence class in $\mathcal{A}^{x, V}$ also appears as an equivalence class in $\mathcal{B}_i^{x, V}$. (D) is verified as follows. Within the (i, m) th sort, the structures appearing as equivalence classes in $\mathcal{B}_i^{x, V}$ are the $\mathcal{B}_j^{x, U_{i, m}}$, which are also equivalence classes of $\mathcal{A}^{x, V}$, and also the $\mathcal{A}^{x, U_{i, m}}$. For each m , since $(x, i) \in V$, there is j such that $(x, i, m, j) \in U$, so that $(x, j) \in U_{i, m}$. Thus, for each m there is j such that $\mathcal{A}^{x, U_{i, m}} \geq_n \mathcal{B}_j^{x, U_{i, m}}$. Since we have checked (A), (B), (C), and (D) of the previous claim, we conclude that $\mathcal{A}^{x, V} \geq_{n+1} \mathcal{B}_i^{x, V}$. \square

Sublemma 2.1.3. *Let \mathcal{A} and \mathcal{B} be structures in the relational language $\mathcal{L} \cup \{\mathbb{E}\}$ which consist of an equivalence relation \mathbb{E} with infinitely many equivalence classes, such that each equivalence class is an \mathcal{L} -structure (and no relations from \mathcal{L} hold between different equivalence classes). Write $\mathcal{A} = \sqcup_i \mathcal{A}_i$ and $\mathcal{B} = \sqcup_i \mathcal{B}_i$ as disjoint unions of equivalence classes, with the \mathcal{A}_i and \mathcal{B}_i being \mathcal{L} -structures. If there is some \mathcal{B}_j such that for every \mathcal{A}_i we have $\mathcal{A}_i \not\geq_{n-1} \mathcal{B}_j$, then $\mathcal{B} \not\geq_n \mathcal{A}$.*

Proof. Suppose to the contrary that $\mathcal{B} \geq_n \mathcal{A}$. Choose $\bar{b} \in \mathcal{B}_j$. There is $\bar{a} \in \mathcal{A}$ such that $(\mathcal{A}, \bar{a}) \geq_{n-1} (\mathcal{B}, \bar{b})$. But, as $\mathcal{A} \not\geq_{n-1} \mathcal{B}$, this yields a contradiction. \square

Claim 2.1.4. *If $(x, i) \notin V$ then $\mathcal{B}_i^{x, V} \not\geq_n \mathcal{A}^{x, V}$.*

Proof. Since $(x, i) \notin V$, there is m such that for all j we have $(x, i, m, j) \notin U$ and so $(x, j) \notin U_{i, m}$. Thus, there is m such that for all j we have $\mathcal{B}_j^{x, U_{i, m}} \not\geq_{n-1} \mathcal{A}^{x, U_{i, m}}$. For this m , we will show that Sublemma 2.1.3 applies when focusing on the (i, m) th sort of $\mathcal{B}_i^{x, V}$ and $\mathcal{A}^{x, V}$

$\mathcal{A}^{x,U_{i,m}}$ is one of the equivalence classes of $\mathcal{B}_i^{x,V}$ for this sort. The equivalence classes of $\mathcal{A}^{x,V}$ for this sort are exactly the structures $\mathcal{B}_j^{x,U_{i,m}}$. Note that for all j we have $(x,j) \notin U_{i,m}$ and so by (2), $\mathcal{B}_j^{x,U_{i,m}} \not\preceq_{n-1} \mathcal{A}^{x,U_{i,m}}$. Thus, we can apply Sublemma 2.1.3 to conclude that $\mathcal{B}_i^{x,V} \not\preceq_n \mathcal{A}^{x,V}$ on the sort indexed by i, m and indeed $\mathcal{B}_i^{x,V} \not\preceq_n \mathcal{A}^{x,V}$ follows at once. \square

Claims 2.1.2, and 2.1.4 finish the inductive argument. We now argue for the full statement of the lemma. Consider a Π_{2m}^0 set $V \subseteq 2^\omega$. Note that $U := V \times \{0\}$ is a Π_{2m} subset of $2^\omega \times \omega$. The above inductive construction (with parameter $n = m + 1$) gives us structures $\mathcal{M}^x = \mathcal{A}^{x,U}$ and $\mathcal{N}^x = \mathcal{B}_0^{x,U}$ such that

$$x \in V \iff (x,0) \in U \iff \mathcal{A}^{x,U} \leq_m \mathcal{B}_0^{x,U} \iff \mathcal{M}^x \leq_m \mathcal{N}^x.$$

This gives the desired effective reduction from V to the sets of pairs of structures $(\mathcal{M}, \mathcal{N})$ related by $\mathcal{M} \leq_m \mathcal{N}$ and completes the proof of the lemma. \square

We delay giving a lower bound on the definability of the back-and-forth relations in the infinite case until Section 5 where we use jump inversions. The finite cases contain the critical combinatorics.

3. UPPER BOUNDS FOR THE DEFINABILITY OF THE α -TYPES OF STRUCTURES

The remaining statements of Theorem 1.3 are concerned with the case where we fix a structure \mathcal{M} and consider whether some other structure \mathcal{N} satisfies $\mathcal{N} \geq_\alpha \mathcal{M}$ or $\mathcal{N} \leq_\alpha \mathcal{M}$. In this section, we will prove the needed upper bounds on the complexities of the sets $\{\mathcal{N} : \mathcal{M} \leq_\alpha \mathcal{N}\}$ and $\{\mathcal{N} : \mathcal{M} \geq_\alpha \mathcal{N}\}$. We begin with general bounds which hold for most α , followed by certain exceptional cases where we can prove better bounds for small α or α near limit ordinals. All of these results are optimal; we provide the corresponding proofs of the lower bounds in future sections.

3.1. General bounds. The following two lemmas provide general bounds for the α -types of structures that are superior to the $\Pi_{2\alpha}^0$ bounds provided by the previous analysis.

Lemma 3.1. *Let \mathcal{M} be a fixed countable structure and let $\bar{a} \in \mathcal{M}$. For each $\alpha < \omega_1$ there is a $\Pi_{\alpha+2}$ formula φ such that $\mathcal{N} \models \varphi(\bar{b})$ if and only if $(\mathcal{N}, \bar{b}) \geq_\alpha (\mathcal{M}, \bar{a})$. Hence the set*

$$\{\mathcal{N} : \mathcal{M} \leq_\alpha \mathcal{N}\}$$

is $\Pi_{\alpha+2}^0$.

Proof. The classical analysis of the back-and-forth relations (see, e.g., Lemma VI.14 in [Mon]) gives several base cases for this claim. The formulas given there are $\Pi_{2\alpha}$, and for $\alpha = 1$ or 2 and $\alpha = \lambda$ or $\lambda + 1$ for a limit level λ we have $2\alpha \leq \alpha + 2$ and so the claim is already shown. (In fact, for $\alpha = 1$, there is even a Π_1 formula satisfying the lemma.)

We now complete the proof by proving the following inductive claim.

Claim. *If for every $\bar{a} \in \mathcal{M}$ there is a formula $\varphi_{\mathcal{M}, \bar{a}}^\alpha(\bar{x})$ satisfying the claim for $\alpha \in \omega_1$ then for every $\bar{a} \in \mathcal{M}$ there is a formula $\varphi_{\mathcal{M}, \bar{a}}^{\alpha+2}(\bar{x})$ satisfying the claim for $\alpha + 2$.*

Proof. Suppose that for each \mathcal{M} and $\bar{q} \in \mathcal{M}$ there is a $\Pi_{\alpha+2}$ formula $\varphi_{\mathcal{M}, \bar{q}}^\alpha(\bar{x})$ such that $\mathcal{N} \models \varphi(\bar{p})$ if and only if $(\mathcal{N}, \bar{p}) \geq_\alpha (\mathcal{M}, \bar{q})$. Given \mathcal{M} and $\bar{a} \in \mathcal{M}$ we must show that there is a $\Pi_{\alpha+4}$ formula $\varphi(\bar{x})$ such that $\mathcal{N} \models \varphi(\bar{p})$ if and only if $(\mathcal{N}, \bar{p}) \geq_{\alpha+2} (\mathcal{M}, \bar{a})$. For each $\bar{b} \in \mathcal{M}$, there is a Π_α formula $\theta_{\bar{b}}(\bar{x}, \bar{y})$ defining the Π_α type of $\bar{a}\bar{b}$ in \mathcal{M} , i.e., such that $\mathcal{M} \models \theta_{\bar{b}}(\bar{a}, \bar{c})$ if and only if $(\mathcal{M}, \bar{a}\bar{c}) \geq_\alpha (\mathcal{M}, \bar{a}\bar{b})$ (see [Mon] Lemma II.62). Let

$$\psi(\bar{x}) := \bigwedge_{\ell} \forall y_1, \dots, y_\ell \bigwedge_{b_1, \dots, b_\ell \in \mathcal{M}} \left(\theta_{\bar{b}}(\bar{x}, \bar{y}) \longrightarrow \varphi_{\mathcal{M}, \bar{a}\bar{b}}^\alpha(\bar{x}, \bar{y}) \right)$$

and let

$$\chi(\bar{x}) := \bigwedge_k \forall y_1, \dots, y_k \bigvee_{b_1, \dots, b_k \in \mathcal{M}} \bigwedge_{c_1, \dots, c_\ell \in \mathcal{M}} \exists z_1, \dots, z_\ell \theta_{\bar{b}\bar{c}}(\bar{x}, \bar{y}, \bar{z}).$$

Note that ψ is $\Pi_{\alpha+2}$ and χ is $\Pi_{\alpha+4}$, so that $\psi \wedge \chi$ is $\Pi_{\alpha+4}$. We claim that $\mathcal{N} \models \psi(\bar{p}) \wedge \chi(\bar{p})$ if and only if $(\mathcal{N}, \bar{p}) \geq_{\alpha+2} (\mathcal{M}, \bar{a})$ and therefore is our desired formula.

First, suppose that $\mathcal{N} \models \psi(\bar{p}) \wedge \chi(\bar{p})$. Given $\bar{q} \in \mathcal{N}$, by $\mathcal{N} \models \chi(\bar{p})$ there is $\bar{b} \in \mathcal{M}$ such that

$$\mathcal{N} \models \bigwedge_{c_1, \dots, c_\ell \in \mathcal{M}} \exists z_1, \dots, z_\ell \theta_{\bar{b}\bar{c}}(\bar{p}, \bar{q}, \bar{z}).$$

Then for any $\bar{c} \in \mathcal{M}$, there is $\bar{r} \in \mathcal{N}$ such that

$$\mathcal{N} \models \theta_{\bar{b}\bar{c}}(\bar{p}, \bar{q}, \bar{r}).$$

By $\mathcal{N} \models \psi(\bar{p})$, we get

$$\mathcal{N} \models \varphi_{\mathcal{M}, \bar{a}\bar{b}\bar{c}}^\alpha(\bar{p}, \bar{q}, \bar{r}).$$

This implies that $(\mathcal{N}, \bar{p}\bar{q}\bar{r}) \geq_\alpha (\mathcal{M}, \bar{a}\bar{b}\bar{c})$. In particular, to win the $(\mathcal{N}, \bar{p}) \geq_{\alpha+2} (\mathcal{M}, \bar{a})$ game, the Duplicator can play \bar{b} in response to \bar{q} on the first move, \bar{r} in response to \bar{c} on the second move, and then play the winning strategy for $(\mathcal{N}, \bar{p}\bar{q}\bar{r}) \geq_\alpha (\mathcal{M}, \bar{a}\bar{b}\bar{c})$ to win the game, as desired.

On the other hand, suppose that $(\mathcal{N}, \bar{p}) \geq_{\alpha+2} (\mathcal{M}, \bar{a})$. We must argue that $\mathcal{N} \models \psi(\bar{p})$ and $\mathcal{N} \models \chi(\bar{p})$. First, we argue that $\mathcal{M} \models \psi(\bar{a})$ and $\mathcal{M} \models \chi(\bar{a})$. The former is due to the definitions of $\theta_{\bar{b}}$ and $\varphi_{\mathcal{M}, \bar{a}\bar{b}}^\alpha$. The latter is because $\mathcal{M} \models \theta_{\bar{b}\bar{c}}(\bar{a}, \bar{b}, \bar{c})$ for any \bar{b} and \bar{c} as the quantifiers and conjunctions/disjunctions are essentially both over elements of \mathcal{M} . Now since ψ is $\Pi_{\alpha+2}$, and $(\mathcal{N}, \bar{p}) \geq_{\alpha+2} (\mathcal{M}, \bar{a})$, we have $\mathcal{N} \models \psi(\bar{p})$. Though χ is $\Pi_{\alpha+4}$, we can also argue that because $(\mathcal{M}, \bar{a}) \models \chi$ and $(\mathcal{N}, \bar{p}) \geq_{\alpha+2} (\mathcal{M}, \bar{a})$ we get $\mathcal{N} \models \chi(\bar{p})$. We can argue this by hand, but one could also note that χ is $\mathfrak{A}_{\alpha+2}$ and apply Proposition 1.6 (and indeed the complexity of the argument below is one justification for considering this complexity class of formulas as we do in Section 6). To show that $\mathcal{N} \models \chi(\bar{p})$, for all k and \bar{q} we must show that

$$\mathcal{N} \models \bigvee_{b_1, \dots, b_k \in \mathcal{M}} \bigwedge_{c_1, \dots, c_\ell \in \mathcal{M}} \exists z_1, \dots, z_\ell \theta_{\bar{b}\bar{c}}(\bar{p}, \bar{q}, \bar{z}).$$

Choose $\bar{u} \in \mathcal{M}$ such that $(\mathcal{N}, \bar{p}\bar{q}) \leq_{\alpha+1} (\mathcal{M}, \bar{a}\bar{u})$. Since $\mathcal{M} \models \chi$, we have

$$\mathcal{M} \models \bigvee_{b_1, \dots, b_k \in \mathcal{M}} \bigwedge_{c_1, \dots, c_\ell \in \mathcal{M}} \exists z_1, \dots, z_\ell \theta_{\bar{b}\bar{c}}(\bar{a}, \bar{u}, \bar{z}).$$

So, for some disjoint \bar{b} (in fact for $\bar{b} = \bar{u}$) for every \bar{c} there are $\bar{v}_{\bar{c}} \in \mathcal{M}$ such that

$$\mathcal{M} \models \theta_{\bar{b}\bar{c}}(\bar{a}, \bar{u}, \bar{v}_{\bar{c}}).$$

For each \bar{c} choose $\bar{r}_{\bar{c}}$ such that $(\mathcal{M}, \bar{a}\bar{u}\bar{v}_{\bar{c}}) \leq_\alpha (\mathcal{N}, \bar{p}\bar{q}\bar{r}_{\bar{c}})$. Since $\theta_{\bar{b}\bar{c}}$ is Π_α we get

$$\mathcal{N} \models \theta_{\bar{b}\bar{c}}(\bar{p}, \bar{q}, \bar{r}_{\bar{c}}).$$

Thus there is \bar{b} such that for every \bar{c} there is $\bar{r}_{\bar{c}}$ such that

$$\mathcal{N} \models \bigvee_{b_1, \dots, b_k \in \mathcal{M}} \bigwedge_{c_1, \dots, c_\ell \in \mathcal{M}} \exists z_1, \dots, z_\ell \theta_{\bar{b}\bar{c}}(\bar{p}, \bar{q}, \bar{z}).$$

Since we have shown this for all k and \bar{q} , we have that $\mathcal{N} \models \chi(\bar{p})$. □

This completes the proof of Lemma 3.1. □

Note that for α odd, finite, and greater than 1 we do not get a better complexity even though the base case is stronger than required by the lemma. This is of no help in the inductive steps—the complexity comes from the formula χ , which does not involve the formula given by the inductive hypothesis.

We now prove the analog of Lemma 3.1 for $\{\mathcal{N} : \mathcal{N} \leq_\alpha \mathcal{M}\}$.

Lemma 3.2. *Let \mathcal{M} be a fixed countable structure. For each $\alpha \in \omega_1$ there is a $\Pi_{\alpha+3}$ formula φ such $\mathcal{N} \models \varphi(\bar{b})$ if and only if $(\mathcal{N}, \bar{b}) \leq_\alpha (\mathcal{M}, \bar{a})$. Hence the set*

$$\{\mathcal{N} : \mathcal{N} \leq_\alpha \mathcal{M}\}$$

is $\Pi_{\alpha+3}^0$.

Proof. If α is a limit ordinal, this claim follows at once from the classical description of back-and-forth relations. Therefore, we can assume that $\alpha = \beta + 1$. We note that

$$(\mathcal{N}, \bar{b}) \leq_\alpha (\mathcal{M}, \bar{a}) \iff \mathcal{N} \models \bigwedge_{\beta < \alpha} \bigwedge_{\bar{c} \in \mathcal{M}} \exists \bar{x} (\mathcal{N}, \bar{b}, \bar{x}) \geq_\beta (\mathcal{M}, \bar{a}, \bar{c}).$$

By Lemma 3.1 each formula $(\mathcal{N}, \bar{b}, \bar{d}) \geq_\beta (\mathcal{M}, \bar{a}, \bar{c})$ can be expressed in a $\Pi_{\beta+2}$ manner. This means that, overall, the formula is $\Pi_{\alpha+3}$. \square

One contrast between the formulas described in Lemmas 3.1 and 3.2 to capture the back-and-forth relations and those described in the classical analysis of Lemma VI.14 in [Mon] lies in the computability of the formulas. The method in Lemma VI.14 in [Mon] is always computable in the simplest presentation of the structure \mathcal{M} . On the other hand, Lemmas 3.1 and 3.2 rely on Π_α formulas that isolate the entire Π_α -type of a tuple *within* a given structure (provided by [Mon] Lemma II.62). Extracting such a formula given a tuple and a structure is not an effective procedure. In sum, there is a tradeoff at play here. The formulas in Lemmas 3.1 and 3.2 may be far simpler in quantifier complexity when compared to the previously known descriptions of back-and-forth relations, but they are less computable. We conjecture that this tradeoff is necessary.

Conjecture 3.3.

(1) *For $n \geq 2$ even, there is a structure \mathcal{M} such that*

$$\{\mathcal{N} : \mathcal{N} \geq_n \mathcal{M}\}$$

is Π_{2n}^0 but not Σ_{2n}^0 .

(2) *For $n \geq 3$ odd, there is a structure \mathcal{M} such that*

$$\{\mathcal{N} : \mathcal{N} \geq_n \mathcal{M}\}$$

is Π_{2n-1}^0 but not Π_{2n-1}^0 .

(3) *For $n \geq 2$ even, there is a structure \mathcal{M} such that*

$$\{\mathcal{N} : \mathcal{N} \leq_n \mathcal{M}\}$$

is Π_{2n-1}^0 but not Σ_{2n-1}^0 .

(4) *For $n \geq 3$ odd, there is a structure \mathcal{M} such that*

$$\{\mathcal{N} : \mathcal{N} \leq_n \mathcal{M}\}$$

is Π_{2n}^0 but not Σ_{2n}^0 .

Moreover, each of these is witnessed by an index set result for countable structures, e.g., for (1), the index set $\{i : \mathcal{N}_i \leq_n \mathcal{M}\}$ is Π_{2n}^0 m -complete.

Note that the difference between the odd and even cases is due to the base case, and whether the last case of the back-and-forth game is in \mathcal{M} or \mathcal{N} . See the difference between Lemma 3.5 and Lemma 3.6.

(1), for example, would be implied by the following conjecture from Scott analysis:

Conjecture 3.4. *For each even $n \geq 2$, there is a computable structure \mathcal{M} with a Π_n Scott sentence but no computable Σ_{2n} Scott sentence.*

Alvir, Knight, and McCoy [AKMC20] produce a structure with a Π_2 Scott sentence but no computable Π_2 Scott sentence. Alvir, Csima, and Harrison-Trainor [ACHT] produce a structure with a Π_2 Scott sentence but no computable Σ_4 Scott sentence. The general case remains open.

3.2. Exceptional bounds. We now consider the simplest cases of Theorem 1.3 with $\alpha = 1$, $\alpha = 2$, or α near a limit ordinal. These small values have improved descriptions for the α -type when compared to the general results of the previous section.

Lemma 3.5. *For any structure \mathcal{M} the set*

$$\{\mathcal{N} : \mathcal{N} \geq_1 \mathcal{M}\}$$

is Π_1^0 .

Proof. The conjunction of the finitary universal sentences true in \mathcal{M} defines the set as all Π_1 formulas are a conjunction of these formulas. \square

Lemma 3.6. *For any structure \mathcal{M} the set*

$$\{\mathcal{N} : \mathcal{N} \leq_1 \mathcal{M}\}$$

is Π_2^0 .

Proof. This is the same upper bound given previously, straight from the definition. Let $\theta_{\bar{a}}(\bar{x})$ be the finitary quantifier-free formula which says that \bar{x} and \bar{a} satisfy the same atomic formulas (from among the first $|\bar{a}|$ -many formulas). Then

$$\mathcal{N} \leq_1 \mathcal{M} \iff \mathcal{N} \models \bigwedge_{\bar{a} \in \mathcal{M}^n} \exists y_1, \dots, y_n \theta_{\bar{a}}(y_1, \dots, y_n). \quad \square$$

Lemma 3.7. *For any structure \mathcal{M} the set*

$$\{\mathcal{N} : \mathcal{N} \leq_2 \mathcal{M}\}$$

is Π_3^0 .

Proof. Note that

$$\mathcal{N} \leq_2 \mathcal{M} \iff \bigwedge_{\bar{a} \in \mathcal{M}} \exists \bar{x} (\mathcal{N}, \bar{x}) \geq_1 (\mathcal{M}, \bar{a})$$

which is Π_3^0 by the upper bound established in Lemma 3.5. \square

Another place where the bounds are exceptional is near limit ordinals. Just like the small finite values, we can improve the bounds for these ordinals. We consider these cases below.

Lemma 3.8. *For any structure \mathcal{M} we have the following.*

- (1) *If $\alpha = \lambda$ where λ is a limit ordinal, the set $\{\mathcal{N} : \mathcal{N} \leq_\alpha \mathcal{M}\} = \{\mathcal{N} : \mathcal{N} \geq_\alpha \mathcal{M}\}$ is Π_α^0 .*
- (2) *If $\alpha = \lambda + 1$ where λ is a limit ordinal, the set $\{\mathcal{N} : \mathcal{N} \geq_\alpha \mathcal{M}\}$ is $\Pi_{\alpha+1}^0$.*
- (3) *If $\alpha = \lambda + 1$ where λ is a limit ordinal, the set $\{\mathcal{N} : \mathcal{N} \leq_\alpha \mathcal{M}\}$ is $\Pi_{\alpha+1}^0$.*

Proof. Each of these follows immediately from the classical description of the back-and-forth relations from Lemma VI.14 in [Mon]. In particular, note that $\lambda = 2\lambda$ and $(\lambda + 1) + 1 = 2(\lambda + 1)$. \square

Lemma 3.9. *If λ is a limit ordinal the set*

$$\{\mathcal{N} : \mathcal{M} \geq_{\lambda+2} \mathcal{N}\}$$

is $\Pi_{\lambda+3}^0$ for all \mathcal{M} .

Proof. For this proof, we let $p_{\bar{a}}$ denote the Π_λ -type of $\bar{a} \in \mathcal{M}$, and note that $p_{\bar{a}}(\bar{x})$ can be expressed as a Π_λ formula (as the conjunction of the formulas $\varphi_{\bar{a}}^\alpha(\bar{x})$ for $\alpha < \lambda$ expressing that $(\mathcal{N}, \bar{x}) \geq_\alpha (\mathcal{M}, \bar{a})$). First, if $\mathcal{M} \geq_{\lambda+2} \mathcal{N}$, then the Π_λ -types realised in \mathcal{N} are the same as the Π_λ -types realised in \mathcal{M} . (Or, what is the same, $\mathcal{M} \equiv_{\lambda+1} \mathcal{N}$). This can be expressed by the $\Pi_{\lambda+2}$ sentence

$$\psi = \bigwedge_{\bar{a} \in \mathcal{M}} \exists \bar{x} p_{\bar{a}}(\bar{x}) \wedge \bigwedge_n \forall x_1, \dots, x_n \bigvee_{\bar{a} \in \mathcal{M}} p_{\bar{a}}(\bar{x}).$$

Now for each $\bar{a} \in \mathcal{M}$, let $P_{\bar{a}}$ be the set of all Π_λ -types extending that of \bar{a} realised in the structure \mathcal{M} ,

$$P_{\bar{a}} = \{p_{\bar{a}\bar{b}}(\bar{x}, \bar{y}) : \bar{b} \in \mathcal{M}\}.$$

Let $\overline{P_{\bar{a}}}$ be the set of all Π_λ types realised in \mathcal{M} but not in $P_{\bar{a}}$. Consider the sentence

$$\theta = \bigwedge_{\bar{a} \in \mathcal{M}} \exists \bar{x} \bigwedge_{p \in \overline{P_{\bar{a}}}} \neg \exists \bar{y} p(\bar{x}, \bar{y}).$$

Note that θ is true in \mathcal{M} by construction and that it is a $\Pi_{\lambda+3}$ formula. In particular, θ is a conjunction of $\Sigma_{\lambda+2}$ formulas. If $\mathcal{M} \geq_{\lambda+2} \mathcal{N}$, then since θ is true in \mathcal{M} , each of its $\Sigma_{\lambda+2}$ conjuncts is true in \mathcal{N} . This means that θ is true in \mathcal{N} . (Alternatively, one can note that θ is $\overline{\mathfrak{E}}_\alpha$ and appeal to Proposition 1.6 which we prove later.) Thus if $\mathcal{M} \geq_{\lambda+2} \mathcal{N}$ then $\mathcal{N} \models \psi \wedge \theta$.

On the other hand, suppose that $\mathcal{N} \models \psi \wedge \theta$. Then \mathcal{M} and \mathcal{N} realise the same Π_λ types. Given $\bar{a} \in \mathcal{M}$, choose $\bar{b} \in \mathcal{N}$ such that

$$\mathcal{N} \models \bigwedge_{p \in \overline{P_{\bar{a}}}} \neg \exists \bar{y} p(\bar{b}, \bar{y}).$$

Now given $\bar{b}' \in \mathcal{N}$, we have that

$$\mathcal{N} \models \bigwedge_{p \in \overline{P_{\bar{a}}}} \neg p(\bar{b}, \bar{b}').$$

But the Π_λ types realised in \mathcal{N} are the same as the Π_λ -types realised in \mathcal{M} because $\mathcal{N} \models \psi$, and so since $\bar{b}\bar{b}'$ does not realise one of the types in $\overline{P_{\bar{a}}}$ it must realise one of the types in $P_{\bar{a}}$, say $p_{\bar{a}\bar{a}'}$. Then $(\mathcal{M}, \bar{a}\bar{a}') \equiv_\lambda (\mathcal{N}, \bar{b}\bar{b}')$. Working backwards through the proof gives that $(\mathcal{M}, \bar{a}) \geq_{\lambda+1} (\mathcal{N}, \bar{b})$ and so $\mathcal{M} \geq_{\lambda+2} \mathcal{N}$. Thus if $\mathcal{N} \models \psi \wedge \theta$ then $\mathcal{M} \geq_{\lambda+2} \mathcal{N}$. \square

4. LOWER BOUNDS FOR THE DEFINABILITY OF THE α -TYPES OF STRUCTURES

We now consider lower bounds for the complexity of the sets $\{\mathcal{N} : \mathcal{M} \leq_\alpha \mathcal{N}\}$ and $\{\mathcal{N} : \mathcal{M} \geq_\alpha \mathcal{N}\}$. We only consider particular values of α and will show how to use these results to prove the complete characterization in Section 5. We begin by considering small finite values for which the back-and-forth relations are very easy to define. We then look at slightly larger finite values that exhibit more general behavior and are therefore suitable base cases for the jump inversion argument in Section 5. Lastly, we analyze exceptional cases near limit ordinals. In all cases, we have already shown that the sets are of the specified complexity, and so it is just the hardness results that remain.

4.1. Lower bounds for small finite cases. The following three cases are too small to exhibit general behavior, so they are treated separately.

Lemma 4.1. *There is a structure \mathcal{M} such that the set*

$$\{\mathcal{N} : \mathcal{N} \geq_1 \mathcal{M}\}$$

is Π_1^0 -complete.

Proof. Let \mathcal{M} be a structure in the language of one unary relation U where U holds of exactly one element. Given a $\mathbf{\Pi}_1^0$ set A , we can build for each $x \in 2^\omega$ a structure \mathcal{N}^x such that U holds of exactly one element in \mathcal{N}^x (and hence $\mathcal{N}^x \geq_1 \mathcal{M}$) if $x \in A$ and U holds of exactly two elements in \mathcal{N}^x (and hence $\mathcal{N}^x \not\leq_1 \mathcal{M}$) if $x \notin A$. \square

Lemma 4.2. *There is a structure \mathcal{M} such that the set*

$$\{\mathcal{N} : \mathcal{N} \leq_1 \mathcal{M}\}$$

is $\mathbf{\Pi}_2^0$ -complete.

Proof. Let \mathcal{M} be the structure in the language of one unary relation U where U holds of an infinite-coinfinite subset of the elements of \mathcal{M} . The set $\text{Inf} \subseteq 2^\omega$ of strings with infinitely many 1's is $\mathbf{\Pi}_2^0$ -complete (see e.g., Section 23.A of [Kec95] for this and other examples of complete sets), so we will reduce it to $\{\mathcal{N} : \mathcal{N} \leq_1 \mathcal{M}\}$. Given $x \in 2^\omega$ we construct \mathcal{N}^x so that $x \in \text{Inf} \iff \mathcal{N}^x \leq_1 \mathcal{M}$. If \mathcal{N}^x has domain \mathbb{N} , let $\mathcal{N}^x \models U(2i)$ if and only if $x(i) = 1$ (U will never hold of any odd elements). If x has infinitely many 1's then $\mathcal{N}^x \cong \mathcal{M}$, so in particular $\mathcal{N}^x \leq_1 \mathcal{M}$. If x has finitely many 1's, say n of them, then in \mathcal{M} at least $n+1$ distinct elements satisfy the relation U , but in \mathcal{N}^x there are only n . Thus $\mathcal{N}^x \not\leq_1 \mathcal{M}$ as desired. \square

Lemma 4.3. *There is a structure \mathcal{M} such that the set*

$$\{\mathcal{N} : \mathcal{N} \leq_2 \mathcal{M}\}$$

is $\mathbf{\Pi}_3^0$ -complete.

Proof. Identify $(2^\omega)^\omega$ with all lists $(a_i)_{i \in \omega}$ of infinite binary strings. Let $A \subseteq (2^\omega)^\omega$ be the set of lists $(a_i)_{i \in \omega}$ such that every a_i has finitely many 1's. This set is $\mathbf{\Pi}_3^0$ -complete.

Consider structures of the following type: they have countably many sorts, labeled by unary relations R_i , each sort consisting of a linear ordering. Given a structure \mathcal{C} of this form, we write \mathcal{C}_i to mean the linear ordering defined by the R_i points. We let \mathcal{M} be the unique structure of this form with $\mathcal{M}_i \cong \omega$ for all i . Given $x = (x_i)_{i \in \omega}$ we construct in a continuous way \mathcal{N}^x , also a structure of the above form, so that $x \in A \iff \mathcal{N}^x \leq_2 \mathcal{M}$. Given i , we explain how to construct \mathcal{N}_i^x the i th sort of \mathcal{N}^x . Start with a copy of ω . Whenever we see another 1 in the string x_i , add an element below all of the previously added elements. If x_i has finitely many 1's, then in \mathcal{N}_i^x we add only finitely many elements below the initial copy of ω , and so $\mathcal{N}_i^x \cong \omega$. If a_i has infinitely many 1's, then $\mathcal{N}_i^x = \omega^* + \omega \cong \zeta$. If $x \in A$, all of the x_i have finitely many 1's and $\mathcal{N}^x \cong \mathcal{M}$, so in particular $\mathcal{N}^x \leq_2 \mathcal{M}$. On the other hand, if $x \notin A$, some x_i has infinitely many 1's, so $\mathcal{N}_i^x \cong \zeta$. Let

$$\varphi := \exists y [R_i(y) \wedge \forall z (R_i(z) \longrightarrow z \geq y)].$$

Note that $\mathcal{M} \models \varphi$, yet $\mathcal{N}^x \models \neg\varphi$, so $\mathcal{N}^x \not\leq_2 \mathcal{M}$ as desired. \square

4.2. Critical finite cases. The next two cases we consider are critical because they exhibit all of the complexity of the behavior of the general case. In Section 5, we will use jump inversion to obtain all of the other non-exceptional cases from these two.

Lemma 4.4. *There is a structure \mathcal{M} such that the set*

$$\{\mathcal{N} : \mathcal{N} \geq_2 \mathcal{M}\}$$

is $\mathbf{\Pi}_4^0$ -complete.

Proof. The structures we work with will be *flower graphs* (sometimes also called bouquet graphs). Let \mathcal{S} be a non-empty family of subsets of ω . (From now on, all of our families will be assumed to be non-empty.) There is a corresponding flower graph $\mathcal{G}_{\mathcal{S}}$. This flower graph

will have, for each $S \in \mathcal{S}$, infinitely many connected components (all isomorphic to each other) consisting of a central vertex with, for each $n \in S$, a loop of length n .

This graph encodes what we will call *positive enumerations* of the family \mathcal{S} . A positive enumeration of \mathcal{S} should be an object from which positive information about \mathcal{S} can be obtained positively, though one has to express this formally. There are many equivalent definitions, but one is as follows. A positive enumeration of a family is a function $f: \omega \times \omega \times \omega \rightarrow \{0, 1\}$ such that, for each x, y , $f(x, y, 0) = 0$ and if $f(x, y, s) = 1$ then $f(x, y, t) = 1$ for all $t \geq s$. We set

$$\hat{f}(x, y) = \lim_{s \rightarrow \infty} f(x, y, s).$$

Let

$$X_i = \{j : \hat{f}(i, j) = 1\}.$$

Then f is a positive enumeration of the family $\{X_i : i \in \omega\}$. Note that the order in which the subsets of ω appear does not matter, and sets can be repeated. The same set can show up multiple times in the enumeration.

The idea is that $\mathcal{G}_{\mathcal{S}}$ has the property, easily verified, that (a) from any enumeration of \mathcal{S} , we can compute a copy of $\mathcal{G}_{\mathcal{S}}$, and (b) from any copy of $\mathcal{G}_{\mathcal{S}}$ we can enumerate a copy of \mathcal{S} . (In particular, there is a continuous map that takes an enumeration of \mathcal{S} to a presentation of $\mathcal{G}_{\mathcal{S}}$, and a continuous map taking a presentation of $\mathcal{G}_{\mathcal{S}}$ to an enumeration of \mathcal{S} .)

We say that a family \mathcal{S} is *closed under finite additions of elements* if whenever $S \in \mathcal{S}$ and F is finite, then $S \cup F \in \mathcal{S}$.

Claim 4.4.1. *Let \mathcal{S} and \mathcal{T} be two families of subsets of ω , and suppose that both \mathcal{S} and \mathcal{T} are closed under finite additions of elements. Then $\mathcal{G}_{\mathcal{S}} \leq_2 \mathcal{G}_{\mathcal{T}}$ if and only if for each $T \in \mathcal{T}$ there is $S \in \mathcal{S}$ such that $S \subseteq T$.*

Proof. Suppose that there is $T \in \mathcal{T}$ such that for all $S \in \mathcal{S}$ we have $S \not\subseteq T$. We describe a winning strategy for Spoiler in the back-and-forth game to witness $\mathcal{G}_{\mathcal{S}} \not\leq_2 \mathcal{G}_{\mathcal{T}}$. First, Spoiler plays the central vertex $u \in \mathcal{G}_{\mathcal{T}}$ of a connected component corresponding to T . Say that Duplicator responds with the central vertex $v \in \mathcal{G}_{\mathcal{S}}$ of a connected component corresponding to $S \in \mathcal{S}$. Since $S \not\subseteq T$, there is some $n \in S$ with $n \notin T$. Then Spoiler plays an n -cycle in $\mathcal{G}_{\mathcal{S}}$ in the connected component of v , to which Duplicator cannot respond with an n -cycle in the connected component of u . If Duplicator responds with a vertex $v \in \mathcal{G}_{\mathcal{S}}$ that is not a central vertex, then Spoiler can play a path connecting v to its central vertex v' along with an n -cycle connected to v' . Again, Duplicator cannot respond with an n -cycle in the connected component of u . Thus $\mathcal{G}_{\mathcal{S}} \not\leq_2 \mathcal{G}_{\mathcal{T}}$.

Now suppose that for each $T \in \mathcal{T}$ there is $S \in \mathcal{S}$ such that $S \subseteq T$. We will describe a winning strategy for Duplicator in the back-and-forth game to witness $\mathcal{G}_{\mathcal{S}} \leq_2 \mathcal{G}_{\mathcal{T}}$. On their first play, Spoiler plays elements of $\mathcal{G}_{\mathcal{T}}$; we may assume that they play central vertices u_1, \dots, u_k together with, for each i , the elements of finitely many loops in the connected component of u_i . For each i , let F_i be the finite set of sizes of loops that Spoiler plays connected to u_i . For each i , let $T_i \in \mathcal{T}$ be the set coded by the connected component of u_i , and choose in \mathcal{S} a set S_i with $F_i \subseteq S_i \subseteq T_i$. Duplicator will respond, in $\mathcal{G}_{\mathcal{S}}$, with vertices v_1, \dots, v_k chosen such that the connected component of v_i codes $S_i \in \mathcal{S}$; using the fact that $F_i \subseteq S_i$, Duplicator also responds to each loop played by Spoiler on u_i with a loop of the same size on v_i . Now in the second round of the game suppose that Spoiler plays, in $\mathcal{G}_{\mathcal{S}}$, finitely many further loops on the v_i , together with new central vertices v'_1, \dots, v'_ℓ and finitely many loops in their connected components. Since $S_i \subseteq T_i$, Duplicator can respond to each loop on v_i with a loop on u_i of the corresponding size. For each j let G_j be the set of sizes of loops played by Spoiler on v'_j , and choose $T_j \in \mathcal{T}$ with $G_j \subseteq T_j$. Then Duplicator can respond to each v'_j with a central vertex u'_j coding T_j , and to the loops played on v'_j with loops of the same lengths on u'_j . Thus $\mathcal{G}_{\mathcal{S}} \leq_2 \mathcal{G}_{\mathcal{T}}$. \square

Let $\mathcal{S} = \{S_i \cup F : i \in \mathbb{N}, F \in [\mathbb{N}]^{<\omega}\}$ where $S_i = \{\langle i, j \rangle\}_{j \in \omega}$.

Claim 4.4.2. *The set of enumerations f of families \mathcal{T} such that for all $T \in \mathcal{T}$ there is $S \in \mathcal{S}$ with $S \subseteq T$ is Π_4^0 -complete. Moreover, given any Π_4^0 set A , there is a Wadge reduction $x \mapsto f_x$ such that each f_x is an enumeration of a family which is closed under finite additions of elements.*

Proof. Let A be the Π_4^0 -complete set of all lists $f = (f_{u,v})_{u,v \in \omega}$ of graphs of functions $f_{u,v} : \omega \rightarrow \omega$, with

$$f \in A \iff \forall u \exists v f_{u,v} \text{ is total.}$$

Given $f = (f_{u,v})$, we produce an enumeration g of the family $\mathcal{T} = \{T_{u,F} : u \in \mathbb{N}, F \in [\mathbb{N}]^{<\omega}\}$ where

$$T_{u,F} = F \cup \{\langle v, j \rangle : v \in \mathbb{N} \text{ and } f_{u,v}(j) \downarrow\}.$$

We must show that $f \in A$ if and only if for all $T \in \mathcal{T}$ there is $S \in \mathcal{S}$ such that $S \subseteq T$.

Say that $f \in A$. In this case, we must show that there is a pair $i \in \mathbb{N}, F' \in [\mathbb{N}]^{<\omega}$ such that $T_{u,F} \supseteq S_i \cup F'$. Let $F = F'$ and take i to be the witness showing that $f_{u,i}$ is total. In particular this means that $f_{u,i}(j) \downarrow$ always holds so for all j , $\langle i, j \rangle \in T_{u,F}$. It follows at once that $T_{u,F} \supseteq S_i \cup F'$ as desired.

Conversely say that there is an $S_i \in \mathcal{S}$ such that $S_i \subseteq T_{u,\emptyset}$. This means that for all j , $f_{u,i}(j) \downarrow$, that is, $f_{u,i}$ is total. This holds for any u , so we have that $f \in A$, as desired. \square

Together, these two claims finish the proof. We take $\mathcal{M} = \mathcal{G}_{\mathcal{S}}$. We reduce to $\{\mathcal{N} : \mathcal{N} \geq_2 \mathcal{M}\}$ any Π_4^0 set B by producing, for each x , an enumeration f_x of a family \mathcal{T}_x which is closed under finite additions of elements, and from this a presentation of $\mathcal{G}_{\mathcal{T}_x}$. Then $\mathcal{G}_{\mathcal{T}_x} \geq_2 \mathcal{G}_{\mathcal{S}}$ if and only if $x \in B$. \square

Lemma 4.5. *There is a structure \mathcal{M} such that*

$$\{\mathcal{N} : \mathcal{N} \leq_3 \mathcal{M}\}$$

is Π_6^0 -complete.

Proof. We will make use of the construction of Lemma 4.4. Let $\mathcal{S} = \{S_i \cup F : i \in \mathbb{N}, F \in [\mathbb{N}]^{<\omega}\}$, where $S_i = \{\langle i, j \rangle\}_{j \in \omega}$, be the family of subsets of ω defined in Lemma 4.4 and let \mathcal{S}_{fin} be the family of all finite subsets of ω .

In addition to what we proved earlier, we will need two additional facts. First, in Claim 4.4.2 we can replace the family \mathcal{T} by $\mathcal{T} \cup \mathcal{S}$ to assume that \mathcal{T} contains \mathcal{S} . This modification does not change whether for all $T \in \mathcal{T}$ there is $S \in \mathcal{S}$ with $S \subseteq T$. Second, under this assumption, we can improve one direction of Claim 4.4.1 as follows.

Claim 4.5.1. *Let \mathcal{S} and \mathcal{T} be two families of subsets of ω , and suppose that both \mathcal{S} and \mathcal{T} are closed under finite additions of elements. Suppose that $\mathcal{S} \subseteq \mathcal{T}$ and that for each $T \in \mathcal{T}$ there is $S \in \mathcal{S}$ such that $S \subseteq T$. Then $\mathcal{G}_{\mathcal{S}} \geq_3 \mathcal{G}_{\mathcal{T}}$.*

Proof. We give a strategy for Duplicator in the back-and-forth game to witness $\mathcal{G}_{\mathcal{S}} \geq_3 \mathcal{G}_{\mathcal{T}}$. On their first play, Spoiler plays elements $\bar{p} \in \mathcal{G}_{\mathcal{S}}$ intersecting connected components coding S_1, \dots, S_k . Because $S \subseteq T$, there is a natural embedding $\iota : \mathcal{G}_{\mathcal{S}} \rightarrow \mathcal{G}_{\mathcal{T}}$. Duplicator plays $\iota(\bar{p})$ in response. We claim that $(\mathcal{G}_{\mathcal{S}}, \bar{p}) \leq_2 (\mathcal{G}_{\mathcal{T}}, \iota(\bar{p}))$, so this is a winning move. Duplicator will play according to ι on the connected components coding S_1, \dots, S_k in each structure. This is a winning strategy as ι restricts to an isomorphism on these components. The remaining connected components on each side only finitely differ from $\mathcal{G}_{\mathcal{S}}$ and $\mathcal{G}_{\mathcal{T}}$. In particular, they are isomorphic to the whole of $\mathcal{G}_{\mathcal{S}}$ and $\mathcal{G}_{\mathcal{T}}$. This means that Duplicator can play a winning strategy for $\mathcal{G}_{\mathcal{S}} \leq_2 \mathcal{G}_{\mathcal{T}}$ on these components to win the $(\mathcal{G}_{\mathcal{S}}, \bar{p}) \leq_2 (\mathcal{G}_{\mathcal{T}}, \iota(\bar{p}))$ game. Under the assumptions provided, such a winning strategy is provided in Claim 4.4.1. \square

The following general claim will allow us to conclude the desired result.

Claim 4.5.2. *Let \mathcal{M} be a structure such that for all $\mathbf{\Pi}_\beta^0$ sets C there is a continuous map $x \mapsto \mathcal{N}_x$ such that:*

- (1) *if $x \in C$ then $\mathcal{N}_x \leq_{\alpha+1} \mathcal{M}$,*
- (2) *if $x \notin C$ then $\mathcal{N}_x \not\leq_\alpha \mathcal{M}$, and*
- (3) *there is a structure \mathcal{F} such that for all x $\mathcal{F} \leq_\alpha \mathcal{N}_x$ and $\mathcal{F} \not\leq_\alpha \mathcal{M}$.*

Then there is a structure $\widetilde{\mathcal{M}}$ such that $\{\widetilde{\mathcal{N}} : \widetilde{\mathcal{N}} \leq_{\alpha+1} \widetilde{\mathcal{M}}\}$ is $\mathbf{\Pi}_{\beta+2}^0$ -hard.

Proof. Let A be a $\mathbf{\Pi}_{\beta+2}^0$ set, and write

$$x \in A \iff \forall m \exists n (x, m, n) \in B$$

where B is $\mathbf{\Pi}_\beta^0$. By hypothesis for each m, n there is a structure $\mathcal{N}_{m,n}^x$ such that

$$(x, m, n) \in B \implies \mathcal{N}_{m,n}^x \leq_{\alpha+1} \mathcal{M}$$

and

$$(x, m, n) \notin B \implies \mathcal{N}_{m,n}^x \not\leq_\alpha \mathcal{M}.$$

Fix also the structure \mathcal{F} . We must define a structure $\widetilde{\mathcal{M}}$ and build, for each x , a structure $\widetilde{\mathcal{N}}^x$ such that

$$x \in A \iff \widetilde{\mathcal{N}}^x \leq_{\alpha+1} \widetilde{\mathcal{M}}.$$

We work in the language of \mathcal{M} together with infinitely many fresh unary relations U_i and a fresh equivalence relation E . Our structures $\widetilde{\mathcal{M}}$ and $\widetilde{\mathcal{N}}^x$ will be divided by the U_i into infinitely many distinguished sorts, each of which consists of infinitely many infinite E -equivalence classes, with the equivalence classes refining the sorts. On each equivalence class, there will be a structure in the language of \mathcal{M} ; the choice of structures that appear in each equivalence class of each sort will be the only difference between the different structures. For $\widetilde{\mathcal{M}}$, in each sort U_n we have one equivalence class which is a copy of \mathcal{M} and each other equivalence class is a copy of \mathcal{F} . For $\widetilde{\mathcal{N}}^x$, in each sort U_m , we put as the $2n$ th equivalence class a copy of $\mathcal{N}_{m,n}^x$, and as the $2n+1$ th equivalence class a copy of \mathcal{F} .

We claim that $\widetilde{\mathcal{N}}^x \leq_{\alpha+1} \widetilde{\mathcal{M}}$ if and only if $x \in A$. First, suppose that $\widetilde{\mathcal{N}}^x \leq_{\alpha+1} \widetilde{\mathcal{M}}$. Given m , let $u \in \widetilde{\mathcal{M}}$ be an element of the sort U_m and within U_m of the equivalence class on which we have a copy of \mathcal{M} . There must be $v \in \widetilde{\mathcal{N}}^x$ such that $(\widetilde{\mathcal{N}}^x, v) \geq_\alpha (\widetilde{\mathcal{M}}, u)$; moreover, v must be in the m th sort. If G_u is the structure on the equivalence class of u , and G_v is the structure on the equivalence class of v , then $G_v \geq_\alpha G_u$. Then v cannot be in the $2n+1$ th equivalence class of the m th sort of \mathcal{N}_x , since $\mathcal{F} \not\leq_\alpha \mathcal{M}$. Thus v must be in the $2n$ th equivalence class for some n , and $G_v \cong \mathcal{N}_{m,n}^x \geq_\alpha \mathcal{M}$. By assumption, this implies that $(x, m, n) \in B$. So for each m there is n such that $(x, m, n) \in B$, which implies that $x \in A$.

On the other hand, suppose that $x \in A$. For each m there is n such that $(x, m, n) \in B$, and so for each m there is n such that $\mathcal{N}_{m,n}^x \leq_{\alpha+1} \mathcal{M}$. We must show that $\widetilde{\mathcal{N}}^x \leq_{\alpha+1} \widetilde{\mathcal{M}}$. Since \mathcal{M} is divided into sorts by the U_i , it suffices to show that this is true on each sort. Consider the m th sort, and fix n such that $\mathcal{N}_{m,n}^x \leq_{\alpha+1} \mathcal{M}$. For each equivalence class of $\widetilde{\mathcal{M}}$ (with induced structure G), there is an equivalence class of $\widetilde{\mathcal{N}}^x$ (with induced structure H) such that $G \geq_{\alpha+1} H$; for the first equivalence class where $G \cong \mathcal{M}$ this is the $2n$ th equivalence class with $H \cong \mathcal{N}_{m,n}^x$, and for any other equivalence class with $G \cong \mathcal{F}$ this is any of the infinitely many equivalence classes with $H \cong \mathcal{F}$. Also, for each equivalence class of $\widetilde{\mathcal{N}}^x$ (with induced graph $H \cong \mathcal{N}_{m,n}^x$ or $H \cong \mathcal{F}$) there are infinitely many equivalence classes of \mathcal{M} (with induced structure $G \cong \mathcal{F}$) such that $H \geq_\alpha G$. This is enough to check that $\widetilde{\mathcal{N}}^x \leq_{\alpha+1} \widetilde{\mathcal{M}}$ (see Sublemma 2.1.1). \square

We now complete the proof of the lemma by using Claim 4.5.2 with $\alpha = 2$. By Claim 4.5.1 along with the construction in Lemma 4.4, there is a continuous map $x \mapsto \mathcal{N}_x$ satisfying the first two conditions of Claim 4.5.2. Therefore, to finish the proof, we need only to construct an \mathcal{F} that satisfies the last condition of Claim 4.5.2 relative to this map. Let $\mathcal{F} = \mathcal{G}_{\mathcal{S}_{\text{fin}}}$. Note that $\mathcal{N}_x \cong \mathcal{G}_{\mathcal{T}}$ where \mathcal{T} is always closed under finite additions of elements. This means that for all finite sets F , there is a $T \in \mathcal{T}$ such that $F \supseteq T$. By Claim 4.4.1, this means that $\mathcal{F} \leq_2 \mathcal{N}_x$. Also, $\mathcal{M} = \mathcal{G}_{\mathcal{S}}$ and \mathcal{S} has only infinite sets; none of these sets are contained in any finite set, so by Claim 4.4.1 $\mathcal{M} \not\leq_2 \mathcal{F}$, as desired. This completes the proof of the lemma. \square

4.3. Exceptional lower bounds at limit levels. The following cases are too close to a limit ordinal to exhibit general behavior, so they are considered separately.

Lemma 4.6. *If α is a limit ordinal, $\mathcal{M} \leq_{\alpha} \mathcal{N}$ if and only if $\mathcal{M} \geq_{\alpha} \mathcal{N}$. There is a structure \mathcal{M} such that the set*

$$\{\mathcal{N} : \mathcal{N} \equiv_{\alpha} \mathcal{M}\}$$

is Π_{α}^0 -complete.

Proof. The first claim is well-known; see for example the proof of Lemma II.67 in [Mon].

To see the second claim, let $\delta_i \rightarrow \alpha$ be a fundamental sequence. By the Ash and Knight Pair of Structures theorem [AK90] the set of models isomorphic to ω^{δ_i} is $\Pi_{\delta_i}^0$ -hard. Let the language of \mathcal{M} be $\{<\}$ along with infinitely many unary relations R_i . Let \mathcal{M} contain exactly one copy of each ω^{δ_i} , where R_i holds exactly of the elements in ω^{δ_i} .

Given a Π_{α}^0 set A , we can write $A = \bigcap_i A_i$ where A_i is $\Pi_{\delta_i}^0$. For $x \in 2^{\omega}$, let $\mathcal{N}_i^x \cong \omega^{\delta_i}$ if and only if $x \in A_i$. Note that this can be done in a continuous fashion. \mathcal{N}^x contains exactly a copy of each \mathcal{N}_i^x and R_i holds exactly of the elements in \mathcal{N}_i^x . As each component of \mathcal{N}^x is constructed continuously, so is all of \mathcal{N}^x . If $x \in A$ it is immediate that $\mathcal{M} \cong \mathcal{N}^x$ and so $\mathcal{M} \equiv_{\alpha} \mathcal{N}$. If $x \notin A$, then for some i $\mathcal{N}_i^x \not\cong \omega^{\delta_i}$. In particular, \mathcal{N}^x fails to satisfy the $\Pi_{2\delta_i+1}$ Scott sentence of ω^{δ_i} (give explicitly in [Mon] Lemma II.5) relativized to the predicate R_i . Thus, $\mathcal{N}^x \not\equiv_{\alpha} \mathcal{M}$, completing the proof. \square

For the next case, we need to make use of Montalbán’s tree of structures theorem as seen in [Mon14]. We only need a specific case of the theorem (namely, the one where the “tree” has only one path that is the length of a limit ordinal), so we will only state the weak version needed.

Theorem 4.7 (Montalbán [Mon14]). *Given a limit ordinal λ and a fundamental sequence $\delta_i \rightarrow \lambda$, let $\{\mathcal{L}_i\}_{i \in \omega}$ and \mathcal{L}_{ω} be structures with $\mathcal{L}_i \equiv_{\delta_i+1} \mathcal{L}_{i+1} \equiv_{\delta_i+1} \mathcal{L}_{\omega}$ for all i . Given a Π_{λ}^0 set $A = \bigcap_i A_i$ where each A_i is $\Pi_{\delta_i}^0$ there is a continuous procedure that given x outputs \mathcal{L}_i if i is the least such that $x \notin A_i$ and \mathcal{L}_{ω} if $i \in A$.*

Note that examples of such \mathcal{L}_i with the additional property that $M_i \not\equiv_{\delta_i+2} \mathcal{L}_{i+1}$ always exist; see for example, [Mon] Example IX.24 where the examples are linear orderings.

Lemma 4.8. *If $\alpha = \lambda + 1$ where λ is a limit ordinal, there is a structure \mathcal{M} such that the set*

$$\{\mathcal{N} : \mathcal{N} \geq_{\alpha} \mathcal{M}\}$$

is $\Pi_{\alpha+1}^0$ -complete.

Proof. We first construct an example \mathcal{K} such that $\{\mathcal{N} : \mathcal{N} \geq_{\alpha} \mathcal{K}\}$ is $\Sigma_{\lambda+1}^0$ complete; we will use such a \mathcal{K} as a piece of our final construction. Let $\{\mathcal{L}_i\}_{i \in \omega}$ and \mathcal{L}_{ω} be linear orderings with $\mathcal{L}_i \equiv_{\delta_i+1} \mathcal{L}_{i+1} \equiv_{\delta_i+1} \mathcal{L}_{\omega}$ and $\mathcal{L}_i \not\equiv_{\delta_i+2} \mathcal{L}_{i+1}$ for all i . \mathcal{K} will be a structure in the signature of two equivalence relations E and F and an ordering $<$. The equivalence relation F will refine E . Given some $\sigma \in (\omega + 1)^{\omega}$, let \mathcal{K}_{σ} be the structure in the language $\{F, <\}$ with infinitely many

F equivalence classes, each of which is a linear order. These linear orders will be infinitely many copies of each $\mathcal{L}_{\sigma(j)}$ for each $j \in \omega$. Then define \mathcal{K} as follows with infinitely many E equivalence classes, each of which is a structure \mathcal{K}_σ . For each $\sigma \in (\omega + 1)^\omega$ with only finitely many finite values, \mathcal{K} will have infinitely many equivalence classes with a copy of \mathcal{K}_σ .

Let B be a $\Sigma_{\lambda+1}^0$ set. Let $x \in B$ if and only if $\exists m x \in C_m$ for a family of Π_λ^0 sets C_m . Let $A_j = C_0 \cup \dots \cup C_j$. For each j , write $A_j = \bigcap_i A_{j,i}$ where each $A_{j,i}$ is $\Pi_{\delta_i}^0$. Let F_j be the continuous map from Theorem 4.7 with respect to the set A_j and let $\mathcal{N}_j^x := F_j(x)$. Note that $\bigsqcup_{j \in \omega} \mathcal{N}_j^x$ is isomorphic to \mathcal{K}_{τ_0} for some $\tau_0 \in (\omega + 1)^\omega$. Let $\{\tau_k\}_{k \in \omega}$ enumerate the set of elements in $(\omega + 1)^\omega$ that differ from τ_0 on only finitely many inputs. Let \mathcal{N}^x contain infinitely many E equivalence classes, countably many for each τ_k where the equivalence class corresponding to each τ_k contains a copy of \mathcal{K}_{τ_k} . Note that \mathcal{N}^x is constructed by countably many disjoint continuous modifications of F_j , and so in particular the map $x \mapsto \mathcal{N}^x$ is continuous. (To modify F_k , first designate countably many equivalence classes for each $p \in (\omega + 1)^{<\omega}$; then on each equivalence class corresponding to p put a copy of $\mathcal{L}_{p(\ell)}$ for $\ell \leq |p|$ and have R_ℓ hold of those points; for each $r > |p|$ put a copy of N_r^x and have R_r hold of those points.)

We now show that $x \in B$ if and only if $\mathcal{N}^x \geq_{\lambda+1} \mathcal{K}$. If $x \in B$, eventually there is a witness m such that $x \in C_m$. For $j \geq m$, this means that $x \in A_j$. In particular, $\tau_0(j)$ for $j \geq m$ is always ω . This means that the set of elements of $(\omega + 1)^\omega$ that are finitely different from τ_0 are exactly those that have finitely many finite values. In particular, $\mathcal{N}^x \cong \mathcal{K}$, so $\mathcal{N}^x \geq_{\lambda+1} \mathcal{K}$ as required. Now say that $x \notin B$. This means that for any j , $x \notin A_j$. Thus τ_0 has only finite values. Consider the formula $\psi := \exists z \bigwedge_{i \in \omega} \varphi_{\tau_0(i)}^E(z)$, where $\varphi_{\tau_0(i)}^E(z)$ is the $\Pi_{<\lambda}$ sentence that expresses that in the E -equivalence class of z there is an F -equivalence class $\delta_{\tau_0(i)} + 2$ -equivalent to $\mathcal{L}_{\tau_0(i)}$. By construction, $\mathcal{N}^x \models \psi$; any element of an equivalence class corresponding to τ_0 is a witness. However, $\mathcal{M} \models \neg\psi$; among the \mathcal{L}_j the $\delta_{\tau_0(i)} + 2$ -type of $\mathcal{L}_{\tau_0(i)}$ isolates it up to isomorphism and no E class of \mathcal{M} has all of the $\mathcal{L}_{\tau_0(i)}$ included (in fact, each such E class only has finitely many \mathcal{L}_j). As ψ is a $\Sigma_{\lambda+1}$ formula we see that $\mathcal{N}^x \not\geq_{\lambda+1} \mathcal{K}$, as desired.

Finally, we construct the desired \mathcal{M} . Let D be a $\Pi_{\lambda+2}^0$ set. Let $x \in D$ if and only if $\forall n x \in B_n$ for a family of $\Sigma_{\lambda+1}^0$ sets B_n . For each n let $\mathcal{K}_n \cong \mathcal{K}$ and \mathcal{N}_n^x be the structures constructed above for B_n . \mathcal{M} will be a structure in the signature of $<$, the linear ordering within each \mathcal{L}_i , two equivalence relations E and F along with infinitely many unary predicates $\{S_n\}_{n \in \omega}$ denoting sorts in the structure. \mathcal{M} contains a copy of each $\mathcal{K}_n \cong \mathcal{K}$, with the predicate S_n holding exactly of the elements of \mathcal{K}_n . Construct $\tilde{\mathcal{N}}_x^x$ with a copy of each of the structures \mathcal{N}_n^x for each n ; let S_n hold exactly of the elements in \mathcal{N}_n^x . It is immediate that $\tilde{\mathcal{N}}_x^x \geq_{\lambda+1} \mathcal{M}$ exactly when $\mathcal{N}_n^x \geq_{\lambda+1} \mathcal{K}_n$ for all n . By the above analysis, this happens exactly when $\forall n x \in B_n$, that is, $x \in D$, and this procedure is continuous as desired. \square

5. JUMP INVERSION

We now move the results from Lemmas 2.1, 4.2, 4.3, 4.4, and 4.5 up the Borel hierarchy. To do this, we use the method of jump inversion following the methods and notation from Chapter X.3 of [Mon], which itself follows Goncharov, Harizanov, Knight, McCoy, R. Miller, and Solomon [GHK⁺05]. The results listed in that chapter are unsuitable for our direct use; they lack uniformity in their statement and discuss constructions of computable reductions on indices rather than continuous Wadge reductions. Nevertheless, the proof techniques presented in Chapter X.3 of [Mon] do have the desired level of uniformity, even if the results are not stated in that way, and the arguments relativize in a manner that is suitable for the construction of a Wadge reduction. We provide a discussion of these techniques and the notation associated with them.

Fix $\alpha < \omega_1$. In Chapter X.3 of [Mon], one works effectively, but as we are working non-effectively, we generally do not need to worry about issues like ordinal presentations. However, if we wanted, we could work relative to a presentation of α . Let \mathcal{L}_0 and \mathcal{L}_1 be rigid and uniformly boldface $\Delta_{\alpha+1}^0$ categorical (equivalently, $\Sigma_{\alpha+1}$ -atomic or having a $\Pi_{\alpha+2}$ Scott sentence; see [Mon] Theorems II.23 and VII.21) structures with one binary relation S which are \equiv_{α} -equivalent but with $\mathcal{L}_0 \not\leq_{\alpha+1} \mathcal{L}_1$ and $\mathcal{L}_1 \not\leq_{\alpha+1} \mathcal{L}_0$. Such structures \mathcal{L}_0 and \mathcal{L}_1 exist by Lemma X.14 of [Mon] (but such examples were also constructed in [AK90]). It follows from the Ash and Knight pair of structures theorem [AK90] that for any $\Delta_{\alpha+1}^0$ set U , there is a continuous function $x \mapsto \mathcal{C}_x$ such that

$$x \in U \implies \mathcal{C}_x \cong \mathcal{L}_0$$

and

$$x \notin U \implies \mathcal{C}_x \cong \mathcal{L}_1.$$

This exact statement is not observed in the original paper of Ash and Knight, but it is a direct corollary; the first explicit published reference known to the authors for a boldface version of the theorem is [HM14] Theorem 2.2. (We also note that the boldface result follows with a small amount of extra work from Louveau and Saint Raymond's separation theorem [LR87].)

We can use these structures \mathcal{L}_0 and \mathcal{L}_1 to define the following map of structures.

Definition 5.1. *Given a graph $\mathcal{A} = (A, E)$ construct $\Phi_{\alpha}(\mathcal{A}) = (C; A, R)$ as follows in the language with a unary relation A and a 4-ary relation R . There is one A -point in $\Phi_{\alpha}(\mathcal{A})$ for each element in \mathcal{A} , and we identify them. The non- A -points in $\Phi_{\alpha}(\mathcal{A})$ are split into disjoint infinite sets $B_{a,b}$ for each pair of $a, b \in A$. Given a fixed $a, b \in A$, $R(a, b, \cdot, \cdot)$ will be a binary relation on $B_{a,b}$. If $E(a, b)$ then $(B_{a,b}, R(a, b, \cdot, \cdot))$ will be isomorphic to \mathcal{L}_0 and if $\neg E(a, b)$ then $(B_{a,b}, R(a, b, \cdot, \cdot))$ will be isomorphic to \mathcal{L}_1 .*

Intuitively, one replaces each edge with a copy of \mathcal{L}_0 and non-edge with a copy of \mathcal{L}_1 . In Chapter X.3 of [Mon], it is proven that Φ_{α} is an infinitary Σ_1 bi-interpretation with the α -canonical structural jump of the image. We collect the relevant consequences of this claim along with other observations about this construction in the theorem below.

Theorem 5.2. *For each $\alpha \in \omega_1$, the map Φ_{α} has the following properties:*

- (1) *For every $\Pi_{\alpha+\beta}$ formula φ there is a Π_{β} formula φ^* such that $\mathcal{A} \models \varphi^* \iff \Phi_{\alpha}(\mathcal{A}) \models \varphi$.*
- (2) *For every Π_{β} formula ψ there is a $\Pi_{\alpha+\beta}$ formula ψ_* such that $\mathcal{A} \models \psi \iff \Phi_{\alpha}(\mathcal{A}) \models \psi_*$.*
- (3) *$\mathcal{A} \leq_{\beta} \mathcal{B}$ if and only if $\Phi_{\alpha}(\mathcal{A}) \leq_{\alpha+\beta} \Phi_{\alpha}(\mathcal{B})$.*

Proof. Using Karp's theorem [Kar65], (3) follows directly from (1) and (2). Essentially, (1) and (2) follow directly from the results in Section 2.2 of [MR23] along with Chapter X.3 of [Mon]. To provide more detail, it follows from the existence of the Σ_1 bi-interpretation and the pull-back theorem (Theorem XI.7 of [Mon]) that

- (1) For every Π_{β} formula φ there is a Π_{β} formula φ^{**} such that

$$\mathcal{A} \models \varphi^{**} \iff \Phi_{\alpha}(\mathcal{A})^{(\alpha)} \models \varphi.$$

- (2) For every Π_{β} formula ψ there is a Π_{β} formula ψ_{**} such that

$$\mathcal{A} \models \psi \iff \Phi_{\alpha}(\mathcal{A})^{(\alpha)} \models \psi_{**}.$$

From here, it is enough to show that

- (1) For every Π_{β} formula φ^{**} there is a $\Pi_{\alpha+\beta}$ formula φ^* such that

$$\Phi_{\alpha}(\mathcal{A}) \models \varphi^* \iff \Phi_{\alpha}(\mathcal{A})^{(\alpha)} \models \varphi^{**}.$$

(2) For every $\Pi_{\alpha+\beta}$ formula ψ_* there is a Π_β formula ψ_{**} such that

$$\Phi_\alpha(\mathcal{A}) \models \psi_{**} \iff \Phi_\alpha(\mathcal{A})^{(\alpha)} \models \psi_*$$

The first item follows immediately by the definition of the α -canonical structural jump. The second item is shown by transfinite induction, with the only interesting case being the base case. We point out the uniformity in Montalbán's proof in Chapter X.3 of [Mon]. In particular, over every possible \mathcal{A} he adds the same countably many Π_α types to the language to Morleyize $\Phi_\alpha(\mathcal{A})$ and obtain $\Phi_\alpha(\mathcal{A})^{(\alpha)}$. As a result, the translation of $\Pi_{\alpha+1}$ formulas about the α -canonical structural jump into Π_1 formulas about the base structure provided in [MR23] Proposition 11 is uniform across all input structures and directly provides our desired base case. \square

The following lemmas are the key technical inputs needed to iterate our previous results through the ordinals.

Lemma 5.3. *Let X be a Polish space. Given a $\Sigma_{\alpha+1}^0$ -measurable map $F : X \rightarrow \text{Mod}(E)$, there is a continuous map $\Phi_\alpha(F) : X \rightarrow \text{Mod}(A, R)$ such that $\Phi_\alpha(F)(x) \cong \Phi_\alpha(F(x))$.*

Proof. Consider a basic clopen set $U_{m,n,i}$ in $\text{Mod}(E)$ for $n, m \in \omega$ and $i \in \{0, 1\}$. If $i = 0$, then this is the set of all structures with $m \not\# n$, and if $i = 1$, then it is the set of all structures with $m E n$. Since F is $\Sigma_{\alpha+1}^0$ -measurable, $C_{m,n,i} = F^{-1}(U_{m,n,i})$ must be $\Delta_{\alpha+1}$. Let $G_{m,n}$ be a continuous map, coming from the pair of structures theorem, reducing $(C_{m,n,0}, C_{m,n,1})$ to $(\mathcal{L}_0, \mathcal{L}_1)$. That is, for each $x \in X$,

$$m \not\#^{F(x)} n \iff x \in C_{m,n,0} \iff G_{m,n}(x) \cong \mathcal{L}_0$$

and

$$m E^{F(x)} n \iff x \in C_{m,n,1} \iff G_{m,n}(x) \cong \mathcal{L}_1.$$

We now define the map $\Phi_\alpha(F)$. Given x , $\Phi_\alpha(F)(x)$ will be the structure in the language $\{A, R\}$ which has an infinite set A which we identify with \mathbb{N} , and for $m, n \in A$ we have disjoint infinite sets $B_{m,n}$ with $R(m, n, \cdot, \cdot)$ defined on $B_{m,n}$ so that $B_{m,n}$ is isomorphic to $G_{m,n}(x)$. This is a continuous construction since each $G_{m,n}$ is continuous, and $\Phi_\alpha(F)(x) \cong \Phi_\alpha(F(x))$ by choice of the $G_{m,n}$. \square

Lemma 5.4. *If a class of graphs \mathbb{D} is Π_n^0 hard as an invariant set in $\text{Mod}(E)$ then the class of structures*

$$\Phi_\alpha(\mathbb{D}) := \{\mathcal{M} : \mathcal{M} \cong \Phi_\alpha(\mathcal{G}) \text{ for some } \mathcal{G} \in \mathbb{D}\}$$

is $\Pi_{\alpha+n}^0$ hard as an invariant set in $\text{Mod}(A, R)$. Moreover, $\Phi_\alpha(\mathbb{D})$ is $\Pi_{\alpha+n}^0$ hard within the image of Φ_α , that is, the maps witnessing these reductions always output structures isomorphic to $\Phi_\alpha(\mathcal{G})$ for some \mathcal{G} .

Proof. Let S be a $\Pi_{\alpha+n}^0$ subset of a Polish space X . We describe a continuous reduction from S to $\Phi_\alpha(\mathbb{D})$, which will immediately yield the desired result. Let S be $\Pi_{\alpha+n}^0$ relative to the parameter Y . Apply a change of topology to X to make all $\Delta_{\alpha+1}^0(Y)$ sets clopen. In particular, S is a Π_n^0 set in this new topology. Let G be the continuous map reducing S to \mathbb{D} . Returning to the viewpoint of the original topology, $G : X \rightarrow \text{Mod}(E)$ is a $\Sigma_{\alpha+1}^0$ -measurable map. By Lemma 5.3, there is a continuous map $\Phi_\alpha(G) : X \rightarrow \text{Mod}(A, R)$ such that $\Phi_\alpha(G)(x) \cong \Phi_\alpha(G(x))$. Note that

$$x \in S \iff G(x) \in \mathbb{D} \iff \Phi_\alpha(G(x)) \in \Phi_\alpha(\mathbb{D}) \iff \Phi_\alpha(G)(x) \in \Phi_\alpha(\mathbb{D}).$$

Therefore, $\Phi_\alpha(G)$ is the required continuous map witnessing the reduction. \square

We are now ready to generalize Lemmas 4.4 and 4.5 which give the general-case lower bounds.

Lemma 5.5. *For α not a limit ordinal or successor of a limit ordinal, there is a structure \mathcal{M} such that the set*

$$\{\mathcal{N} : \mathcal{N} \geq_\alpha \mathcal{M}\}$$

is $\Pi_{\alpha+2}^0$ -complete.

Proof. Let \mathcal{M} be the graph from Lemma 4.4 such that $\{\mathcal{N} : \mathcal{N} \geq_2 \mathcal{M}\}$ is Π_4^0 -complete. If $\alpha = 2$, we are done; otherwise, if $\alpha = \beta + 2 > 2$, we argue that for the jump inverse $\Phi_\beta(\mathcal{M})$ we have that $\{\mathcal{N} : \mathcal{N} \geq_\alpha \Phi_\beta(\mathcal{M})\}$ is $\Pi_{\alpha+2}^0$ -hard.

In $\text{Mod}(A, R)$ let $\text{Im}(\Phi_\beta)$ be the (closure under isomorphisms) of the image of Φ_β . By Theorem 5.2 we have $\mathcal{N} \geq_2 \mathcal{M}$ if and only if $\Phi_\beta(\mathcal{N}) \geq_\alpha \Phi_\beta(\mathcal{M})$. Thus

$$\{\mathcal{N} : \mathcal{N} \geq_\alpha \Phi_\beta(\mathcal{M})\} \cap \text{Im}(\Phi_\beta) = \{\mathcal{N}^* : \mathcal{N}^* \cong \Phi_\beta(\mathcal{N}) \text{ for some } \mathcal{N} \geq_2 \mathcal{M}\}.$$

By Lemma 5.4 this latter set is $\Pi_{\beta+4}^0$ hard, and moreover it is $\Pi_{\beta+4}^0$ hard within $\text{Im}(\Phi_\beta)$. Thus

$$\{\mathcal{N} : \mathcal{N} \geq_\alpha \Phi_\beta(\mathcal{M})\}$$

is $\Pi_{\beta+4}^0$ -hard or, equivalently, $\Pi_{\alpha+2}^0$ -hard. \square

Lemma 5.6. *For $\alpha = \beta + 3$, there is a structure \mathcal{M} such that the set*

$$\{\mathcal{N} : \mathcal{N} \leq_\alpha \mathcal{M}\}$$

is $\Pi_{\alpha+3}^0$ -complete.

Proof. Let \mathcal{M} be the structure from Lemma 4.5 such that $\{\mathcal{N} : \mathcal{N} \leq_3 \mathcal{M}\}$ is Π_6^0 -complete. Abusing notation, we can assume that \mathcal{M} is a graph up to effective bi-interpretation. If $\alpha = 3$, we are done; otherwise, if $\alpha = \beta + 3 > 3$, we argue that for the jump inverse $\Phi_\beta(\mathcal{M})$ we have that $\{\mathcal{N} : \mathcal{N} \leq_\alpha \Phi_\beta(\mathcal{M})\}$ is $\Pi_{\alpha+3}^0$ -hard. The argument is the same as in Lemma 5.5. \square

We also generalize the exceptional cases Lemmas 4.2 and 4.3 to obtain lower bounds for the remaining exceptional cases close to limit ordinals.

Lemma 5.7. *If λ is a limit ordinal there is a structure \mathcal{M} such that the set*

$$\{\mathcal{N} : \mathcal{M} \geq_{\lambda+2} \mathcal{N}\}$$

is $\Pi_{\lambda+3}^0$ -complete.

Proof. Let \mathcal{M} be structure from Lemma 4.3 such that $\{\mathcal{N} : \mathcal{N} \geq_2 \mathcal{M}\}$ is Π_3^0 -complete. Abusing notation, we can assume that \mathcal{M} is a graph up to effective bi-interpretation. If $\alpha = \lambda + 2$ for our limit ordinal λ , we argue that for the jump inverse $\Phi_\lambda(\mathcal{M})$ we have that $\{\mathcal{N} : \mathcal{N} \geq_\alpha \Phi_\lambda(\mathcal{M})\}$ is $\Pi_{\alpha+1}^0$ -hard. This is argued exactly the same as in Lemma 5.5. \square

Lemma 5.8. *If $\alpha = \lambda + 1$ where λ is a limit ordinal, there is a structure \mathcal{M} such that the set*

$$\{\mathcal{N} : \mathcal{N} \leq_\alpha \mathcal{M}\}$$

is $\Pi_{\alpha+1}^0$ -complete.

Proof. Let \mathcal{M} be the structure from Lemma 4.2 such that $\{\mathcal{N} : \mathcal{N} \leq_1 \mathcal{M}\}$ is Π_2^0 -complete. Abusing notation, we can assume that \mathcal{M} is a graph up to effective bi-interpretation. If $\alpha = \lambda + 1$ for limit ordinal λ , we argue that for the jump inverse $\Phi_\lambda(\mathcal{M})$ we have that $\{\mathcal{N} : \mathcal{N} \leq_\alpha \Phi_\lambda(\mathcal{M})\}$ is $\Pi_{\alpha+1}^0$ -hard. This is argued exactly the same as in Lemma 5.5. \square

To generalize Lemma 2.1, we need slightly more than Lemma 5.4, but the idea is essentially the same. The only difference is that we are now considering pairs of structures.

Lemma 5.9. *For α not a limit ordinal, the set of pairs of structures*

$$\{(\mathcal{M}, \mathcal{N}) : \mathcal{M} \leq_\alpha \mathcal{N}\}$$

is $\Pi_{2\alpha}^0$ -complete.

Proof. For finite ordinals we have Lemma 2.1. So assume that α is infinite and write $\alpha = \lambda + n$ for a limit ordinal λ and $n \geq 1$. Note that $2\alpha = \lambda + 2n$. Consider the construction from Lemma 2.1 showing that the set $\{(\mathcal{M}, \mathcal{N}) : \mathcal{M} \leq_n \mathcal{N}\}$ is Π_{2n}^0 -complete. Abusing notation, we can assume that \mathcal{M} and \mathcal{N} are graphs up to effective bi-interpretation. We argue using the λ -jump inversion that $\{(\mathcal{M}, \mathcal{N}) : \mathcal{M} \leq_\alpha \mathcal{N}\}$ is $\Pi_{\lambda+2n}^0$ -hard.

Consider a $\Pi_{\lambda+2n}^0$ set A . In a manner analogous to the argument in Lemma 5.4, we may refine the topology so that A is Π_{2n}^0 and find a map G that witnesses the hardness of $\{(\mathcal{M}, \mathcal{N}) : \mathcal{M} \leq_n \mathcal{N}\}$ with respect to this map. With respect to the original topology, this map G is $\Sigma_{\lambda+1}^0$ -measurable. Note that $G(x)$ is a pair $(\mathcal{M}_x, \mathcal{N}_x)$. The map G factors into two $\Sigma_{\lambda+1}^0$ -measurable maps $G_1(x) = \mathcal{M}_x$ and $G_2(x) = \mathcal{N}_x$. Lemma 5.3 gives us continuous maps $\Phi_\lambda(G_i)$ such that $\Phi_\lambda(G_i)(x) = \Phi_\lambda(G_i(x))$. Note that by Theorem 5.2,

$$x \in A \iff G_1(x) \leq_n G_2(x) \iff \Phi_\lambda(G_1(x)) \leq_{\lambda+n} \Phi_\lambda(G_2(x)) \iff \Phi_\lambda(G_1)(x) \leq_{\lambda+n} \Phi_\lambda(G_2)(x).$$

Therefore, the map $(\Phi_\lambda(G_1), \Phi_\lambda(G_2))$ is a reduction from A to $\{(\mathcal{M}, \mathcal{N}) : \mathcal{M} \leq_\alpha \mathcal{N}\}$, so the set is $\Pi_{\lambda+2n}^0$ hard. \square

6. \mathfrak{E}_α AND \mathfrak{A}_α FORMULAS

As we have seen, the Σ_α/Π_α hierarchy is not optimal for defining back-and-forth types. In particular, for a fixed \mathcal{M} , defining the set

$$\{\mathcal{N} : \mathcal{N} \geq_\alpha \mathcal{M}\}$$

requires a $\Pi_{\alpha+2}$ formula, but it is not true that if φ is $\Pi_{\alpha+2}$ and $\mathcal{M} \leq_\alpha \mathcal{N}$ and $\mathcal{M} \models \varphi$ then $\mathcal{N} \models \varphi$. In this section, we define a hierarchy of complexity classes of $\mathcal{L}_{\omega_1\omega}$ formulas which more precisely capture the back-and-forth relations. We call these the *back-and-forth complexity classes* of $\mathcal{L}_{\omega_1\omega}$ formulas.

Definition 6.1. *We define \mathfrak{A}_α and \mathfrak{E}_α (and simultaneously $\overline{\mathfrak{A}}_\alpha$ and $\overline{\mathfrak{E}}_\alpha$) by mutual transfinite recursion.*

- $\mathfrak{A}_1 := \Pi_1$
- $\mathfrak{E}_1 := \Sigma_1$
- $\mathfrak{A}_\alpha :=$ closure of $\bigcup_{\beta < \alpha} \overline{\mathfrak{E}}_\beta$ under \forall and $\mathbb{M}_{i \in \omega}$
- $\mathfrak{E}_\alpha :=$ closure of $\bigcup_{\beta < \alpha} \overline{\mathfrak{A}}_\beta$ under \exists and $\mathbb{W}_{j \in \omega}$
- $\overline{\mathfrak{E}}_\alpha :=$ closure of \mathfrak{E}_α under $\mathbb{W}_{j \in \omega}, \mathbb{M}_{i \in \omega}$
- $\overline{\mathfrak{A}}_\alpha :=$ closure of \mathfrak{A}_α under $\mathbb{W}_{j \in \omega}, \mathbb{M}_{i \in \omega}$

Each \mathfrak{A}_α formula can be written in the form

$$\mathbb{M}_{i \in I} \forall \bar{y}_i \varphi_i(\bar{x}, \bar{y}_i)$$

where the formulas φ_i are $\overline{\mathfrak{E}}_\beta$ for some $\beta < \alpha$, and similarly a \mathfrak{E}_α formula can be written in the form

$$\mathbb{W}_{i \in I} \exists \bar{y}_i \varphi_i(\bar{x}, \bar{y}_i)$$

where each φ_i is $\overline{\mathfrak{A}}_\beta$ for some $\beta < \alpha$.




FIGURE 6.1. Hierarchies of Σ_α/Π_α and $\mathcal{E}_\alpha/\bar{\mathcal{A}}_\alpha$ formulas. Solid arrows denote inclusions. The dotted arrows show how syntactic operations affect the complexity of formulas.

One could also make these definitions for $\mathcal{L}_{\infty, \omega}$ formulas by allowing the conjunctions and disjunctions to be over arbitrary sets; 1.6 still holds in this context. For simplicity we limit ourselves to $\mathcal{L}_{\omega_1 \omega}$.

The following are some basic properties of these complexity classes that can all be confirmed by straightforward induction arguments. The complexity classes \exists_α and \forall_α are defined by counting alternations of quantifiers but not counting infinite conjunctions and disjunctions.

Proposition 6.2. *Up to equivalence in countable structures:*

- $\Sigma_\alpha \subseteq \mathcal{E}_\alpha \subseteq \bar{\mathcal{E}}_\alpha \subseteq \exists_\alpha \cap \bar{\mathcal{A}}_{\alpha+1} \cap \mathcal{E}_{\alpha+2}$
- $\forall \mathcal{E}_\alpha \subseteq \mathcal{E}_\alpha, \forall \bar{\mathcal{E}}_\alpha \subseteq \bar{\mathcal{E}}_\alpha, \exists \mathcal{E}_\alpha \subseteq \mathcal{E}_{\alpha+1}, \forall \mathcal{E}_\alpha \subseteq \bar{\mathcal{A}}_{\alpha+1}$
- $\mathcal{E}_\alpha \subseteq \mathcal{E}_{\alpha+1}, \bar{\mathcal{A}}_\alpha \subseteq \bar{\mathcal{A}}_{\alpha+1}, \bar{\mathcal{E}}_\alpha \subseteq \bar{\mathcal{E}}_{\alpha+1}, \bar{\mathcal{A}}_\alpha \subseteq \bar{\mathcal{A}}_{\alpha+1}$
- $\forall \bar{\mathcal{E}}_\alpha \subseteq \bar{\mathcal{E}}_\alpha, \forall \bar{\mathcal{A}}_\alpha \subseteq \bar{\mathcal{A}}_\alpha, \exists \bar{\mathcal{E}}_\alpha \subseteq \mathcal{E}_{\alpha+2}, \forall \bar{\mathcal{E}}_\alpha \subseteq \bar{\mathcal{A}}_{\alpha+1}$
- $\neg \mathcal{E}_\alpha = \bar{\mathcal{A}}_\alpha, \neg \bar{\mathcal{A}}_\alpha = \mathcal{E}_\alpha, \neg \bar{\mathcal{E}}_\alpha = \bar{\mathcal{A}}_\alpha, \neg \bar{\mathcal{A}}_\alpha = \bar{\mathcal{E}}_\alpha.$
- for limit α : $\bar{\mathcal{E}}_\alpha = \bar{\mathcal{A}}_\alpha = \text{closure of } \bigcup_{\beta < \alpha} (\bar{\mathcal{E}}_\beta \cup \bar{\mathcal{A}}_\beta) \text{ under } \forall, \exists, \text{ thus } \bar{\mathcal{E}}_\alpha \subseteq \mathcal{E}_{\alpha+1} \text{ (not } \mathcal{E}_{\alpha+2})$
- $\bar{\mathcal{E}}_\alpha \subseteq \text{arbitrary } \forall \exists \text{ of } \Sigma_\alpha$

See Figure 6.1 for a diagram summarizing the inclusions of these syntactic classes and a diagram from the classical Σ_α hierarchy for comparison.

As indicated above, the key properties of this hierarchy lie in the way that it interacts with the α back-and-forth relations. We show first that the α back-and-forth relations guarantee agreement on $\bar{\mathcal{E}}_\alpha$ and $\bar{\mathcal{A}}_\alpha$ formulas the same way that they guarantee agreement on Σ_α and Π_α formulas even though there are more $\bar{\mathcal{E}}_\alpha$ and $\bar{\mathcal{A}}_\alpha$ formulas. In particular, we aim to extend the following theorem of Karp [Kar65] (see e.g. [Mon], Theorem II.36).

Theorem 6.3. *For any non-zero ordinal α , structures \mathcal{M} and \mathcal{N} and tuples $\bar{a} \in \mathcal{M}$ and $\bar{b} \in \mathcal{N}$, the following are equivalent:*

- (1) $(\mathcal{M}, \bar{a}) \leq_\alpha (\mathcal{N}, \bar{b})$.
- (2) Every Π_α formula true about \bar{a} in \mathcal{M} is true about \bar{b} in \mathcal{N} .
- (3) Every Σ_α formula true about \bar{b} in \mathcal{N} is true about \bar{a} in \mathcal{M} .

Given this tight connection between the complexity classes of the hyperarithmetic hierarchy and the back-and-forth game, one may wonder why these classes should not themselves be considered the classes of “back-and-forth complexity”. As it turns out, the analog of Karp’s theorem holds with Π_α replaced by $\overline{\mathfrak{A}}_\alpha$ and Σ_α replaced by $\overline{\mathfrak{E}}_\alpha$. Furthermore, the back-and-forth complexity classes will enjoy many connections with the back-and-forth relations that ordinary Π_α and Σ_α formulas do not enjoy. We begin by demonstrating the analog of Karp’s theorem for $\overline{\mathfrak{A}}_\alpha$ and $\overline{\mathfrak{E}}_\alpha$. Because these classes are bigger than Π_α and Σ_α , it is enough to demonstrate that the back-and-forth relations also preserve all $\overline{\mathfrak{A}}_\alpha$ and $\overline{\mathfrak{E}}_\alpha$ in the appropriate manner.

Proposition 1.6. *Suppose that $(\mathcal{M}, \bar{a}) \leq_\alpha (\mathcal{N}, \bar{b})$ for $\alpha \geq 1$. Then given a $\overline{\mathfrak{E}}_\alpha$ formula $\varphi(\bar{x})$ and a $\overline{\mathfrak{A}}_\alpha$ formula $\psi(\bar{x})$,*

$$\mathcal{N} \models \varphi(\bar{b}) \implies \mathcal{M} \models \varphi(\bar{a})$$

and

$$\mathcal{M} \models \psi(\bar{a}) \implies \mathcal{N} \models \psi(\bar{b})$$

Proof. We argue by induction on complexity. Our base case is $\alpha = 1$ and \mathfrak{A}_1 and \mathfrak{E}_1 formulas, which are just Π_1 and Σ_1 formulas. For these, the proposition follows directly from Theorem 6.3. Now, we have the inductive cases. In the inductive argument, we use the fact that we can group the outside quantifiers in a \mathfrak{A}_α or \mathfrak{E}_α sentence.

Suppose that $\varphi(\bar{x})$ is \mathfrak{E}_α , say $\varphi(\bar{x}) = \mathbb{W}_i \exists \bar{y}_i \theta_i(\bar{x}, \bar{y}_i)$ where the $\theta_i(\bar{x}, \bar{y}_i)$ are $\overline{\mathfrak{A}}_\beta$ for some $\beta < \alpha$. Suppose that $\mathcal{N} \models \varphi(\bar{b})$; then there is i and \bar{b}' such that $\mathcal{N} \models \theta_i(\bar{b}, \bar{b}')$. Then there is \bar{a}' such that $(\mathcal{N}, \bar{b}\bar{b}') \leq_\beta (\mathcal{M}, \bar{a}\bar{a}')$. By the induction hypothesis, $\mathcal{M} \models \theta_i(\bar{a}, \bar{a}')$. Thus $\mathcal{M} \models \varphi(\bar{a})$ as desired.

Suppose that $\varphi(\bar{x})$ is \mathfrak{A}_α , say $\varphi(\bar{x}) = \mathbb{A}_i \forall \bar{y}_i \theta_i(\bar{x}, \bar{y}_i)$ where the $\theta_i(\bar{x}, \bar{y}_i)$ are $\overline{\mathfrak{E}}_\beta$ for some $\beta < \alpha$. This case is dual to the previous one. Suppose that $\mathcal{M} \models \varphi(\bar{a})$; then for every $\bar{a}' \in \mathcal{M}$ and i , $\mathcal{M} \models \theta_i(\bar{a}, \bar{a}')$. Suppose there is some $\bar{b}' \in \mathcal{N}$ and i such that $\mathcal{N} \not\models \theta_i(\bar{b}, \bar{b}')$. Then there is \bar{a}' such that $(\mathcal{N}, \bar{b}\bar{b}') \leq_\beta (\mathcal{M}, \bar{a}\bar{a}')$. By the induction hypothesis, $\mathcal{M} \not\models \theta_i(\bar{a}, \bar{a}')$, a contradiction. Thus $\mathcal{N} \models \varphi(\bar{b})$ as desired.

For the case of $\overline{\mathfrak{E}}_\alpha$, we induct on the number of infinitary conjunctions or disjunctions in front of the base \mathfrak{E}_α formula. Suppose that $\varphi(\bar{x})$ is $\overline{\mathfrak{E}}_\alpha$, and say $\varphi(\bar{x}) = \mathbb{W}_i \theta_i(\bar{x})$ where each $\theta_i(\bar{x})$ is $\overline{\mathfrak{E}}_\alpha$ with fewer infinitary conjunctions or disjunctions in the front. Suppose that $\mathcal{N} \models \varphi(\bar{b})$; then for some i , $\mathcal{N} \models \theta_i(\bar{b})$, and so by the induction hypothesis, $\mathcal{M} \models \theta_i(\bar{a})$. Thus $\mathcal{M} \models \varphi(\bar{a})$. Similarly, suppose that $\varphi(\bar{x})$ is $\overline{\mathfrak{A}}_\alpha$, and say $\varphi(\bar{x}) = \mathbb{A}_i \theta_i(\bar{x})$ where each $\theta_i(\bar{x})$ is $\overline{\mathfrak{E}}_\alpha$. Suppose that $\mathcal{N} \models \varphi(\bar{b})$; then for all i , $\mathcal{N} \models \theta_i(\bar{b})$, and so by the induction hypothesis, $\mathcal{M} \models \theta_i(\bar{a})$ for all i . Thus $\mathcal{M} \models \varphi(\bar{a})$.

The cases where $\varphi(\bar{x})$ is $\overline{\mathfrak{A}}_\alpha$, and either of the form $\varphi(\bar{x}) = \mathbb{W}_i \theta_i(\bar{x})$ or $\varphi(\bar{x}) = \mathbb{A}_i \theta_i(\bar{x})$ where each $\theta_i(\bar{x})$ is $\overline{\mathfrak{A}}_\alpha$, is identical to the previous case. \square

The following is immediate from the above lemma, along with Karp’s theorem.

Corollary 6.4. *For any $\alpha \geq 1$, structures \mathcal{M} and \mathcal{N} , and tuples $\bar{a} \in \mathcal{M}$ and $\bar{b} \in \mathcal{N}$, the following are equivalent:*

- (1) $(\mathcal{M}, \bar{a}) \leq_\alpha (\mathcal{N}, \bar{b})$.
- (2) Every Π_α formula true about \bar{a} in \mathcal{M} is true about \bar{b} in \mathcal{N} .
- (3) Every Σ_α formula true about \bar{b} in \mathcal{N} is true about \bar{a} in \mathcal{M} .
- (4) Every \mathfrak{A}_α formula true about \bar{a} in \mathcal{M} is true about \bar{b} in \mathcal{N} .
- (5) Every \mathfrak{E}_α formula true about \bar{b} in \mathcal{N} is true about \bar{a} in \mathcal{M} .
- (6) Every $\overline{\mathfrak{A}}_\alpha$ formula true about \bar{a} in \mathcal{M} is true about \bar{b} in \mathcal{N} .
- (7) Every $\overline{\mathfrak{E}}_\alpha$ formula true about \bar{b} in \mathcal{N} is true about \bar{a} in \mathcal{M} .

This means that when it comes to interactions with back-and-forth relations, our new notions of complexity act at least as nicely as the classical notions. That said, our new notions have properties that the classical ones do not. The key difference is distilled in the following result. Classically, there may be no maximal Π_α or Σ_α formula, i.e., one which captures the entire Π_α or Σ_α theory of a given structure or tuple (we give examples of such structures in the following section). In the past, a $\Pi_{2\alpha}$ formula was used to describe theories at this level (see [Mon] Lemma VI.14), though we showed in Lemma 3.1 that $\Pi_{\alpha+2}$ is sufficient. Our larger classes contrast with this; they are always able to capture an entire theory at level α with a formula at that same level.

Proposition 1.5. *For \mathcal{M} a countable structure, $\bar{a} \in \mathcal{M}$, and $\alpha \geq 1$, there are a $\bar{\mathfrak{E}}_\alpha$ formula $\varphi_{\bar{a}, \mathcal{M}, \alpha}(\bar{x})$ and \mathfrak{A}_α formula $\psi_{\bar{a}, \mathcal{M}, \alpha}(\bar{x})$ such that for \mathcal{N} any structure,*

$$\mathcal{N} \models \varphi_{\bar{a}, \mathcal{M}, \alpha}(\bar{b}) \iff (\mathcal{M}, \bar{a}) \geq_\alpha (\mathcal{N}, \bar{b})$$

and

$$\mathcal{N} \models \psi_{\bar{a}, \mathcal{M}, \alpha}(\bar{b}) \iff (\mathcal{M}, \bar{a}) \leq_\alpha (\mathcal{N}, \bar{b}).$$

Proof. We argue by induction, starting with $\alpha = 1$. For $\psi_{\bar{a}, \mathcal{M}, 1}$, for each m there are only finitely many possible atomic m -types (using only the first m atomic formulas). Let $\theta_{\bar{a}}(\bar{x})$ be the finitary quantifier-free formula which says that \bar{x} and \bar{a} satisfy the same atomic formulas (from among the first $|\bar{a}|$ -many formulas). In a finite relational language, like that of linear orders, we can take $\theta_{\bar{a}}(\bar{x})$ to say that \bar{a} and \bar{x} have the same atomic type, i.e., are ordered in the same way. Then $(\mathcal{N}, \bar{b}) \geq_1 (\mathcal{M}, \bar{a})$ if and only if

$$\mathcal{N} \models \bigwedge_{n \in \mathbb{N}} \forall \bar{y}^n \bigvee_{\bar{a}' \in \mathcal{M}^n} \theta_{\bar{a}\bar{a}'}(\bar{b}, \bar{y}).$$

The inner disjunction looks like it is infinite, but in fact, it is not; this is because there are only finitely many possible formulas $\theta_{\bar{a}\bar{a}'}(\bar{x})$. Thus, this formula is \mathfrak{A}_1 (and in fact Π_1). For $\varphi_{\bar{a}, \mathcal{M}, 1}$, we have $(\mathcal{N}, \bar{b}) \leq_1 (\mathcal{M}, \bar{a})$ if and only if

$$\mathcal{N} \models \bigwedge_{\bar{a}' \in \mathcal{M}} \exists \bar{y} \theta_{\bar{a}\bar{a}'}(\bar{b}, \bar{y}).$$

This is $\bar{\mathfrak{E}}_1$.

Now, given $\alpha > 1$, suppose that we have $\bar{\mathfrak{E}}_\beta$ formulas $\varphi_{\bar{a}, \mathcal{M}, \beta}$ and \mathfrak{A}_β $\psi_{\bar{a}, \mathcal{M}, \beta}$ for $\beta < \alpha$. Then $(\mathcal{N}, \bar{b}) \leq_\alpha (\mathcal{M}, \bar{a})$ if and only if

$$\bigwedge_{\beta < \alpha} \bigwedge_{\bar{a}' \in \mathcal{M}} \exists \bar{y} \psi_{\bar{a}\bar{a}', \mathcal{M}, \beta}(\bar{b}, \bar{y})$$

and $(\mathcal{N}, \bar{b}) \geq_\alpha (\mathcal{M}, \bar{a})$ if and only if

$$\bigwedge_{\beta < \alpha} \bigwedge_{m \in \mathbb{N}} \forall \bar{y}^m \bigvee_{\bar{a}' \in \mathcal{M}} \varphi_{\bar{a}\bar{a}', \mathcal{M}, \beta}(\bar{b}, \bar{y}).$$

These define our desired $\bar{\mathfrak{E}}_\alpha$ formulas $\varphi_{\bar{a}, \mathcal{M}, \alpha}$ and \mathfrak{A}_α formulas $\psi_{\bar{a}, \mathcal{M}, \alpha}$. □

7. HENKIN CONSTRUCTIONS

In countable model theory, Henkin constructions with restricted sets of formulas—often in some countable fragment, or in a complexity class such as Σ_α —have proved useful, especially for type omitting arguments. We will describe two applications of the $\mathfrak{A}_\alpha/\bar{\mathfrak{E}}_\alpha$ formulas in Henkin constructions.

7.1. Scott ranks of models of a theory of linear orders. In [GHT], Gonzalez and Harrison-Trainor prove the following theorem:

Theorem 7.1 (Gonzalez and Harrison-Trainor [GHT]). *Given a satisfiable Π_α sentence T of linear orders, there is a linear order $\mathcal{B} \models T$ with a $\Pi_{\alpha+4}$ Scott sentence.*

The proof is a Henkin construction using formulas of bounded complexity. One of the two key facts about linear orders used in this proof is the well-known characterization of the back-and-forth relations between tuples in linear orders which appears, for example, as Lemma 15.8 of [AK00].

Lemma 7.2. *Given linear orders (\mathcal{A}, \bar{a}) and (\mathcal{B}, \bar{b}) (with the tuples in increasing order), $(\mathcal{A}, \bar{a}) \geq_\alpha (\mathcal{B}, \bar{b})$ if and only if for each $i = 0, \dots, n$ we have $(a_i, a_{i+1})^{\mathcal{A}} \geq_\alpha (b_i, b_{i+1})^{\mathcal{B}}$.*

In other words, the α -back-and-forth type of a tuple “factors” into the back-and-forth types of the intervals defined by the tuples. That said, this correspondence cannot necessarily be performed on a formula-by-formula basis. In other words, given a Σ_α formula that holds of \bar{c} , it is not necessarily implied by a collection of Σ_α formulas about the intervals it defines. It turns out, as proved in [GHT], that the $\mathfrak{E}_\alpha/\mathfrak{A}_\alpha$ complexity classes are exactly the right complexity classes to make this true. This lemma is provably false when \mathfrak{E}_α is replaced by Σ_α .

Lemma 7.3. *Let \mathcal{A} be a countable linear order and $a_1 < \dots < a_n$ elements of \mathcal{A} . Suppose that $\mathcal{A} \models \varphi(a_1, \dots, a_n)$ with φ a \mathfrak{E}_α formula in the language of linear orders. Then there are \mathfrak{E}_α sentences $\theta_0, \dots, \theta_n$ such that (a) for every $k = 0, \dots, n$ we have $(a_k, a_{k+1}) \models \theta_k$, and (b) if \mathcal{B} is any linear order and $b_1 < \dots < b_n$, if for every $k = 0, \dots, n$ we have $(b_k, b_{k+1}) \models \theta_k$ then $\mathcal{B} \models \varphi(b_1, \dots, b_n)$.*

The Henkin construction for proving Theorem 7.1 is a Henkin construction where all of the formulas are $\mathfrak{E}_{\alpha+1}$.

7.2. Omitting types. In this section, we will demonstrate how the $\mathfrak{E}_\alpha/\mathfrak{A}_\alpha$ formulas can be used in a Henkin construction by proving Theorem 1.7. This theorem was already known to Gonzalez (unpublished), but the proof given here is simpler than the original proof.

Scott’s isomorphism theorem states that infinitary sentences can always be used as isomorphism invariants for countable structures [Sco65]. Based on this, Montalbán defined his notion of robust Scott rank in a way that robustly aligns with the hyperarithmetic hierarchy [Mon15]. In this section, we show that his notion of Scott rank also aligns well with our new notion of syntactic complexity. This gives a new set of criteria equivalent to Montalbán’s notion of Scott rank, showing that it is even more robust than we previously thought. In particular, the addition of more infinitary conjunctions and disjunctions to our formulas does not increase their expressive power too much when it comes to defining an isomorphism invariant. To demonstrate these claims, we perform a type-omitting argument that extends and supersedes that of Montalbán from [Mon15].

We begin by defining a new syntactic class closely related to the \mathfrak{E}_α hierarchy.

Definition 7.4. *A formula is $\forall\mathfrak{E}_\alpha$ if it is of the form $\mathbb{M}_{i \in \omega} \forall \bar{x}_i \varphi_i(x_i)$ with $\varphi_i \in \mathfrak{E}_\alpha$.*

To be explicit, the difference between $\forall\mathfrak{E}_\alpha$ and $\mathfrak{A}_{\alpha+1}$ is that $\mathfrak{A}_{\alpha+1}$ adds universal quantifiers and conjunctions to \mathfrak{E}_α formulas while $\forall\mathfrak{E}_\alpha$ only adds universal quantifiers and conjunctions to the slightly more restricted class of \mathfrak{E}_α formulas. We begin with a helpful lemma and then prove a type-omitting theorem for $\forall\mathfrak{E}_\alpha$ formulas.

Lemma 7.5. *Given a countable structure \mathcal{M} and any $\varphi(\bar{x}) \in \mathfrak{E}_\alpha$ there is a Σ_α formula $\theta(\bar{x})$ such that*

$$\mathcal{M} \models \forall \bar{x} (\theta(\bar{x}) \iff \varphi(\bar{x})).$$

Proof. Write the $\varphi(\bar{x})$ as $\bigvee_{i \in \omega} \exists \bar{x}_i \psi_i(\bar{x}_i, \bar{y})$ where each ψ_i is $\bar{\mathfrak{A}}_{\beta_i}$ for $\beta_i < \alpha$. Consider S , the set of witnesses to $\varphi(\bar{x})$ in \mathcal{M} (note that if S is empty, the statement of the lemma is trivial). For each $\bar{a} \in \mathcal{M}$, one of the disjuncts holds, say $\mathcal{M} \models \exists \bar{x}_j \psi_{i,\bar{a}}(\bar{x}_j, \bar{a})$. Let $\bar{b}_{\bar{a}}$ be a witness giving $\mathcal{M} \models \psi_{i,\bar{a}}(\bar{b}_{\bar{a}}, \bar{a})$. By [Mon] Lemma II.62, there is a Π_{β_i} formula $\varphi_{\bar{a}}$ that isolates the Π_{β_i} type of $\bar{b}_{\bar{a}}, \bar{a}$ in \mathcal{M} . Consider the formula,

$$\theta(\bar{x}) := \bigvee_{\bar{a} \in S} \exists \bar{y} \varphi_{\bar{a}}(\bar{y}, \bar{x}).$$

This is a Σ_α formula. If $\bar{a} \in S$ then $\mathcal{M} \models \theta(\bar{a})$ by construction. Moreover, if $\mathcal{M} \models \theta(\bar{c})$ then there is some witness $\bar{a} \in S$ and $\bar{d} \in \mathcal{M}$ such that $\mathcal{M} \models \varphi_{\bar{a}}(\bar{d}, \bar{c})$. This means that $\bar{d}, \bar{c} \geq_{\beta_i} \bar{b}_{\bar{a}}, \bar{a}$ and so $\mathcal{M} \models \psi_{i,\bar{a}}(\bar{d}, \bar{c})$. In particular, this gives that $\bar{c} \in S$, or what is the same, $\mathcal{M} \models \varphi(\bar{c})$ as desired. \square

We can now prove our type-omitting theorem.

Theorem 7.6. *Let \mathcal{M} be a countable structure. Let $(\Gamma_i)_{i \in \omega}$ be a list of Π_α -types which are not Σ_α -supported in \mathcal{M} . Let χ be a $\forall \mathfrak{E}_\alpha$ sentence true in \mathcal{M} . Then there is \mathcal{N} such that $\mathcal{N} \models \chi$ and \mathcal{N} omits all of the Γ_i .*

We begin by noting that, by Lemma 7.5, if Γ_i is not Σ_α -isolated, then it is not \mathfrak{E}_α isolated either. This is essentially the only step of this proof that differs from the proof of Montalbán's type omitting theorem from [Mon15]. The key point is that by expanding the class of formulas considered, the same proof goes through but we obtain Theorem 1.7 as a consequence.

To obtain Theorem 1.7 from Theorem 7.6, we must note that in Proposition 1.5 we defined, given \mathcal{M} , a $\bar{\mathfrak{E}}_\alpha$ sentence $\varphi_{\mathcal{M},\alpha}$ such that for any structure \mathcal{N} ,

$$\mathcal{N} \models \varphi_{\mathcal{M},\alpha} \iff \mathcal{N} \leq_\alpha \mathcal{M}.$$

Looking at the proof, $\varphi_{\mathcal{M},\alpha}$ is actually a countable conjunction of \mathfrak{E}_α formulas and hence $\forall \mathfrak{E}_\alpha$. Then to obtain Theorem 1.7 we apply Theorem 7.6 with $\varphi_{\mathcal{M},\alpha}$ as χ . Gonzalez had previously given a proof of Theorem 1.7 which differed significantly from Montalbán's; by using the right class of formulas, we obtain a simple proof of this stronger result. We include the full proof of Theorem 7.6 for completeness (in part because it does not appear in full in [Mon15]).

Proof. Write χ as $\bigwedge_{i \in \omega} \forall \bar{x}_i \varphi_i(\bar{x}_i)$ with $\varphi_i \in \mathfrak{E}_\alpha$. We build a set of \mathfrak{E}_α formulas T over the vocabulary of \mathcal{M} enriched with countably many constants C to perform a Henkin-like construction. We insist on the following properties:

- (1) If $\bigvee \psi_i \in T$, then for some i , $\psi_i \in T$.
- (2) If $\exists \bar{y} \psi(\bar{y}) \in T$, then $\psi(\bar{c}) \in T$ for some constants $\bar{c} \in C$.
- (3) If $\bigwedge \psi_i \in T$, then for all i , $\psi_i \in T$.
- (4) If $\forall \bar{y} \psi(\bar{y}) \in T$, then $\psi(\bar{c}) \in T$ for all $\bar{c} \in C$.
- (5) For every atomic sentence ψ over $\tau \cup C$, either $\psi \in T$ or $\neg \psi \in T$.
- (6) For every i and tuple \bar{c} of length $|\bar{x}_i|$, $\varphi_i(\bar{c}) \in T$.
- (7) For every tuple \bar{c} of length $|\bar{z}|$, there is a $\theta \in \Phi(\bar{z})$ such that $\neg \theta(\bar{c}) \in T$.

Along the way, at each stage s , we will make sure that T is satisfiable in the sense that there is some interpretation $\nu_s : C \rightarrow \mathcal{M}$ that assigns the constants we have used so far to a set of elements that satisfy all of the formulas in T that we have asserted about these constants. If we can do this, the claim is shown, as the Henkin model will satisfy χ by item (6) and omit $\Phi(\bar{z})$ by item (7).

We begin with empty sets T_0, C_0 , and ν_0 . At each stage we are given C_s a finite subset of C , T_s , a finite set of \mathfrak{E}_α formulas only mentioning constants C_s , and $\nu_s : C_s \rightarrow \mathcal{M}$ with the property that $\mathcal{M} \models T_s(\nu_s(C_s))$. At each stage, we address one of the properties (1) – (7), one instance at a time. We describe below how T_s, C_s , and ν_s are modified to achieve this.

- (1) Property (1). Given $\forall \psi_i \in T_s$ we know that $\mathcal{M} \models \forall \psi_i(\nu_s(C_s))$. Pick a j with the property that $\mathcal{M} \models \psi_j(\nu_s(C_s))$ and add ψ_j to T_s to make T_{s+1} . Let $\nu_s = \nu_{s+1}$ and $C_{s+1} = C_s$. It is straightforward to confirm that the desired properties are maintained.
- (2) Property (2). Given $\exists \bar{y}\psi(\bar{y}) \in T_s$ we know that $\mathcal{M} \models \exists \bar{y}\psi(\bar{y}, \nu_s(C_s))$. Let $\bar{a} \in \mathcal{M}$ have the property that $\mathcal{M} \models \psi(\bar{a}, \nu_s(C_s))$. Take new constants $\bar{d} \in C$ and let $C_{s+1} = C_s \cup \{\bar{d}\}$ and $T_{s+1} = T_s \cup \{\psi(\bar{d})\}$. Extend ν_s by defining $\nu_{s+1}(\bar{d}) = \bar{a}$. It is straightforward to confirm that the desired properties are maintained.
- (3) Property (3). Given $\bigwedge \psi_i \in T_s$ and $i \in \omega$ let $T_{s+1} = T_s \cup \{\psi_i\}$, $C_{s+1} = C_s$ and $\nu_{s+1} = \nu_s$. It is straightforward to confirm that the desired properties are maintained.
- (4) Property (4). Given $\forall \bar{y}\psi(\bar{y}) \in T_s$ and $\bar{c} \in C_s$ of the appropriate length $T_{s+1} = T_s \cup \{\psi(\bar{c})\}$, $C_{s+1} = C_s$ and $\nu_{s+1} = \nu_s$. It is straightforward to confirm that the desired properties are maintained.
- (5) Property (5). For any atomic sentence ψ and $\bar{c} \in C_s$ of the appropriate length check if $\mathcal{M} \models \psi(\nu_s(\bar{c}))$ or if $\mathcal{M} \models \neg\psi(\nu_s(\bar{c}))$. In the former case, add $\psi(\bar{c})$ to T_s to obtain T_{s+1} and in the later case add $\neg\psi(\bar{c})$ to T_s to obtain T_{s+1} . Either way, let $C_{s+1} = C_s$ and $\nu_{s+1} = \nu_s$. It is straightforward to confirm that the desired properties are maintained.
- (6) Property (6). Given φ_i and $\bar{c} \in C_s$ of the appropriate length $T_{s+1} = T_s \cup \{\varphi_i(\bar{c})\}$, $C_{s+1} = C_s$ and $\nu_{s+1} = \nu_s$. It is straightforward to confirm that the desired properties are maintained.
- (7) Property (7). This is the key property of the type omitting argument and the only property that necessitates shifting ν_s from its already established action to obtain ν_{s+1} . This step follows the standard type omitting playbook in a Henkin construction. Fix a tuple $\bar{c} \in C_s$ of length $|\bar{z}|$ and let \bar{d} be the elements of C_s that are not among \bar{c} . We can write $\bigwedge T_s = \rho(\bar{c}, \bar{d})$ where ρ is a \mathfrak{E}_α formula. Note that $\exists \bar{y}\rho(\bar{c}, \bar{y})$ is also a \mathfrak{E}_α formula that is satisfied in \mathcal{M} by $\nu_s(\bar{c})$. In particular, because $\Phi(\bar{z})$ is not \mathfrak{E}_α supported, there is some $\bar{a} \in \mathcal{M}$ that satisfies $\exists \bar{y}\rho(\bar{a}, \bar{y})$ but does not have type $\Phi(\bar{z})$. Let $C_{s+1} = C_s$ and let $A_{s+1} = A_s \cup \{-\theta(\bar{c})\}$ where $\theta \in \Phi(\bar{z})$ has the property that $\mathcal{M} \models \neg\theta(\bar{a})$. Note that $-\theta \in \mathfrak{E}_\alpha$ as $\theta \in \mathfrak{A}_\alpha$, so this is an allowable extension of T . Finally let $\nu_{s+1}(\bar{c}) = \bar{a}$ and let $\nu_{s+1}(\bar{d}) = \bar{b}$ where $\mathcal{M} \models \rho(\bar{a}, \bar{b})$. It is straightforward to confirm that the desired properties are maintained.

With this, we can achieve a model with properties (1) – (7), and therefore the proof is complete. \square

We now show that having a $\forall \mathfrak{E}_\alpha$ Scott sentence is equivalent to having one that is $\Pi_{\alpha+1}$. This adds to a long list of equivalent conditions for having $SR(\mathcal{M}) \leq \alpha$ (see Theorem 1.1 of [Mon15]). Parts of the proof below, in particular (1) \implies (2) and (2) \implies (3), are very similar to the corresponding parts of the proof of that theorem.

Theorem 7.7. *The following are equivalent when the classes \mathfrak{A} and \mathfrak{E} are restricted to $\mathcal{L}_{\omega_1, \omega}$:*

- (1) \mathcal{M} has a $\forall \mathfrak{E}_\alpha$ Scott sentence.
- (2) The \mathfrak{A}_α type of every tuple in \mathcal{M} is supported by an \mathfrak{E}_α formula.
- (3) Every automorphism orbit of every tuple in \mathcal{M} is definable by a \mathfrak{E}_α formula.
- (4) \mathcal{M} has a $\Pi_{\alpha+1}$ Scott sentence, i.e., $SR(\mathcal{M}) \leq \alpha$.

Proof. We begin by showing that (1) \implies (2). Say that \mathcal{M} has a \mathfrak{A}_α type $\Phi(\bar{z})$ that is not supported by any \mathfrak{E}_α formula. Note that an \mathfrak{A}_α type is determined by its Π_α restriction by Proposition 1.6. Also, note that $\Phi(\bar{z})$ is not supported by any Σ_α formula. By Theorem 7.6 any $\chi \in \forall \mathfrak{E}_\alpha$ is also true of a model \mathcal{N} that omits Φ and therefore is not isomorphic to \mathcal{M} .

We now move to prove that (2) \implies (3). Let $\varphi_{\bar{a}}(\bar{x})$ be the \mathfrak{E}_α formula that supports the \mathfrak{A}_α type of \bar{a} . We claim that $\varphi_{\bar{a}}(\bar{x})$ defines the automorphism orbit of \bar{a} . To demonstrate this,

we claim that the set $T := \{(\bar{a}, \bar{b}) \mid \mathcal{M} \models \varphi_{\bar{a}}(\bar{b})\}$ is a back-and-forth set. As $\varphi_{\bar{a}}$ supports the whole \mathfrak{A}_α type of \bar{a} it is certainly the case that for $(\bar{a}, \bar{b}) \in T$, \bar{a} and \bar{b} have the same atomic diagram. Now consider $c \in \mathcal{M}$; we find a $d \in \mathcal{M}$ such that $\mathcal{M} \models \varphi_{\bar{a}, c}(\bar{b}, d)$. In other words, we must show that $\mathcal{M} \models \exists x \varphi_{\bar{a}, c}(\bar{b}, x)$. If this does not hold, we have that $\mathcal{M} \models \neg \exists x \varphi_{\bar{a}, c}(\bar{b}, x)$. Note that this is a \mathfrak{A}_α formula, therefore, it is implied by $\varphi_{\bar{a}}(\bar{b})$ which isolates the \mathfrak{A}_α type of \bar{b} . However, this means that $\mathcal{M} \models \neg \exists x \varphi_{\bar{a}, c}(\bar{a}, x)$ as $\mathcal{M} \models \varphi_{\bar{a}}(\bar{a})$. That said, $\mathcal{M} \models \exists x \varphi_{\bar{a}, c}(\bar{a}, x)$ as witnessed by c , a contradiction. Therefore, there is such a d as required.

To see the “forth” part of the argument, it is enough to show that $\mathcal{M} \models \varphi_{\bar{a}}(\bar{b}) \iff \varphi_{\bar{b}}(\bar{a})$ and appeal to a symmetrical argument as the one above. Say that $\mathcal{M} \models \varphi_{\bar{a}}(\bar{b}) \wedge \neg \varphi_{\bar{b}}(\bar{a})$. As $\neg \varphi_{\bar{b}}$ is \mathfrak{A}_α , we obtain that it is implied by $\varphi_{\bar{a}}$ which supports the whole \mathfrak{A}_α type in \mathcal{M} . This is a contradiction as it yields $\mathcal{M} \models \neg \varphi_{\bar{b}}(\bar{b})$. Therefore, T is a back-and-forth set and any \bar{b} with $\mathcal{M} \models \varphi_{\bar{a}}(\bar{b})$ is automorphic to \bar{a} .

We now show that (3) \implies (4). This is just the same as showing that there is a Σ_α description of any automorphism orbit in \mathcal{M} if there is a \mathfrak{E}_α description of any automorphism orbit in \mathcal{M} . This follows immediately from Lemma 7.5.

Lastly, we note that (4) \implies (1). If $\text{SR}(\mathcal{M}) \leq \alpha$ then \mathcal{M} has a $\Pi_{\alpha+1}$ Scott sentence. As every $\Pi_{\alpha+1}$ formula is $\forall \mathfrak{E}_\alpha$, it follows that \mathcal{M} also has a $\forall \mathfrak{E}_\alpha$ Scott sentence. \square

Corollary 1.8. *Let \mathcal{M} be a countable structure.*

(1) *If, for all countable \mathcal{N} ,*

$$\mathcal{M} \leq_\alpha \mathcal{N} \implies \mathcal{M} \cong \mathcal{N}$$

then \mathcal{M} has a $\Pi_{\alpha+2}$ Scott sentence.

(2) *If, for all countable \mathcal{N} ,*

$$\mathcal{N} \leq_\alpha \mathcal{M} \implies \mathcal{M} \cong \mathcal{N}$$

then \mathcal{M} has a $\Pi_{\alpha+1}$ Scott sentence.

Proof. For (1), $\mathcal{M} \leq_\alpha \mathcal{N} \implies \mathcal{M} \cong \mathcal{N}$. There is a $\Pi_{\alpha+2}$ sentence defining the set of \mathcal{N} such that $\mathcal{M} \leq_\alpha \mathcal{N}$. By assumption, this is a Scott sentence.

For (2), suppose that $\mathcal{N} \leq_\alpha \mathcal{M} \implies \mathcal{M} \cong \mathcal{N}$. Then, by Theorem 1.7, there is a structure $\mathcal{N} \leq_\alpha \mathcal{M}$ such that \mathcal{N} omits every Π_α type which is not Σ_α -supported in \mathcal{M} . Thus $\mathcal{N} \cong \mathcal{M}$ and every Π_α type realized in \mathcal{M} is Σ_α -supported in \mathcal{M} . Thus \mathcal{M} has a $\Pi_{\alpha+1}$ Scott sentence. \square

REFERENCES

- [ACHT] Rachael Alvir, Barbara Csima, and Matthew Harrison-Trainor. On the computability of optimal Scott sentences. preprint.
- [AGHTT21] Rachael Alvir, Noam Greenberg, Matthew Harrison-Trainor, and Dan Turetsky. Scott complexity of countable structures. *J. Symb. Log.*, 86(4):1706–1720, 2021.
- [AK90] C. J. Ash and J. F. Knight. Pairs of recursive structures. *Ann. Pure Appl. Logic*, 46(3):211–234, 1990.
- [AK00] C.J. Ash and J. Knight. *Computable Structures and the Hyperarithmetical Hierarchy*. Elsevier Science, 2000.
- [AKMC20] Rachael Alvir, Julia F. Knight, and Charles McCoy CSC. Complexity of Scott sentences. *Fund. Math.*, 251(2):109–129, 2020.
- [Ehr61] A. Ehrenfeucht. An application of games to the completeness problem for formalized theories. *Fund. Math.*, 49:129–141, 1960/61.
- [Fra50] Roland Fraïssé. Sur une nouvelle classification des systèmes de relations. *C. R. Acad. Sci. Paris*, 230:1022–1024, 1950.
- [GHK⁺05] Sergey Goncharov, Valentina Harizanov, Julia Knight, Charles McCoy, Russell Miller, and Reed Solomon. Enumerations in computable structure theory. *Ann. Pure Appl. Logic*, 136(3):219–246, 2005.
- [GHT] David Gonzalez and Matthew Harrison-Trainor. Scott spectral gaps are bounded for linear orderings. Preprint: <https://arxiv.org/pdf/2411.12084>.

- [HM14] Kenneth Harris and Antonio Montalbán. Boolean algebra approximations. *Trans. Amer. Math. Soc.*, 366(10):5223–5256, 2014.
- [Kar65] Carol R. Karp. Finite-quantifier equivalence. In *Theory of Models (Proc. 1963 Internat. Sympos. Berkeley)*, pages 407–412. North-Holland, Amsterdam, 1965.
- [Kec95] Alexander S. Kechris. *Classical descriptive set theory*, volume 156 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995.
- [LE66] E. G. K. Lopez-Escobar. On defining well-orderings. *Fund. Math.*, 59:13–21, 1966.
- [LR87] A. Louveau and J. Saint Raymond. Borel classes and closed games: Wadge-type and hurewicz-type results. *Transactions of the American Mathematical Society*, 304(2):431–467, 1987.
- [Mon] Antonio Montalbán. Computable structure theory: Beyond the arithmetic. draft.
- [Mon14] Antonio Montalbán. Priority arguments via true stages. *Journal of Symbolic Logic*, 79(4):1315–1335, 2014.
- [Mon15] A. Montalbán. A robust Scott rank. *Proc. Amer. Math. Soc.*, 143(12):5427–5436, 2015.
- [Mon21] Antonio Montalbán. *Computable Structure Theory: Within the Arithmetic*. Perspectives in Logic. Cambridge University Press, 2021.
- [MR23] Antonio Montalbán and Dino Rossegger. The structural complexity of models of arithmetic. *Journal of Symbolic Logic*, To appear:1–17, 2023.
- [Nad74] Mark Nadel. Scott sentences and admissible sets. *Ann. Math. Logic*, 7:267–294, 1974.
- [Sco65] Dana Scott. Logic with denumerably long formulas and finite strings of quantifiers. In *Theory of Models (Proc. 1963 Internat. Sympos. Berkeley)*, pages 329–341. North-Holland, Amsterdam, 1965.
- [Vau75] Robert Vaught. Invariant sets in topology and logic. *Fund. Math.*, 82:269–294, 1974/75.