

CONFLICT-FREE HYPERGRAPH MATCHINGS AND COVERINGS

FELIX JOOS, DHURV MUBAYI, AND ZAK SMITH

ABSTRACT. Recent work showing the existence of conflict-free almost-perfect hypergraph matchings has found many applications. We show that, assuming certain simple degree and codegree conditions on the hypergraph $\mathcal{H}$ and the conflicts to be avoided, a conflict-free almost-perfect matching can be extended to one covering all vertices in a particular subset of $V(\mathcal{H})$, by using an additional set of edges; in particular, we ensure that our matching avoids all additional conflicts, which may consist of both old and new edges.

This setup is useful for various applications in design theory and Ramsey theory. For example, our main result provides a crucial tool in the recent proof of the high girth existence conjecture due to Delcourt and Postle. It also provides a black box which encapsulates many long and tedious calculations, greatly simplifying the proofs of results in generalised Ramsey theory.

1. INTRODUCTION

Hypergraph matching problems can be used to model various central questions in combinatorics, and consequently have been studied for many years. Notably, Frankl and Rödl [11], as well as Pippenger [17], proved that any k -uniform hypergraph (k fixed) on n vertices, in which each vertex belongs to roughly d edges and any pair of vertices belongs to at most $o_d(d)$ edges, contains a matching covering $(1 - o_d(1))n$ vertices. These theorems have undoubtedly had a vast number of applications.

More recently, Delcourt and Postle [7], as well as Glock, Joos, Kim, Kühn, and Lichev [12], generalised this result by introducing so-called conflict-free matchings. Here a conflict is a set of disjoint edges which is forbidden to be a subset of the matching. The main contribution in [7, 12] is as follows: under the same conditions as Frankl, Rödl, and Pippenger, and under sensible conditions on the set of conflicts, one can find an almost perfect matching that avoids all conflicts. This can again be applied to a number of problems, such as high girth decompositions, problems considered by Brown, Erdős and Sós [4] and various questions in generalised Ramsey theory [1, 3, 13, 14, 16].

The unfortunate drawback of these theorems is that they deal only with almost-perfect matchings, whereas in many applications it is desirable to obtain perfect matchings or at least almost perfect matchings that cover a specified vertex subset entirely. Delcourt and Postle describe a setup in which this obstacle can be overcome in a particular setting. Specifically, they consider hypergraphs which are bipartite in the sense that the vertex set can be partitioned into two parts A and B with each edge containing exactly one vertex from A (and they assume that the vertex degrees in A are slightly higher than in B), in which case they find a conflict-free matching covering all vertices of A (the matching is *A-perfect*). The primary limitation of the bipartite hypergraphs in [7] is the fact that edges are only allowed to contain exactly one vertex from A , whereas in several applications this is not the case.

Our main contribution is to obtain a stronger theorem in a more general setup: we work with a ‘tripartite’ hypergraph for which the vertex set has a partition into three sets P, Q, R , and we seek a P -perfect matching. The edge set is divided into two parts: edges containing p vertices from P and q vertices from Q , and edges containing one vertex from P and r vertices from R . In this setup we prove the existence of a P -perfect matching, avoiding conflicts which may consist of both types of edges (almost all vertices in P are covered by edges of the first

Date: 7th November 2025.

The research leading to these results was partially supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – 428212407 (F. Joos and Z. Smith).




FIGURE 1. Two matchings $\mathcal{M}_1 \subseteq \mathcal{H}_1$ and $\mathcal{M}_2 \subseteq \mathcal{H}_2$ in the hypergraph $\mathcal{H}$, whose union forms a P -perfect matching $\mathcal{M}$; here $p = 2, q = 4, r = 2$.

type and a few vertices in P are covered by edges of the second type). The proof generalises a ‘two-stage method’ for constructions in generalised Ramsey theory, first introduced by Bennett, Cushman, Dudek, and Prałat [2], and subsequently used to prove various other results in this area; our theorem simplifies all such proofs significantly so that all technical computations are no longer needed, and mostly back-of-the-envelope calculations are sufficient.

This framework also turns out to be useful in other settings, including high-girth coverings and designs; these applications are discussed further in Section 2. We expect that there will be further applications in future.

1.1. Tripartite Matching Theorem. Given two hypergraphs $\mathcal{H}$ and $\mathcal{C}$, say that $\mathcal{C}$ is a *conflict hypergraph* for $\mathcal{H}$ if $V(\mathcal{C}) = E(\mathcal{H})$, and in this case call the edges of $\mathcal{C}$ *conflicts*. Say that a set of edges $E \subseteq \mathcal{H}$ is $\mathcal{C}$ -free if it does not contain any conflict from $\mathcal{C}$. Suppose that we are given the following setup:

- (S1) integers $\ell \geq 2, d > 0, p \geq 1, q \geq 0, r \geq 1$ with $p + q \geq 2$ and real $\varepsilon > 0$;
- (S2) disjoint sets P, Q, R with $d^\varepsilon \leq |P| \leq |P \cup Q| \leq \exp(d^{\varepsilon^3})$;
- (S3) hypergraph $\mathcal{H}_1$ whose edges consist of p vertices from P and q vertices from Q ;
- (S4) hypergraph $\mathcal{H}_2$ whose edges consist of a single vertex from P and r vertices from R ;
- (S5) conflict hypergraph $\mathcal{C}$ for $\mathcal{H}_1$;
- (S6) conflict hypergraph $\mathcal{D}$ for $\mathcal{H} := \mathcal{H}_1 \cup \mathcal{H}_2$.

Assume that $\mathcal{H}$ satisfies suitable degree conditions, and further that both $\mathcal{C}$ and $\mathcal{D}$ satisfy suitable boundedness conditions, all of which are specified in Section 3 in terms of d and ε .

Theorem 1.1. *Given p, q, r, ℓ as above, there exists $\varepsilon_0 > 0$ such that for all $\varepsilon \in (0, \varepsilon_0)$, there exists d_0 such that given the above setup, the following holds for all $d \geq d_0$: there exists a P -perfect $\mathcal{C} \cup \mathcal{D}$ -free matching $\mathcal{M} \subseteq \mathcal{H}$. Furthermore, at most $d^{-\varepsilon^4}|P|$ vertices of P belong to an edge in $\mathcal{H}_2 \cap \mathcal{M}$.*

The proof of Theorem 1.1, given in Section 5, consists of two stages, as mentioned: we first apply Theorem 4.2, a variant of the original conflict-free hypergraph matching theorem [12], to the hypergraph $\mathcal{H}_1$ in order to obtain a $\mathcal{C}$ -free matching $\mathcal{M}_1$ covering all but at most a $d^{-\varepsilon^3}$ fraction of the vertices of P with edges from $\mathcal{H}_1$. In the proof of Theorem 4.2 (which we apply as a black box), this matching is chosen randomly one edge at a time, and accordingly can be thought of as ‘pseudorandom’ in an appropriate sense; specifically, we are able to show that certain weight functions on edge sets have roughly their ‘expected’ value when summed over $\mathcal{M}_1$. To extend $\mathcal{M}_1$ to a P -perfect matching, for each vertex $x \in P$ which is not already covered, we randomly choose some edge from $\mathcal{H}_2$ to cover x , and use the Lovász Local Lemma to show that with non-zero probability the resulting set $\mathcal{M}_2$ of edges is indeed a matching, and the union $\mathcal{M}$ of our two matchings is $\mathcal{D}$ -free. In order to apply the local lemma, we use

the aforementioned weight functions to bound the number of potential conflicts which could be introduced when choosing an edge from $\mathcal{H}_2$ to cover each $x \in P$.

1.2. Notation. Write $[i, n] = \{i, \dots, n\}$, so $[n] = [1, n]$. Unless otherwise stated we identify hypergraphs with their edge sets, writing $e \in \mathcal{G}$ to mean $e \in E(\mathcal{G})$. Given a set of vertices $U \subseteq V(\mathcal{G})$ in a hypergraph $\mathcal{G}$, write $d_{\mathcal{G}}(U)$ for the degree of U in $\mathcal{G}$, that is the number of edges of $\mathcal{G}$ containing U ; in the cases $U = \{u\}$ and $U = \{u, v\}$, where U consists of one or two vertices, we just write $d_{\mathcal{G}}(u)$ and $d_{\mathcal{G}}(u, v)$ respectively. We omit the subscript if $\mathcal{G}$ is obvious from context. Write $\Delta_j(\mathcal{G})$ for the maximum degree $d_{\mathcal{G}}(U)$ among sets $U \subseteq V(\mathcal{G})$ of j vertices. Given a subset of the vertices $V \subseteq V(\mathcal{G})$, write $\Delta_V(\mathcal{G})$ for the maximum degree $d_{\mathcal{G}}(u)$ of any single vertex $u \in V$, and similarly $\delta_V(\mathcal{G})$ for the minimum degree; assume $V = V(\mathcal{G})$ if not specified. Given $j \in \mathbb{N}$, write $\mathcal{G}^{(j)} := \{E \in \mathcal{G} : |E| = j\}$ for the subhypergraph of $\mathcal{G}$ containing only those edges of size j .

We omit ceiling and floor symbols whenever they do not affect the argument. We refer readers to the Glossary section at the end of the paper for a summary of more specific notation which appears throughout the paper.

1.3. This paper. We start by outlining two significant applications of Theorem 1.1 in Section 2. We then complete the formal statement in Section 3 by listing the various conditions required on our hypergraph and conflicts. In Section 4, we state our prerequisites, in particular Theorem 4.2, a variant of the main theorem from [12], as well as a more complicated but weaker set of conditions, for which we will in fact prove our main theorem. Finally, the proof itself is given in Section 5, which begins with a more detailed outline.

2. APPLICATIONS

We briefly discuss here two applications of Theorem 1.1.

2.1. Conflict-free coverings and large girth designs. For a hypergraph $\mathcal{H}$, we call a set of edges $\mathcal{M} \subseteq \mathcal{H}$ a *covering* of $\mathcal{H}$ if all vertices belong to some edge in $\mathcal{M}$, and say it is *perfect* if each vertex belongs to exactly one edge. In the setting of Frankl, Rödl and Pippenger, the existence of almost perfect matchings and almost perfect coverings (coverings in which most vertices belong to exactly one edge) is equivalent and one object can easily be transformed into the other one. However, it is not obvious that just greedily adding edges to turn an almost perfect matching into an almost perfect covering can be done without introducing conflicts; Theorem 2.1, which follows easily from Theorem 1.1, resolves this problem. It represents the natural analogue of Theorem 4.1, providing a covering in place of a matching.

Theorem 2.1. *Fix $\ell, k \geq 2$. There exists ε_0 such that for all $\varepsilon \in (0, \varepsilon_0)$, there exists d_0 such that for all $d \geq d_0$ the following holds. Suppose $\mathcal{H}$ is a k -graph on $n \leq \exp(d^{\varepsilon^3})$ vertices. Assume that $(1 - d^{-\varepsilon})d \leq \delta(\mathcal{H}) \leq \Delta(\mathcal{H}) \leq d$ and $\Delta_2(\mathcal{H}) \leq d^{1-\varepsilon}$. Let $\mathcal{C}$ be a (d, ℓ, ε) -bounded conflict hypergraph for $\mathcal{H}$. Then there is a $\mathcal{C}$ -free covering $\mathcal{M} \subseteq \mathcal{H}$ such that all but $d^{-\varepsilon^5}n$ vertices are covered exactly once, and no vertex is covered more than twice.*

One of the original motivations for studying hypergraph matchings, and in particular conflict-free matchings, was the problem of finding almost-perfect Steiner systems of large girth. In general, a *partial (m, s, t) -Steiner system* is a collection $\mathcal{S}$ of subsets of $[m]$, each of size s , such that every subset of $[m]$ of size t is contained in at most one element of $\mathcal{S}$; it is *approximate* if it has size $(1 - o(1))\binom{m}{t}/\binom{s}{t}$. The *girth* of $\mathcal{S}$ is the smallest integer $g \geq 2$ such that some set of $(s - t)g + t$ vertices induces at least g sets in $\mathcal{S}$.

Recently, Delcourt and Postle [8] proved the existence of perfect (m, s, t) -Steiner systems of large girth, via a new *refined absorption* method. Their result provides a common generalization of the existence conjecture for designs originating from the 1800s and Erdős' conjecture from 1973 on the existence of high girth Steiner triple systems. Our main result, Theorem 1.1, is a crucial tool in their proof (Theorem 2.10 in [8]). As noted earlier, our proof method uses the main result of [12] which employs the random greedy process. On the other hand, Delcourt and Postle's proof (to appear in a forthcoming version of [7]) of their Theorem 2.10 from [8]

uses the nibble method. A simpler version of Theorem 1.1, without conflicts, is also used by Delcourt and Postle [9], as well as by Delcourt, Postle, and Kelly [5, 6] in other applications of their refined absorption method.

Conflict-free hypergraph matchings can also be used to find approximate systems for a much more general class of quasirandom hypergraphs, where, for example, we restrict the choice of elements of $\mathcal{S}$ to a randomly chosen subset of $\binom{[m]}{s}$; see Theorem 1.4 of [12]. The following analogous covering result can be easily deduced from our Theorem 2.1. Given real numbers a, b, c , we write $a = b \pm c$ to mean $b - c \leq a \leq b + c$.

Theorem 2.2. *For all $c_0 > 0, \ell \geq 2$ and $s > t \geq 2$, there exists $\varepsilon_0 > 0$ such that for all $\varepsilon \in (0, \varepsilon_0)$, there exists m_0 such that the following holds for all $m \geq m_0$ and $c \geq c_0$. Let G be a t -graph on m vertices and let $\mathcal{K}$ be a collection of sets of size s which induce cliques in G such that any edge is contained in $(1 \pm m^{-\varepsilon})cm^{s-t}$ elements of $\mathcal{K}$.*

Then, there exists a subset $\mathcal{S} \subseteq \mathcal{K}$ such that every edge of G is contained in at least one element of $\mathcal{S}$, the proportion of edges of G contained in more than one element of $\mathcal{S}$ is $o_m(1)$, none are contained in more than two, and any subset of $\mathcal{S}$ of size j , where $j \in [2, \ell]$, whose elements have pairwise intersections of size at most $t - 1$, spans more than $(s - t)j + t$ points.

2.2. Generalised Ramsey numbers. Given graphs G and H , and $q \in \mathbb{N}$, define the *generalised Ramsey number* $r(G, H, q)$ to be the minimum number of colours needed to colour the edges of G in such a way that every copy of H receives at least q distinct colours. Bennett, Cushman, Dudek, and Prałat [2] showed that $r(K_n, K_4, 5) = 5n/6 + o(n)$, answering a question of Erdős and Gyárfás, by introducing the aforementioned two-stage method in which they first colour most of the edges of K_n using a modified triangle removal process (requiring complicated technical analysis), and then complete this to a full colouring using the Lovász Local Lemma. Joos and Mubayi [14] simplified this method greatly by encoding the first stage as a suitable conflict-free hypergraph matching problem and applying the main theorem from [12] as a black box, and demonstrated its versatility further by showing that $r(K_n, C_4, 3) = n/2 + o(n)$. This approach has subsequently been used to prove various similar and more general results [1, 3, 13, 14, 16].

Our main theorem (Theorem 1.1) formalises this two-stage method in a single statement, from which all of these colouring results follow; another main contribution of this paper is to consolidate all of the calculations required for the second stage, so that applications need only focus on constructing appropriate hypergraphs and conflicts satisfying our conditions. This massively simplifies the proofs of all existing results, since it now suffices to only check the orders of magnitude of the numbers of different types of conflicts. For example, one may obtain a concise proof of the following result which was stated (without proof) very recently by Bal, Bennett, Heath, and Zerbib [1]. Given $k \geq 2$, write K_n^k for the complete k -graph on n vertices, and C_ℓ^k for the k -uniform tight cycle of length ℓ ; that is, edges $e_1, \dots, e_\ell$ on vertices $v_1, \dots, v_\ell$ such that $e_i = \{v_i, \dots, v_{i+k-1}\}$ (modulo ℓ) for each $i \in [\ell]$.

Theorem 2.3. *For all $k \geq 2$ and $\ell \geq k + 2$, we have $r(K_n^k, C_\ell^k, k + 1) \leq n/(\ell - k) + o(n)$.*

We remark that the factor $\ell - k$ above is best possible assuming a well-known conjecture about the Turán number of tight paths in hypergraphs.

The proofs of Theorems 2.1, 2.2 and 2.3 are fairly routine and short. For the convenience of the reader we provide them in the arXiv version of this paper in the appendix [15].

3. CONDITIONS FOR THE MAIN THEOREM

We now specify the conditions on $\mathcal{H}, \mathcal{C}$, and $\mathcal{D}$ required for Theorem 1.1 to hold, thus completing its formal statement. Throughout this section, we work in the setup from Section 1.1, recalling (S1)–(S6).

3.1. Degree conditions on $\mathcal{H}$. We require that the hypergraph $\mathcal{H} = \mathcal{H}_1 \cup \mathcal{H}_2$ with $\mathcal{H}_1, \mathcal{H}_2 \neq \emptyset$ satisfies the following conditions.

(H1) $(1 - d^{-\varepsilon})d \leq \delta_P(\mathcal{H}_1) \leq \Delta(\mathcal{H}_1) \leq d$;

- (H2) $\Delta_2(\mathcal{H}_1) \leq d^{1-\varepsilon}$;
- (H3) $\Delta_R(\mathcal{H}_2) \leq d^{\varepsilon^4} \delta_P(\mathcal{H}_2)$;
- (H4) $d(x, v) \leq d^{-\varepsilon} \delta_P(\mathcal{H}_2)$ for each $x \in P$ and $v \in R$.

This means that $\mathcal{H}_1$ is essentially regular for vertices in P and has small codegrees, although vertices in Q are allowed to have much lower (but not higher) degrees. Meanwhile in $\mathcal{H}_2$, every vertex in P must have degree at least a $d^{-\varepsilon^4}$ proportion of the maximum degree in R , and few edges in common with any particular vertex in R . These conditions are used to ensure that we may choose an edge of $\mathcal{H}_2$ containing each $x \in P$ such that the set of edges chosen do not overlap.

3.2. Boundedness conditions on $\mathcal{C}$. The conflicts of $\mathcal{C}$ consist only of edges from $\mathcal{H}_1$, and are avoided directly by Theorem 4.2, so require the same boundedness conditions as in [12]; the conditions we present here are not the most general possible, but suffice for most applications. For any edge $e \in \mathcal{H}_1$, we write $N_{\mathcal{C}}^{(2)}(e) := \{f \in \mathcal{H}_1 : \{e, f\} \in \mathcal{C}\}$. Given $\ell \geq 2$ and $d, \varepsilon > 0$, say that $\mathcal{C}$ is (d, ℓ, ε) -bounded if

- (C1) $2 \leq |C| \leq \ell$ for all $C \in \mathcal{C}$;
- (C2) $\Delta(\mathcal{C}^{(j)}) \leq \ell d^{j-1}$ for all $j \in [2, \ell]$;
- (C3) $\Delta_{j'}(\mathcal{C}^{(j)}) \leq d^{j-j'-\varepsilon}$ for all $j \in [2, \ell]$ and $j' \in [2, j-1]$;
- (C4) $|\{f \in N_{\mathcal{C}}^{(2)}(e) : v \in f\}| \leq d^{1-\varepsilon}$ for all $e \in E(\mathcal{H}_1)$ and $v \in V(\mathcal{H}_1)$;
- (C5) $|N_{\mathcal{C}}^{(2)}(e) \cap N_{\mathcal{C}}^{(2)}(f)| \leq d^{1-\varepsilon}$ for all disjoint $e, f \in E(\mathcal{H}_1)$.

Note that, in many applications, all conflicts of $\mathcal{C}$ have size at least 3, so conditions (C4) and (C5) are vacuously true. For further intuition behind conditions (C1)–(C5), we refer the reader to the proof overview (Section 1.1) in [12].

3.3. Boundedness conditions on $\mathcal{D}$. Since it may be the case that the conflicts of $\mathcal{D}$ consist only of edges from $\mathcal{H}_2$, or of two parts from $\mathcal{H}_1$ and $\mathcal{H}_2$, they must satisfy a new set of conditions to be avoided. We extend our previous notation by writing $\mathcal{D}^{(j_1, j_2)}$ for the set of conflicts in $\mathcal{D}$ consisting of j_1 edges from $\mathcal{H}_1$ and j_2 edges from $\mathcal{H}_2$. Similarly, we now write

$$(3.1) \quad \Delta_{j'_1, j'_2}(\mathcal{D}) := \max \left\{ d_{\mathcal{D}}(F_1 \cup F_2) : F_1 \in \binom{\mathcal{H}_1}{j'_1}, F_2 \in \binom{\mathcal{H}_2}{j'_2} \right\}.$$

Given a vertex $x \in P$, write also $\mathcal{D}_x$ for the set of conflicts in $\mathcal{D}$ containing x in their $\mathcal{H}_2$ -part, and likewise $\mathcal{D}_{x,y}$ for those containing both x and y . Then say that $\mathcal{D}$ is (d, ℓ, ε) -simply-bounded if the following hold for all $x, y \in P$, and $j_1 \in [0, \ell], j_2 \in [2, \ell]$.

- (D1) $2 \leq |D \cap \mathcal{H}_2| \leq |D| \leq \ell$ for each conflict $D \in \mathcal{D}$;
- (D2) $|\mathcal{D}_x^{(j_1, j_2)}| \leq d^{j_1 + \varepsilon^4} \delta_P(\mathcal{H}_2)^{j_2}$;
- (D3) $\Delta_{j'_1, 0}(\mathcal{D}_x^{(j_1, j_2)}) \leq d^{j_1 - j'_1 - \varepsilon} \delta_P(\mathcal{H}_2)^{j_2}$ for each $j'_1 \in [j_1]$;
- (D4) $|\mathcal{D}_{x,y}^{(j_1, j_2)}| \leq d^{j_1 - \varepsilon} \delta_P(\mathcal{H}_2)^{j_2}$.

Referring to the brief proof outline in Section 1.1, the bound in (D2) can be understood heuristically as follows. For $D \in \mathcal{D}_x$ to be contained in $\mathcal{M}$, all edges of $D \cap \mathcal{H}_1$ must be chosen in $\mathcal{M}_1$ and all edges of $D \cap \mathcal{H}_2$ must be chosen in $\mathcal{M}_2$; note in particular that, for the latter to be possible, all vertices $y \in P \cap \bigcup (D \cap \mathcal{H}_2)$ must be left uncovered by $\mathcal{M}_1$. Under the heuristic that edges are chosen roughly uniformly and independently by Theorem 4.2, the probability of the j_1 edges of $D \cap \mathcal{H}_1$ all being contained in $\mathcal{M}_1$ is roughly d^{-j_1} . Under the heuristic that $\mathcal{M}_1$ covers vertices of P roughly uniformly, the probability that even one vertex $y \in P \cap \bigcup (D \cap \mathcal{H}_2) \setminus \{x\}$ is left uncovered is roughly $d^{-\varepsilon^3}$. Therefore, after choosing $\mathcal{M}_1$, we expect there to be at most $d^{\varepsilon^4 - \varepsilon^3} \delta_P(\mathcal{H}_2)^{j_2}$ conflicts $D \in \mathcal{D}_x$ which pose a threat when choosing $\mathcal{M}_2$. Ignoring issues of dependence, each of the remaining j_2 edges of D are then included in $\mathcal{M}_2$ with probability at most $\delta_P(\mathcal{H}_2)^{-1}$, so in total we expect at most $d^{\varepsilon^4 - \varepsilon^3} = o(1)$ conflicts from $\mathcal{D}_x$ to be contained in $\mathcal{M}$. Intuitively, (D3) and (D4) behave similarly to (C3), to ensure that conflicts are ‘well spread-out’ in the hypergraph, which avoids the number of potential conflicts being dominated by rare events with large effects.

We may now formally re-state Theorem 1.1.

Theorem 3.1. *Given p, q, r, ℓ satisfying (S1), there exists $\varepsilon_0 > 0$ such that for all $\varepsilon \in (0, \varepsilon_0)$, there exists d_0 such that, for all $d \geq d_0$, given $\mathcal{H}, \mathcal{C}, \mathcal{D}$ satisfying (S2)–(S6), the following holds. Assume that $\mathcal{H}$ satisfies (H1)–(H4), $\mathcal{C}$ is (d, ℓ, ε) -bounded, and $\mathcal{D}$ is (d, ℓ, ε) -simply-bounded. Then there exists a P -perfect $\mathcal{C} \cup \mathcal{D}$ -free matching $\mathcal{M} \subseteq \mathcal{H}$. Furthermore, at most $d^{-\varepsilon^4}|P|$ vertices of P belong to an edge in $\mathcal{H}_2 \cap \mathcal{M}$.*

We actually prove Theorem 3.1 for a slightly more general set of conditions, given in Section 4.3, which in particular allow for conflicts D with $j_2 = |D \cap \mathcal{H}_2| = 1$; in many applications, however, this type of conflict does not occur.

4. PREPARATION FOR THE PROOF

In this section, we begin by deducing our required variant of the conflict-free hypergraph matchings theorem of [12], as well as giving the more general set of conditions under which we prove Theorem 3.1. We then proceed with the proof itself in Section 5.

4.1. The Conflict-Free Hypergraph Matchings Theorem. To state the theorem we need, we must first make a further definition; assume ℓ is a given integer, as in Section 1.1. Recalling that we write $\mathcal{H}$ to refer to $E(\mathcal{H})$, and given $j \in \mathbb{N} \cup \{0\}$,¹ say that $w: \binom{\mathcal{H}}{j} \rightarrow [0, \ell]$ is a j -uniform ℓ -test function for $\mathcal{H}$ if $w(E) = 0$ whenever $E \in \binom{\mathcal{H}}{j}$ is not a matching. Write $w(X) := \sum_{x \in X} w(x)$ for $X \subseteq \binom{\mathcal{H}}{j}$, and $w(E) := w(\binom{E}{j})$ for general $E \subseteq \mathcal{H}$.

Given a vertex v in a hypergraph $\mathcal{H}$, the *link of v in $\mathcal{H}$* is the hypergraph $\mathcal{H}_{-v} = \{E \setminus \{v\} : E \in \mathcal{H}, v \in E\}$ on vertex set $V(\mathcal{H}) \setminus \{v\}$, that is the set of all *partial* edges which are completed by v to form an edge of $\mathcal{H}$. Now say that a pair of edges $e, f \in \mathcal{H}$ is an $(\varepsilon, \mathcal{C})$ -*conflict-sharing pair* (or just ε -*conflict-sharing pair* since $\mathcal{C}$ is usually obvious from context) if $|(C_{-e})^{(j')} \cap (C_{-f})^{(j')}| > d^{j'-\varepsilon}$ for some $j' \in [\ell - 1]$. Then given a conflict hypergraph $\mathcal{C}$ for $\mathcal{H}$, a j -uniform ℓ -test function w for $\mathcal{H}$, and $d, \varepsilon > 0$, we say that w is $(d, \varepsilon, \mathcal{C})$ -*trackable* if

- (W1) $w(\mathcal{H}) \geq d^{j+\varepsilon}$;
- (W2) $w(\{E \in \binom{\mathcal{H}}{j} : E \supseteq E'\}) \leq w(\mathcal{H})/d^{j'+\varepsilon/2}$ for all $j' \in [j - 1]$ and $E' \in \binom{\mathcal{H}}{j'}$;
- (W3) $w(E) = 0$ for any $E \in \binom{\mathcal{H}}{j}$ with $e, f \in E$ for some 2ε -conflict-sharing pair $e, f \in \mathcal{H}$;
- (W4) $w(E) = 0$ for all $E \in \binom{\mathcal{H}}{j}$ which are not $\mathcal{C}$ -free.

We will make use of trackable test functions in our proof to bound the number of mixed conflicts whose $\mathcal{H}_1$ -part is chosen in the first matching, meaning that their $\mathcal{H}_2$ -part needs to be avoided in the second matching. For some intuition behind conditions (W1)–(W4), we refer the reader again to [12], specifically the explanation of their conditions (Z1)–(Z4) in Section 3. We may now state the original theorem, a simplified variant of the main result from [12]. Note that the absence of conflict-sharing pairs in the matching is not given by any of the theorem statements in the original paper, but follows from the proof, in which such pairs are added as conflicts of size 2.²

Theorem 4.1. *For all $k, \ell \geq 2$, there exists $\varepsilon_0 > 0$ such that for all $\varepsilon \in (0, \varepsilon_0)$, there exists d_0 such that the following holds for all $d \geq d_0$. Suppose $\mathcal{H}$ is a k -graph on $n \leq 2 \exp(d^{\varepsilon^3})$ vertices with $(1 - d^{-\varepsilon})d \leq \delta(\mathcal{H}) \leq \Delta(\mathcal{H}) \leq d$ and $\Delta_2(\mathcal{H}) \leq d^{1-\varepsilon}$ and suppose $\mathcal{C}$ is a (d, ℓ, ε) -bounded conflict hypergraph for $\mathcal{H}$. Suppose also that $\mathcal{Y}$ is a set of $(d, \varepsilon/10, \mathcal{C})$ -trackable ℓ -test functions for $\mathcal{H}$ of uniformity at most ℓ with $|\mathcal{Y}| \leq \exp(d^{\varepsilon^3})$. Then, there exists a $\mathcal{C}$ -free matching*

¹Note that in the original paper j is assumed to be strictly positive, but the conclusion of the theorem still holds in the trivial case $j = 0$. This will simplify our notation later.

²See Lemmas 8.5 and 8.6 in [12], which refer to conflict-sharing pairs as *bad pairs*; note that while the statements given there only avoid $\varepsilon/4$ -conflict-sharing pairs, this can easily be improved to $\varepsilon/2$ by being more conservative with ε , provided that all test functions are zero on sets containing an $\varepsilon/2$ -conflict-sharing pair; this follows from (W3), noting that our condition (W3) uses 2ε in place of ε , unlike in the original paper. Similarly, taking such extra care allows us to require only that test functions are $(d, \varepsilon/4, \mathcal{C})$ -trackable, rather than with ε as in the original statement, and to replace ε by $\varepsilon/2$ in condition (W2).

$\mathcal{M} \subseteq \mathcal{H}$ of size at least $(1 - d^{-2\varepsilon^3})\frac{n}{k}$ with $w(\mathcal{M}) = (1 \pm d^{-\varepsilon^3})d^{-j}w(\mathcal{H})$ for all j -uniform $w \in \mathcal{Y}$. Furthermore, $\mathcal{M}$ contains no $(\varepsilon/2, \mathcal{C})$ -conflict-sharing pairs.

In order to prove Theorem 3.1, we require a slight extension, for which we make another definition; we say a j -uniform test function w is $(d, \varepsilon, \mathcal{C})$ -semi-trackable if it satisfies the alternative conditions

$$(W1^*) \quad w(\mathcal{H}) \leq d^{j+2\varepsilon};$$

$$(W2^*) \quad w(\{E \in \binom{\mathcal{H}}{j} : E \supseteq E'\}) \leq \varepsilon^{-1}d^{j-j'} \text{ for all } j' \in [j-1] \text{ and } E' \in \binom{\mathcal{H}}{j'};$$

as well as (W3) and (W4). Observe also here that, if (W1) holds, then (W2*) implies (W2), so we will usually just check (W2*) when proving that test functions are trackable.

For such functions, it is not in general possible to guarantee the same estimate $w(\mathcal{M}) = (1 \pm d^{-\varepsilon^3})d^{-j}w(\mathcal{H})$, because the heuristically expected value is too small and therefore subject to outlying events; for a simple example, let $v \in V(\mathcal{H})$, choose Z to be some set of size $d(v)/2$ of edges that contain v , and take w as the indicator function of Z , then (heuristically, imagining $\mathcal{M}$ to be appropriately pseudorandom), the expected value of $w(\mathcal{M})$ is (approximately) $1/2$ but clearly we cannot guarantee concentration close to $1/2$ since w takes values in $\{0, 1\}$. It is however obvious in this case that $w(\mathcal{M})$ can be bounded from above, namely by 1. More generally, we might hope that for semi-trackable test functions, we can make use of the bound on $w(\mathcal{H})$ to guarantee that at least $w(\mathcal{M})$ is not too large; indeed, this turns out to be possible.

Theorem 4.2. *Assume the setup of Theorem 4.1, but allow $\mathcal{Y}$ to contain also some $(d, \varepsilon/10, \mathcal{C})$ -semi-trackable ℓ -test functions. Then there exists a $\mathcal{C}$ -free matching $\mathcal{M} \subseteq \mathcal{H}$ of size at least $(1 - d^{-\varepsilon^3})\frac{n}{k}$, containing no $(\varepsilon/2, \mathcal{C})$ -conflict-sharing pairs, with $w(\mathcal{M}) = (1 \pm d^{-\varepsilon^3})d^{-j}w(\mathcal{H})$ for all j -uniform trackable $w \in \mathcal{Y}$ and $w(\mathcal{M}) \leq d^{\varepsilon/4}$ for all semi-trackable $w \in \mathcal{Y}$.*

Proof. Our strategy will be to extend $\mathcal{H}$ using some new dummy vertices and edges, allowing us to extend w to a new test function w' which agrees with w on $\mathcal{H}$, but is also positive on sufficiently many subsets of the new edges to satisfy (W1). We then obtain the usual estimate for the value of $w'(\mathcal{M})$, which in particular gives us the required crude upper bound for $w(\mathcal{M})$.

Let S be a set of m new vertices, disjoint from $V(\mathcal{H})$, with m chosen such that $(1 - d^{-\varepsilon})d \leq \binom{m-1}{k-1} \leq d$, noting that this is possible for $\varepsilon < (k-1)^{-1}$. Note that $m \leq kd^{1/(k-1)}$ and $d^\varepsilon \leq m \leq n$ since $d \leq \binom{n-1}{k-1}$. Let $\mathcal{K}$ be the complete k -graph on vertex set S , and define $\mathcal{H}'$ to be the (disjoint) union of $\mathcal{H}$ with ℓ vertex-disjoint copies of $\mathcal{K}$, say $\mathcal{K}_1, \dots, \mathcal{K}_\ell$ on vertex sets $S_1, \dots, S_\ell$, respectively. Then, by definition, $\mathcal{H}'$ is essentially regular as required for Theorem 4.1, and $\Delta_2(\mathcal{K}) \leq m^{k-2} \leq k^{k-2}d^{1-1/(k-1)} \leq d^{1-\varepsilon}$, as we can choose $\varepsilon_0 < (k-1)^{-1}/2$.

For each $i \in [\ell]$, choose any subset $T_i \subseteq S_i$ of size $t := 2d^{\varepsilon/5}$, and enumerate its vertices arbitrarily as $v_1^i, \dots, v_t^i$. For each $j \in [\ell]$, define

$$\mathcal{Z}_j := \left\{ \{e_1, \dots, e_j\} \in \binom{\mathcal{H}'}{j} : \exists s \in [t] \text{ such that } e_i \cap T_i = \{v_s^i\} \ \forall i \in [j] \right\} \subseteq \binom{\mathcal{H}'}{j}.$$

We claim that the indicator function $\mathbf{1}(\mathcal{Z}_j)$ is a $(d, \varepsilon/5, \mathcal{C})$ -trackable ℓ -test function, for each $j \in [\ell]$.

Clearly every element of $\mathcal{Z}_j$ is a matching. For (W1), observe that $d^{j+\varepsilon/5} \leq |\mathcal{Z}_j| \leq 2d^{j+\varepsilon/5}$, since there are $t = 2d^{\varepsilon/5}$ possible choices of s , and then at least $(1 - d^{-\varepsilon/3})d$ (and at most d) choices for each edge e_i . Indeed, the number of choices for e_i is exactly the number of edges containing v_s^i but no other element of T_i ; by (H2), the number of edges containing v_s^i and any other element $u \in T_i$ is bounded by $2d^{\varepsilon/5} \cdot d^{1-\varepsilon} \leq 2d^{1-\varepsilon/5}$, so the desired bound follows.

For (W2*), note that given a set $E' \in \binom{\mathcal{H}'}{j'}$ for some $j' \in [j-1]$, which belongs to some set $E \in \mathcal{Z}_j$, the choice of $s \in [t]$ is fixed by E' , as are j' of the edges of E . Hence $\mathbf{1}(\mathcal{Z}_j)(\{E \in \binom{\mathcal{H}'}{j} : E \supseteq E'\}) \leq d^{j-j'}$. The conditions (W3) and (W4) are trivial since there are no conflicts containing edges from $\mathcal{H}' \setminus \mathcal{H}$.

Now we may define, for each³ $j \in [\ell]$ and each j -uniform semi-trackable $w \in \mathcal{Y}$, a new test function $w' : \binom{\mathcal{H}'}{j} \rightarrow [0, 1]$ by $w'(E) := w(E) + \mathbf{1}(\mathcal{Z}_j)(E)$ for $E \subseteq \mathcal{H}$, and $w'(E) := 0$ otherwise, and observe that w' is $(d, \varepsilon/10, \mathcal{C})$ -trackable. Indeed, $w'(\mathcal{H}') = w'(\mathcal{H}) \geq |\mathcal{Z}_j| \geq d^{j+\varepsilon/5}$ so (W1) is satisfied, and (W2) follows from (W2*) for w and $\mathbf{1}(\mathcal{Z}_j)$, using the fact that $(\varepsilon^{-1} + 1)d^{j-j'} \leq d^{-j'-\varepsilon/10}w'(\mathcal{H})$. The remaining conditions (W3) and (W4) for w' follow immediately from the fact that they hold for w and $\mathbf{1}(\mathcal{Z}_j)$. Note that $\mathcal{C}$ is also (d, ℓ, ε) -bounded for $\mathcal{H}'$ by definition. Since also $w'(\mathcal{H}) \leq 3d^{j+\varepsilon/5}$, applying Theorem 4.1 to $\mathcal{H}'$ with each w replaced by the corresponding w' gives us a $\mathcal{C}$ -free matching $\mathcal{M}' \subseteq \mathcal{H}'$ containing no $(\varepsilon/2, \mathcal{C})$ -conflict-sharing pairs such that $w'(\mathcal{M}') = (1 \pm d^{-\varepsilon^3})d^{-j}w'(\mathcal{H}') \leq 4d^{\varepsilon/5}$.

Let $\mathcal{M} := \mathcal{M}' \cap \mathcal{H}$. Then $w(\mathcal{M}) \leq w'(\mathcal{M}') \leq d^{\varepsilon/4}$ as required. Clearly $\mathcal{M}$ is $\mathcal{C}$ -free and contains no conflict-sharing pairs. Since $|V(\mathcal{H}')| \leq (\ell + 1)|V(\mathcal{H})|$, it follows that $\mathcal{M}$ covers all but a $d^{-\varepsilon^3}$ -fraction of $\mathcal{H}$. $\square$

4.2. The Lovász Local Lemma. We now state our other prerequisite, the well-known Lovász Local Lemma, a version of which was originally introduced by Erdős and Lovász [10]. The form we use is an immediate corollary to the general form proved by Spencer [18].

Lemma 4.3. *Let $\mathfrak{A} = \{A_1, \dots, A_n\}$ be a finite set of events in a probability space, and suppose that, for each $i \in [n]$, there exists a set $\mathfrak{B}(i) \subseteq [n]$ such that A_i is mutually independent from $\{A_j : j \in [n] \setminus \mathfrak{B}(i)\}$. Suppose also that for each $i \in [n]$ we have $\mathbb{P}[A_i] < 1/2$ and $\sum_{j \in \mathfrak{B}(i)} \mathbb{P}[A_j] \leq 1/4$. Then $\mathbb{P}[A_1^C \cap \dots \cap A_n^C] > 0$.*

4.3. More general conditions. We prove Theorem 3.1 using slightly weaker conditions than those given in Section 1.1, allowing the case $j_2 = |D \cap \mathcal{H}_2| = 1$, and giving more freedom to high-degree vertices in P . Given a set of edges $E \subseteq \mathcal{H}$, write

$$V_P(E) := \{y \in P : y \in e \text{ for some } e \in E \cap \mathcal{H}_2\}$$

for the set of vertices of P found in edges of $\mathcal{H}_2$ in E . A limitation of conditions (D1)–(D4) is that the number of conflicts is governed uniformly by the minimum degree in $\mathcal{H}_2$. Intuitively, given some conflict $E \in \mathcal{D}$, if the vertices in $V_P(E)$ have much higher degrees in $\mathcal{H}_2$, then it should be easier to avoid E because we have more flexibility when choosing edges of $\mathcal{H}_2$ which are used to cover the vertices of $V_P(E)$ in $\mathcal{M}_2$. As such, we can afford to have more conflicts involving vertices of P with higher degrees in $\mathcal{H}_2$; this motivates the following quantity, which allows us to weight these conflicts accordingly when bounding their number.

We define the *unavoidability* of E to be

$$(4.1) \quad A(E) := \prod_{y \in V_P(E)} d_{\mathcal{H}_2}^{-1}(y),$$

and extend this definition to a conflict hypergraph $\mathcal{G}$ by taking the sum over all conflicts, that is $A(\mathcal{G}) := \sum_{E \in \mathcal{G}} A(E)$. This has a natural interpretation. Suppose we select for each vertex $v \in P$ one edge in $\mathcal{H}_2$ containing v independently and uniformly at random. Then $A(\mathcal{G})$ is the expected number of conflicts in $\mathcal{G}$ of which all edges are selected.

Given $j'_1, j'_2 \in [\ell]$ and sets $C \in \binom{\mathcal{H}_1}{j'_1}$ and $D \in \binom{\mathcal{H}_2}{j'_2}$, write $\mathcal{G}_{[C,D]} := \{E \in \mathcal{G} : C \cup D \subseteq E\}$. Then we may define

$$(4.2) \quad \Delta_{j'_1, j'_2}^A(\mathcal{G}) := \max_{C \in \binom{\mathcal{H}_1}{j'_1}, D \in \binom{\mathcal{H}_2}{j'_2}} A(\mathcal{G}_{[C,D]}),$$

which can be thought of as analogous to the maximum (j'_1, j'_2) -degree $\Delta_{j'_1, j'_2}(\mathcal{G})$ (recalling the definition (3.1)), but with each conflict weighted by its unavoidability. Say that $\mathcal{D}$ is $(d, \ell, \varepsilon, \delta)$ -mixed-bounded if the following hold for all $x, y \in P$ and $j_1 \in [0, \ell], j_2 \in [\ell]$.

- (E1) $|D \cap \mathcal{H}_2| \geq 1$ and $|D| \in [2, \ell]$ for each conflict $D \in \mathcal{D}$;
- (E2) $A(\mathcal{D}_x^{(j_1, j_2)}) \leq d^{j_1 + \delta}$;
- (E3) $\Delta_{j', 0}^A(\mathcal{D}_x^{(j_1, j_2)}) \leq d^{j_1 - j' - \varepsilon}$ for each $j' \in [j_1]$;

³Note that the case $j = 0$ is trivial.

- (E4) $A(\mathcal{D}_{x,y}^{(j_1,j_2)}) \leq d^{j_1-\varepsilon}$ whenever $j_2 \geq 2$;
 (E5) $\Delta_{0,1}(\mathcal{D}_x^{(j_1,1)}) \leq \ell d^{j_1}$;
 (E6) $\Delta_{j',1}(\mathcal{D}_x^{(j_1,1)}) \leq d^{j_1-j'-\varepsilon}$ for each $j' \in [j_1 - 1]$.

Observe that the conditions (E2), (E3), (E4) are analogous to (D2), (D3), (D4), except conflicts are now weighted by their unavoidability; in particular, for a conflict $E \in \mathcal{D}^{(j_1,j_2)}$, it is always the case that $A(E) \leq \delta_P(\mathcal{H}_2)^{-j_2}$, so (d, ℓ, ε) -simply-bounded implies $(d, \ell, \varepsilon, \varepsilon^4)$ -mixed-bounded. Indeed, (D2), (D3), (D4) imply (E2), (E3), (E4), respectively, using the fact that $A(E) \leq |E|\delta_P(\mathcal{H}_2)^{-j_2}$ for every $E \subseteq \mathcal{D}^{(j_1,j_2)}$. If (D1) holds then so does (E1), and in this case both of the conditions (E5) and (E6) are vacuous. Note also that conditions (E5) and (E6) are stronger versions of (E2) and (E3) in the case that $j_2 = 1$, requiring small degrees and codegrees for every individual edge $e \in \mathcal{H}_2$, rather than just for the sum over all such e containing a particular vertex $x \in P$.

In a similar fashion, we may also weaken the degree conditions (H3) and (H4) slightly. Define analogously the *unavoidability of a vertex* $v \in R$ to be

$$A(v) := \sum_{x \in P} \frac{d_{\mathcal{H}_2}(x, v)}{d_{\mathcal{H}_2}(x)}.$$

Then we may require only that for all $x \in P$ and $v \in R$,

- (H3') $A(v) \leq d^{\varepsilon^4}$ (and in particular $\delta_P(\mathcal{H}_2) \geq 1$);
 (H4') $d_{\mathcal{H}_2}(x, v) \leq d^{-\varepsilon} d_{\mathcal{H}_2}(x)$.

Recalling that each edge of $\mathcal{H}_2$ contains exactly one vertex of P , note that by definition $A(v)\delta_P(\mathcal{H}_2) \leq \sum_{x \in P} d_{\mathcal{H}_2}(x, v) = d_{\mathcal{H}_2}(v)$ so (H3) implies (H3'), and clearly (H4) implies (H4'). Observe furthermore that (H3') and (H4') together imply that $\delta_P(\mathcal{H}_2) \geq d^{\varepsilon}$; indeed, by (H3') we must have $d_{\mathcal{H}_2}(x) \geq 1$ for every $x \in P$, so there is some $v \in R$ with $d_{\mathcal{H}_2}(x, v) \geq 1$, and then by (H4') this means $d_{\mathcal{H}_2}(x) \geq d^{\varepsilon}$.

5. PROOF OF THEOREM 3.1

We assume the setting described in Theorem 3.1, and devote the entirety of this section to its proof. However, write $\mathcal{D}'$ in place of $\mathcal{D}$, and assume that it is a $(d, \ell, \varepsilon, \varepsilon^4)$ -mixed-bounded conflict hypergraph. Introduce new constants $\ell', \Gamma, i^* \in \mathbb{N}$ according to the hierarchy $0 < 1/d \ll \varepsilon \ll 1/\ell' \ll 1/\Gamma \ll 1/i^* \ll 1/\ell, 1/k$.

5.1. Proof outline. Recall our two-stage proof method outlined in Section 1.1. We first apply Theorem 4.2 to $\mathcal{H}_1$ to give a conflict-free matching $\mathcal{M}_1 \subseteq \mathcal{H}_1$ covering most of the vertices of P . We then extend this to a covering of P by randomly choosing, for each vertex $x \in P$ which is not already covered by $\mathcal{M}_1$, some edge from $\mathcal{H}_2$ to cover x ; call the resulting set of edges $\mathcal{M}_2 \subseteq \mathcal{H}_2$. Using the Lovász Local Lemma, we show that with non-zero probability $\mathcal{M}_2$ is indeed a matching, and the full matching $\mathcal{M} := \mathcal{M}_1 \cup \mathcal{M}_2$ is conflict-free. This approach presents several challenges which must be overcome.

Firstly, Theorem 4.2 requires $\mathcal{H}_1$ to be essentially vertex-regular, but (H1) provides no lower bound on the degrees of vertices in Q . We solve this in Section 5.2 by adding dummy vertices and edges to increase the vertex degrees in Q . Secondly, it need not be the case that the randomly chosen edges of $\mathcal{M}_2$ form a matching. We solve this in Section 5.3 by simply adding non-disjoint pairs of edges in $\mathcal{H}_2$ as additional conflicts of size 2, replacing $\mathcal{D}'$ by a larger conflict hypergraph $\mathcal{D}$, and thus forcing $\mathcal{M}_2$ to be a matching.

The next challenge is to ensure that we can choose $\mathcal{M}_2$ such that $\mathcal{M}$ is conflict-free. The conflicts we must avoid are of the form $C \cup D$, where two conditions hold: $C \subseteq \mathcal{M}_1$ and no edge in D contains a vertex of P which has already been chosen in $\mathcal{M}_1$. If the second condition does not hold, we say that D is *blocked*. In order to ensure that no such D is contained in $\mathcal{M}_2$, we seek to bound from above the total number of unblocked potential conflicts involving a given vertex $x \in P$ in some edge of $D \subseteq \mathcal{H}_2$; since most vertices of P are already covered by $\mathcal{M}_1$, this significantly reduces the number of conflicts we need to consider. In Section 5.4, we define test functions to track the number of partial conflicts appearing in the matching $\mathcal{M}_1$, as well as the

number of those which are blocked; by taking the difference of these two values we later obtain the desired bound.

The final challenge to overcome is that this approach to blocking conflicts fails when $j_2 = 1$, since the conflicts contain no other edges of $\mathcal{H}_2$ which might be blocked. We therefore require the stricter bound ℓd^{j_1} in (E5) on the number of such conflicts of a given size j_1 containing a given edge $e \in \mathcal{H}_2$. Under the assumption that edges in $\mathcal{H}_1$ appear in the matching $\mathcal{M}_1$ ‘pseudorandomly with probability d^{-1} ’ in some suitable sense, this allows the ‘expected number’ of potential conflicts (of any size) for each e to be up to ℓ^2 ; so choosing e purely at random still will not suffice. To overcome this, we observe that, assuming that the Poisson paradigm applies to our concept of pseudorandomness, there should be a constant proportion (specifically, at least $\exp(-\ell^2)$) of the edges e containing any given vertex x with no potential conflicts arising from $\mathcal{M}_1$. Hence for each vertex $x \in P$ we may restrict our random choice of $\mathcal{M}_2$ to only these *safe* edges. We encode this ‘pseudorandom’ behaviour in a further set of test functions, defined in Section 5.5.

In Section 5.6, we apply Theorem 4.2 to obtain our matching $\mathcal{M}_1$, randomly choose $\mathcal{M}_2$, and use the previously defined test functions to bound the number of potential conflicts in each case, so that $\mathcal{M}_2$ is conflict-free with non-zero probability.

5.2. Regularising $\mathcal{H}_1$. First, in order to apply Theorem 4.2 to the hypergraph $\mathcal{H}_1$, we need to ensure that all vertices in $\mathcal{H}_1$ have degree roughly d , rather than just those in P ; since we only require a P -perfect matching, and make no statement about which vertices of Q are covered, we can achieve this by simply adding dummy edges to boost the degrees of vertices in Q .

Assume that $q > 0$ (otherwise we may skip this step), and let $m := |Q|$. For each vertex $v \in Q$, let $d' := d - d_{\mathcal{H}_1}(v)$ and add d' new edges, each containing v and a set of $k - 1$ new vertices, such that each new vertex is only ever contained in one edge. Refer to the set of new vertices as Q' and the set of new edges as $\mathcal{E}_Q$. Add further new vertices until Q' contains $|Q|dm$ vertices. Next, let $\mathcal{F}'$ be a binomial random k -graph with vertex set Q' and expected degree $(1 - d^{-\varepsilon}/2)d$. It is routine to show that, with high probability, $(1 - d^{-\varepsilon})d \leq d_{\mathcal{F}'}(v) \leq d - 1$ and $d_{\mathcal{F}'}(u, v) \leq d^{1-\varepsilon}$ for all distinct $u, v \in Q'$; take $\mathcal{F}$ to be a graph satisfying these properties. Note that each $v \in Q'$ belongs to at most one edge in $\mathcal{E}_Q$, so in total $d_{\mathcal{H}}(v) \leq d$. Define the hypergraph $\mathcal{H}'_1 := \mathcal{H}_1 \cup \mathcal{E}_Q \cup \mathcal{F}$, which we will use in place of $\mathcal{H}_1$ later in the proof.

5.3. Ensuring a Matching in $\mathcal{H}_2$. Let $\mathcal{E} := \{\{e, f\} \subseteq \mathcal{H}_2 : \emptyset \neq e \cap f \subseteq R\}$ be the conflict hypergraph containing as conflicts all pairs of edges in $\mathcal{H}_2$ that overlap (only) in R . We show that $\mathcal{E}$ is a $(d, \ell, \varepsilon/2, 2\varepsilon^4)$ -mixed bounded hypergraph by verifying that it satisfies (E1)-(E6). Note that in particular this implies that $\mathcal{D} := \mathcal{D}' \cup \mathcal{E}$ is $(d, \ell, \varepsilon/3, 3\varepsilon^4)$ -mixed-bounded; indeed, (E1) is trivial, the bounds in conditions (E2), (E3), (E4) are additive and the bounds in (E5) and (E6) are unaffected by $\mathcal{E}$.

Indeed, the condition (E1) is clear and (E3), (E5), (E6) are trivial. For (E2), note that we need only consider the case $(j_1, j_2) = (0, 2)$ and suppose that $D = \{e, f\}$ is a conflict containing $x \in P$; say $x \in e$, and let $v \in e \cap f \cap R$ and $y \in f \cap P$, noting that $x \neq y$. The number of such conflicts D , given x, v, y , is exactly $d_{\mathcal{H}_2}(x, v)d_{\mathcal{H}_2}(y, v)$. Moreover, recalling definitions (4.1) and (4.2), as $|e \cap P| = |f \cap P| = 1$, we have $A(D) = d_{\mathcal{H}_2}(x)^{-1}d_{\mathcal{H}_2}(y)^{-1}$, so we obtain

$$A(\mathcal{E}_x) = \sum_{D \in \mathcal{E}_x} A(D) \leq \sum_{v \in R} \sum_{y \in P} \frac{d_{\mathcal{H}_2}(x, v)d_{\mathcal{H}_2}(y, v)}{d_{\mathcal{H}_2}(x)d_{\mathcal{H}_2}(y)} = \sum_{v \in R} \frac{d_{\mathcal{H}_2}(x, v)}{d_{\mathcal{H}_2}(x)} A(v) \leq r \cdot d^{\varepsilon^4},$$

which suffices, using the fact that $A(v) \leq d^{\varepsilon^4}$ by (H3'), and noting that we allow the case $y = x$ in the sum, as we seek only an upper bound. For (E4), observe that similarly

$$A(\mathcal{E}_{x,y}) \leq \sum_{v \in R} \frac{d_{\mathcal{H}_2}(x, v)}{d_{\mathcal{H}_2}(x)} \frac{d_{\mathcal{H}_2}(y, v)}{d_{\mathcal{H}_2}(y)} \stackrel{(H4')}{\leq} d^{-\varepsilon} \sum_{v \in R} \frac{d_{\mathcal{H}_2}(x, v)}{d_{\mathcal{H}_2}(x)} = rd^{-\varepsilon},$$

which suffices. The rest of the proof will show that all of the conflicts in $\mathcal{D}$ can indeed be avoided when choosing $\mathcal{M}_2$.

5.4. Tracking mixed conflicts: the case $j_2 \geq 2$. In this section, we define a set of test functions and show that they are either trackable or semi-trackable. These will be useful in Section 5.6.4 for showing that the expected number of unblocked conflicts can be appropriately bounded from above such that the Lovász Local Lemma can be used to avoid them.

For the remainder of this section, we fix $j_1 \in [0, \ell]$ and $j_2 \in [2, \ell]$, and consider only conflicts with j_1 and j_2 edges from $\mathcal{H}_1$ and $\mathcal{H}_2$ respectively; for ease of notation, let $\mathcal{G} = \mathcal{D}^{(j_1, j_2)}$ (recalling the notation from Section 3.3). For $i \in [2]$, we write $E_i(\mathcal{G}) := \{E \cap \mathcal{H}_i : E \in \mathcal{G}\}$ for the set of $\mathcal{H}_i$ -parts of conflicts in $\mathcal{G}$.

5.4.1. Defining test functions. We extend our previous notation from Section 4.3 by writing $\mathcal{G}_{[C]} = \mathcal{G}_{[C, \emptyset]}$, $\mathcal{G}_{[C], x} = (\mathcal{G}_{[C]})_x$ and $\mathcal{G}_{[C], x, y} = (\mathcal{G}_{[C]})_{x, y}$, for any $C \subseteq \mathcal{H}_1$ and $x, y \in P$. For the remainder of this section, fix a vertex $x \in P$. We first define $w_x: \binom{\mathcal{H}_1}{j_1} \rightarrow \mathbb{R}_{\geq 0}$ by

$$w_x(C) := A(\mathcal{G}_{[C], x}),$$

recalling (4.1). Given a matching $\mathcal{M}_1 \subseteq \mathcal{H}_1$ containing C (but not covering x), if we choose $\mathcal{M}_2 \subseteq \mathcal{H}_2$ randomly, then the function $w_x(C)$ represents the expected number of conflicts from $\mathcal{G}$ present in $\mathcal{M} = \mathcal{M}_1 \cup \mathcal{M}_2$ which have C as their $\mathcal{H}_1$ -part and contain x in their $\mathcal{H}_2$ -part. As such, $w_x(\mathcal{M}_1)$ represents the expected number of conflicts containing x in their $\mathcal{H}_2$ -part for which the $\mathcal{H}_1$ -part is covered by $\mathcal{M}_1$.

Next, in order to ensure that the test functions we define are trackable, we make an intermediate definition. Say that a set of edges $E \subseteq \mathcal{H}$ is *testable* if it satisfies the following conditions:

- $E \cap \mathcal{H}_1$ is a matching;
- E is $\mathcal{C}$ -free;
- E contains no $\varepsilon/2$ -conflict-sharing pairs of edges (as defined in Section 4.1).

Say that E is *untestable* if it is not testable.⁴ We may (and do henceforth) assume without loss of generality that all of the conflicts in $\mathcal{D}$ are testable. Indeed, letting $\overline{\mathcal{D}} \subseteq \mathcal{D}$ be the set of all testable conflicts, it is clear that $\overline{\mathcal{D}}$ is also $(d, \ell, \varepsilon/3, 3\varepsilon^4)$ -mixed-bounded. Furthermore, if we are able to find a matching $\mathcal{M}_2 \subseteq \mathcal{H}_2$ such that $\mathcal{M} = \mathcal{M}_1 \cup \mathcal{M}_2$ is $\overline{\mathcal{D}}$ -free, with $\mathcal{M}_1 \subseteq \mathcal{H}_1$ obtained from Theorem 4.2, then $\mathcal{M}_1$ may not contain any untestable set, which in particular means that $\mathcal{M}$ is also $\mathcal{D}$ -free. Similarly we may assume that $w_x(C) = 0$ for any C containing x , because if C is covered by $\mathcal{M}_1$ then no edge containing x can be chosen in $\mathcal{M}_2$.

We may now define further $w'_x: \binom{\mathcal{H}_1}{j_1+1} \rightarrow \mathbb{R}_{\geq 0}$ by

$$w'_x(C') := \mathbb{1}(C' \text{ testable}) \sum_{e \in C'} \sum_{y \in (e \cap P) \setminus \{x\}} A(\mathcal{G}_{[C' \setminus \{e\}], x, y}).$$

Observe that the function $w'_x(C')$ represents the sum, over each possible partition $C' = C \cup \{e\}$, of the expected number of those conflicts counted by $w_x(C)$ whose $\mathcal{H}_2$ -part shares some vertex $y \in P \setminus \{x\}$ with the edge $e \in \mathcal{H}_1$. In particular this means $w'_x(\mathcal{M}_1)$ represents the expected number of those conflicts counted by $w_x(\mathcal{M}_1)$ whose $\mathcal{H}_2$ -part shares some vertex $y \in P$ with some edge $e \in \mathcal{M}_1$. In our proof, we will consider the quantity $w_x(\mathcal{M}_1) - w'_x(\mathcal{M}_1)$, which counts the number of conflicts (containing x in their $\mathcal{H}_2$ -part) for which the $\mathcal{H}_1$ -part is contained in $\mathcal{M}_1$, and the $\mathcal{H}_2$ -part is at risk of being contained in $\mathcal{M}_2$. By bounding this quantity from above, we can ensure that all such conflicts can be avoided when choosing $\mathcal{M}_2$. Observe also that both function definitions still make sense when $j_1 = 0$, in which case w_x is defined only on the empty set, and w'_x is defined on single edges.

5.4.2. Ignoring untestable sets. The goal of this section is to prove (5.1) below. For any given $C \in E_1(\mathcal{G})$, and any vertex $y \in P$ not contained in an edge of C , we write $b_{C, y} := |\{e \in \mathcal{H}_1 : y \in e \text{ and } C \cup e \text{ is testable}\}|$, and claim that

$$(5.1) \quad (1 - d^{-\varepsilon/3})d \leq b_{C, y} \leq d.$$

⁴This name reflects the fact that we must ignore such sets when defining our test functions, in order to ensure that the functions are trackable (as per the definition in Section 4.1).

In other words, restricting to testable sets only removes a negligible proportion of the total number of edges e containing y for which $w'_x(C \cup \{e\})$ would otherwise be positive. Indeed, we estimate $b_{C,y}$ by bounding from above the number of $e \in \mathcal{H}_1$ such that $y \in e$ but $C \cup e$ is untestable. **Firstly**, for an $\mathcal{H}_1$ -part C of size j_1 , observe that C covers in total kj_1 vertices z , and for each of these there are at most $\Delta_2(\mathcal{H}_1)$ edges $e \in \mathcal{H}_1$ with $y, z \in e$. Thus the total number of $e \in \mathcal{H}_1$ with $y \in e$ for which $C \cup e$ is not a matching is at most $j_1 k d^{1-\varepsilon}$ by (H2). **Secondly**, by (C3), the number of such e for which $C \cup e$ contains a conflict from $\mathcal{C}$ is at most

$$\sum_{j=2}^{\ell} \sum_{F \in \binom{C}{j-1}} \Delta_{j-1}(\mathcal{C}^{(j)}) \leq \ell 2^{j_1} d^{1-\varepsilon},$$

where F represents a subset of C for which $F \cup e$ is a conflict of size j .⁵ **Thirdly**, we bound the number of e for which $C \cup e$ contains a conflict-sharing pair.⁶ For each edge $f \in \mathcal{H}_1$ and $j \in [\ell]$, let $P_f^j := \{e \in \mathcal{H}_1 : |(\mathcal{C}_e)^{(j)} \cap (\mathcal{C}_f)^{(j)}| \geq d^{j-\varepsilon/2}\}$. Observe that by definition we may rewrite

$$\sum_{e \in P_f^j} |(\mathcal{C}_e)^{(j)} \cap (\mathcal{C}_f)^{(j)}| = \sum_{C' \in (\mathcal{C}_f)^{(j)}} \sum_{e \in P_f^j} \mathbf{1}(C' \cup e \in \mathcal{C}^{(j+1)}) \leq \sum_{C' \in (\mathcal{C}_f)^{(j)}} |\{e \in \mathcal{H}_1 : C' \cup e \in \mathcal{C}^{(j+1)}\}|,$$

and applying conditions (C2) and (C3), we obtain that

$$\sum_{C' \in (\mathcal{C}_f)^{(j)}} |\{e \in \mathcal{H}_1 : C' \cup e \in \mathcal{C}^{(j+1)}\}| \leq \Delta(\mathcal{C}^{(j)}) \Delta_j(\mathcal{C}^{(j+1)}) \leq d^{j+1-\varepsilon},$$

where the second expression is the product of the number of choices for C' given f and the number of choices for e given C' . Then, by the definition of P_f^j , we see that

$$(5.2) \quad |P_f^j| \leq d^{\varepsilon/2-j} \sum_{e \in P_f^j} |(\mathcal{C}_e)^{(j)} \cap (\mathcal{C}_f)^{(j)}| \leq d^{1-\varepsilon/2}.$$

Hence, summing over each $f \in C$ and $j \in [\ell]$, this means that in total there are at most $j_1 \ell d^{1-\varepsilon/2}$ edges $e \in \mathcal{H}_1$ for which $C \cup \{e\}$ contains a conflict-sharing pair. We finish by subtracting the three bounds we have obtained from (H1) to see that (5.1) holds.

5.4.3. Estimating values of test functions. In this section we use (5.1) to show that ignoring untestable sets in the definition of w'_x does not significantly affect the value of $w'_x(\mathcal{H}_1)$, compared to what its value would be if they were included. Specifically, we evaluate $w_x(\mathcal{H}_1)$ and $w'_x(\mathcal{H}_1)$ and show that $w'_x(\mathcal{H}_1) \approx (j_2 - 1)dw_x(\mathcal{H}_1)$. Recalling the definition of $V_P(\cdot)$ from Section 4.3, begin by rewriting

$$\begin{aligned} w'_x(\mathcal{H}_1) &= \sum_{C \in \binom{\mathcal{H}_1}{j_1}} \sum_{e \in \mathcal{H}_1 \setminus C} \mathbf{1}(C \cup e \text{ testable}) \sum_{y \in (e \cap P) \setminus \{x\}} A(\mathcal{G}_{[C],x,y}) \\ &= \sum_{C \in \binom{\mathcal{H}_1}{j_1}} \sum_{e \in \mathcal{H}_1 \setminus C} \mathbf{1}(C \cup e \text{ testable}) \sum_{y \in (e \cap P) \setminus \{x\}} \sum_{D \in E_2(\mathcal{G}_{[C],x})} \mathbf{1}(y \in V_P(D)) A(D) \\ &= \sum_{C \in \binom{\mathcal{H}_1}{j_1}} \sum_{D \in E_2(\mathcal{G}_{[C],x})} A(D) \sum_{y \in V_P(D) \setminus \{x\}} |\{e \in \mathcal{H}_1 \setminus C : y \in e \text{ and } C \cup e \text{ testable}\}| \\ &= \sum_{E \in \mathcal{G}_x} A(E) \sum_{y \in V_P(E) \setminus \{x\}} b_{E \cap \mathcal{H}_1, y}. \end{aligned}$$

Note here that $E = C \cup D$ and recall that $\mathcal{G} := \mathcal{D}^{(j_1, j_2)}$, so there is an obvious bijection between the set of pairs C, D in the sum and the set $\mathcal{G}_x$. Now observe that by definition

⁵Recall here that C is testable by our assumption on $\mathcal{D}$, so this is the only way in which $C \cup \{e\}$ may contain a conflict.

⁶Note that e must be one element of this pair, by testability of C .

$w_x(\mathcal{H}_1) = A(\mathcal{G}_x) = \sum_{E \in \mathcal{G}_x} A(E)$. Therefore, recalling that $|V_P(E) \setminus \{x\}| = j_2 - 1$ and applying each of the two bounds in (5.1) to the inner sum above, we obtain that

$$(5.3) \quad (1 - d^{-\varepsilon/3})(j_2 - 1)dw_x(\mathcal{H}_1) \leq w'_x(\mathcal{H}_1) \leq (j_2 - 1)dw_x(\mathcal{H}_1).$$

We use these estimates in Section 5.6.4 to ensure that the choice of $\mathcal{M}_1$ leaves very few unblocked potential conflicts, by showing that $(j_2 - 1)w_x(\mathcal{M}_1) \approx w'_x(\mathcal{M}_1)$, so that most potential conflicts are in fact blocked; this will give an upper bound on the expected number of unblocked conflicts containing x which are present in $\mathcal{M}_1$.

5.4.4. Checking test function conditions. In order to use Theorem 4.2 to track the values of w_x and w'_x , we must first scale them appropriately so that they take values in the interval $[0, \ell']$. In order to do this, define $\alpha_x := \max\{\alpha'_x, \alpha''_x\}$, where

$$\alpha'_x := \max_{j' \in [j_1]} d^{j' - j_1} \Delta_{j', 0}^A(\mathcal{G}_x), \text{ and } \alpha''_x := d^{-j_1} \max_{y \in P} A(\mathcal{G}_{x, y}),$$

recalling the definitions in (4.1) and (4.2). Note that $\alpha_x \leq d^{-\varepsilon/3}$ by (E3) and (E4). We show now that either both of the functions $\alpha_x^{-1}w_x$ and $\alpha_x^{-1}w'_x$ are $(d, \varepsilon/10, \mathcal{C})$ -trackable, or the function $\alpha_x^{-1}w_x$ is $(d, \varepsilon/10, \mathcal{C})$ -semi-trackable. Observe that we need not consider the case $\alpha_x = 0$, since in this case $w_x = w'_x = 0$ are trivially trackable.

Firstly, the case $j' = j_1$ in α'_x ensures that $\alpha_x^{-1}w_x(C) \leq 1 \leq \ell'$ for all $C \in \binom{\mathcal{H}_1}{j_1}$. Also $\alpha_x^{-1}w'_x(C') \leq \alpha_x^{-1}(j_1 + 1)p\Delta_{j_1, 0}^A(\mathcal{G}_x) \leq (j_1 + 1)p \leq \ell'$ by the definition of α'_x , recalling that $\ell' \ll 1/\ell, 1/k$. Clearly both $\alpha_x^{-1}w_x$ and $\alpha_x^{-1}w'_x$ are zero on any set of edges which is not a matching, recalling the assumption that all conflicts in $\mathcal{G}$ are testable, so they are indeed both ℓ' -test functions (which are j_1 -uniform and $(j_1 + 1)$ -uniform, respectively).

Observe that conditions (W3) and (W4) for $\alpha_x^{-1}w_x$ are immediate from the assumption that no element of $\mathcal{G}$ contains a conflict from $\mathcal{C}$ or conflict-sharing pair, and they both hold for $\alpha_x^{-1}w'_x$ by definition. We prove that both functions satisfy (W2*). Given $j' \in [0, \ell]$ and $E \in \binom{\mathcal{H}_1}{j'}$, write $\mathcal{Z}_E^{(j)} := \{Z \in \binom{\mathcal{H}_1}{j} : Z \supset E\}$ for each $j \in [j', \ell]$. Then for $j' \in [j_1 - 1]$, we have $\alpha_x^{-1}w_x(\mathcal{Z}_E^{(j_1)}) = \alpha_x^{-1}A(\mathcal{G}_{[E], x}) \leq d^{j_1 - j'}$ by the definition of α'_x , so (W2*) holds for w_x .

For $\alpha_x^{-1}w'_x$, let $j' \in [j_1 + 1 - 1] = [j_1]$, consider a sum over $C' = C \cup \{e\}$ with $E \subseteq C'$, and split into two cases depending on whether $E \subseteq C$. Specifically, write

$$(5.4) \quad w'_x(\mathcal{Z}_E^{(j_1+1)}) \leq \sum_{C' \in \mathcal{Z}_E^{(j_1+1)}} \sum_{e \in C'} \sum_{y \in (e \cap P) \setminus \{x\}} A(\mathcal{G}_{[C' \setminus \{e\}], x, y}) \leq S_1 + S_2,$$

where

$$S_1 := \sum_{C \in \mathcal{Z}_E^{(j_1)}} \sum_{e \in \mathcal{H}_1} \sum_{y \in (e \cap P) \setminus \{x\}} A(\mathcal{G}_{[C], x, y}),$$

and

$$S_2 := \sum_{e \in E} \sum_{C \in \mathcal{Z}_{E \setminus \{e\}}^{(j_1)}} \sum_{y \in (e \cap P) \setminus \{x\}} A(\mathcal{G}_{[C], x, y}).$$

Firstly for the case $E \subseteq C$, we may rearrange sums to see that

$$\begin{aligned} S_1 &= \sum_{C \in \mathcal{Z}_E^{(j_1)}} \sum_{e \in \mathcal{H}_1} \sum_{y \in P \setminus \{x\}} \sum_{D \in E_2(\mathcal{G}_{[C], x})} \mathbf{1}(y \in V_P(D), y \in e) A(D) \\ &= \sum_{C \in \mathcal{Z}_E^{(j_1)}} \sum_{D \in E_2(\mathcal{G}_{[C], x})} A(D) \sum_{y \in V_P(D) \setminus \{x\}} d_{\mathcal{H}_1}(y). \end{aligned}$$

We can bound this from above by

$$(5.5) \quad S_1 \leq (j_2 - 1)d \sum_{C \in \mathcal{Z}_E^{(j_1)}} \sum_{D \in E_2(\mathcal{G}_{[C], x})} A(D) = (j_2 - 1)dw_x(\mathcal{Z}_E^{(j_1)}) \leq (j_2 - 1)\alpha_x d^{j_1 - j' + 1},$$

using the fact that $d_{\mathcal{H}_1}(y) \leq d$ and $|V_P(D) \setminus \{x\}| = j_2 - 1$, and then the definition of w_x and the condition (W2*) for w_x .

For the case that $e \in E$, we in fact split into two subcases, according to whether $j' > 1$. If $j' > 1$, then we can rearrange and use (W2*) for w_x similarly to see that

$$(5.6) \quad S_2 \leq \sum_{e \in E} \sum_{C \in \mathcal{Z}_{E \setminus \{e\}}^{(j_1)}} pA(\mathcal{G}_{[C],x}) = p \sum_{e \in E} w_x(\mathcal{Z}_{E \setminus \{e\}}^{(j_1)}) \leq j' p \alpha_x d^{j_1-j'+1}.$$

If however $j' = 1$, then $E = \{e\}$ and we are summing over all possible $C \in \binom{H_1}{j_1}$, which is why we need the definition of α_x'' to give

$$(5.7) \quad S_2 = \sum_{y \in (e \cap P) \setminus \{x\}} A(\mathcal{G}_{x,y}) \leq p \alpha_x d^{j_1-j'+1}.$$

Plugging the bounds (5.5), (5.6), and (5.7) back into (5.4) gives $\alpha_x^{-1} w'_x(\mathcal{Z}_E^{(j_1+1)}) \leq C_{\ell,k} d^{j_1+1-j'}$ for some constant $C_{\ell,k}$ depending only on ℓ and k , as required for (W2*).

Hence if $\alpha_x^{-1} w_x(\mathcal{H}_1) \leq d^{j_1+\varepsilon/5}$, then $\alpha_x^{-1} w_x$ is $(d, \varepsilon/10, \mathcal{C})$ -semi-trackable. If instead $\alpha_x^{-1} w_x(\mathcal{H}_1) \geq d^{j_1+\varepsilon/5}$, then by the estimate (5.3) we have $\alpha_x^{-1} w'_x(\mathcal{H}_1) \geq d^{j_1+1+\varepsilon/10}$ so (W1) holds for both functions. In this case, the condition (W2*) implies (W2), so they are both $(d, \varepsilon/10, \mathcal{C})$ -trackable.

5.5. Tracking mixed conflicts: the case $j_2 = 1$. In this section, we suppose $j_2 = 1$, noting that $j_1 \geq 1$ by (E1), and define a further set of test functions; these are used in Section 5.6.2 to show that such conflicts can be avoided. We fix $x \in P$ for the remainder of this section, and for ease of notation we write $N_x := \{e \in \mathcal{H}_2 : x \in e\}$ and $d_x := d_{\mathcal{H}_2}(x) = |N_x|$. Furthermore, in this section, we define $\mathcal{G} := \cup_{j_1 \in [\ell]} \mathcal{D}^{(j_1,1)}$, noting that we now consider only $j_2 = 1$, but for all values of j_1 together.

5.5.1. Defining test functions. Recall that we fixed a constant $i^* \in \mathbb{N}$ with $1/i^* \ll 1/\ell$. Let $m \in [i^*]$, $s' \in [m\ell]$, $s \in [s']$, and consider any collection $\bar{C} := \{C_t\}_{t \in [m]}$ of m edge sets $C_t \subseteq \mathcal{H}_1$, each with size $|C_t| \in [\ell]$, such that $\sum_{t \in [m]} |C_t| = s'$ but $|\bigcup_{t \in [m]} C_t| = s$. Let $\mathcal{P}(\bar{C}) := (|\bigcap_{t \in T} C_t|)_{T \subseteq [m]}$ encode the sizes of the C_t and their intersections, and let $\mathcal{P}_{m,s,s'}^*$ be the set of all possible values of $\mathcal{P}(\bar{C})$. We consider $\bar{C}$ to be unordered, and identify elements in $\mathcal{P}_{m,s,s'}^*$ which arise from different orderings of the same $\bar{C}$, thus ensuring that $\mathcal{P}(\bar{C})$ is still well-defined. Given $\mathcal{P} \in \mathcal{P}_{m,s,s'}^*$ and a set $E \subseteq \mathcal{H}_1$ of size $|E| = s$, write $C(E, \mathcal{P})$ for the set of all such collections $\bar{C}$ with $\bigcup_{t \in [m]} C_t = E$ and $\mathcal{P}(\bar{C}) = \mathcal{P}$. We may now define a function $\bar{w}_x^{m,s,s'} : \binom{\mathcal{H}_1}{s} \rightarrow \mathbb{R}_{\geq 0}$ by

$$\bar{w}_x^{m,s,s'}(E) := d_x^{-1} \sum_{\mathcal{P} \in \mathcal{P}_{m,s,s'}^*} \sum_{\bar{C} \in C(E, \mathcal{P})} \sum_{e \in N_x} \prod_{t=1}^m \mathbb{1}(C_t \cup e \in \mathcal{G}).$$

Given a matching $\mathcal{M}_1 \subseteq \mathcal{H}_1$ containing E (but not covering x), if we choose $e \in N_x \subseteq \mathcal{H}_2$ randomly, then the function $\bar{w}_x^{m,s,s'}(E)$ represents the expected number of collections $\bar{C} \in \bigcup_{\mathcal{P} \in \mathcal{P}_{m,s,s'}^*} C(E, \mathcal{P})$ for which $C_t \cup e$ is a conflict from $\mathcal{G}$ for every $t \in [m]$. As such $\bar{w}_x^{m,s,s'}(\mathcal{M}_1)$ represents the expected number of m -sets of $\mathcal{H}_1$ -parts $C_1, \dots, C_m$ present in $\mathcal{M}_1$ such that $\sum_{t \in [m]} |C_t| = s'$ but $|\bigcup_{t \in [m]} C_t| = s$ and each part individually forms a conflict with the randomly chosen $\mathcal{H}_2$ -edge $e \in N_x$. We are interested in the value of $\bar{w}_x^{m,s,s'}(\mathcal{M}_1)$, but the function $\bar{w}_x^{m,s,s'}$ itself may not technically be trackable, so we define further

$$w_x^{m,s,s'}(E) := \mathbb{1}(E \text{ is testable}) \bar{w}_x^{m,s,s'}(E),$$

which approximates $\bar{w}_x^{m,s,s'}$ and is trackable, as we will show.

5.5.2. Checking trackability. We again define a normalising constant β_x , analogously to α_x , by $\beta_x := \max\{\beta'_x, \beta''_x\}$, where

$$\beta'_x := \max_{j_1 \in [\ell]} \max_{j' \in [j_1-1]} d^{j'-j_1} \Delta_{j',1}(\mathcal{G}_x^{(j_1,1)}), \text{ and } \beta''_x := d_x^{-1} \max_{j_1 \in [\ell]} \Delta_{j_1,0}(\mathcal{G}_x^{(j_1,1)}).$$

Note again that $\beta'_x \leq d^{-\varepsilon/3}$ by (E6). Likewise, observe that $A(E) = d_x^{-1}$ for any $E \in \mathcal{G}_x^{(j_1,1)}$, so $\Delta_{j_1,0}^A(\mathcal{G}_x^{(j_1,1)}) = d_x^{-1} \Delta_{j_1,0}(\mathcal{G}_x^{(j_1,1)})$, from which it follows that $\beta''_x \leq d^{-\varepsilon/3}$ by the $j' = j_1$ case of (E3). We now fix m, s, s' and aim to prove that the function $\beta_x^{-1} w_x^{m,s,s'}$ is either $(d, \varepsilon/10, \mathcal{C})$ -trackable or $(d, \varepsilon/10, \mathcal{C})$ -semi-trackable. By the definition of β''_x , we have that

$$\bar{w}_x^{m,s,s'}(E) \leq d_x^{-1} \sum_{\mathcal{P} \in \mathcal{P}_{m,s,s'}^*} \sum_{\bar{C} \in C(E, \mathcal{P})} d_{\mathcal{G}_x^{(C_1,1)}}(C_1) \leq d_x^{-1} \Gamma^2 d_x \beta''_x \leq \beta_x \ell',$$

recalling that $1/\ell' \ll 1/\Gamma \ll 1/i^*$ and thus noting that $|\mathcal{P}_{m,s,s'}^*| + |C(E, \mathcal{P})| \leq \Gamma$. We hence see that $\beta_x^{-1} w_x^{m,s,s'} \leq \beta_x^{-1} \bar{w}_x^{m,s,s'}$ is an ℓ' -test function, and observe also that $\beta_x^{-1} w_x^{m,s,s'}$ satisfies (W3) and (W4) by the definition of testability.

Next, we verify that $\beta_x^{-1} \bar{w}_x^{m,s,s'}$ satisfies (W2*). To this end, let $F \subseteq \mathcal{H}_1$ be an edge set of size $j' \in [s-1]$, and recall that $\mathcal{Z}_F^{(s)} = \{Z \in \binom{\mathcal{H}_1}{s} : Z \supset F\}$. Given $\mathcal{P} \in \mathcal{P}_{m,s,s'}^*$, let $\mathcal{F}(F, \mathcal{P})$ be the set of all collections $\bar{F} := \{F_t\}_{t \in [m]}$ of m subsets $F_t \subseteq F$ such that $\bigcup_{t \in [m]} F_t = F$ and there exists some collection $\bar{C} \subseteq 2^{\mathcal{H}_1}$ with $\mathcal{P}(\bar{C}) = \mathcal{P}$ and $C_t \cap F = F_t$ for every $t \in [m]$. Note that any collection $\bar{C}$ with $F \subseteq \bigcup \bar{C}$ uniquely determines some such $\bar{F} \in \mathcal{F}(F, \mathcal{P}(\bar{C}))$. Given $\bar{F} \in \mathcal{F}(F, \mathcal{P})$ and $e \in N_x$, we aim to bound $q(\bar{F}, \mathcal{P}) := \sum_{e \in N_x} q(\bar{F}, \mathcal{P}, e)$, where we define $q(\bar{F}, \mathcal{P}, e)$ as the number of collections $\bar{C} = \{C_t\}_{t \in [m]}$ with $\mathcal{P}(\bar{C}) = \mathcal{P}$ for which $F_t = C_t \cap F$ and $C_t \cup e \in \mathcal{G}$ for every $t \in [m]$.

Suppose we have already chosen $C_1, \dots, C_{t-1}$ such that their intersections are compatible with $\mathcal{P}$, and for every $t' \in [t-1]$ we have $F_{t'} \subseteq C_{t'}$ and $C_{t'} \cup e \in \mathcal{G}$. Write $C'_t := C_1 \cup \dots \cup C_{t-1} \cup F$ and let a_t and a'_t denote the desired sizes of $C_t \cap C'_t$ and $C_t \setminus C'_t$, respectively, noting that these are uniquely determined by $\mathcal{P}$ and $\bar{F}$. If $a_t, a'_t > 0$, then to choose C_t given e , we have at most 2^s choices for $C_t \cap C'_t$, each yielding at most $\Delta_{a_t,1}(\mathcal{G}^{(a_t+a'_t,1)})$ choices for $C_t \setminus C'_t$, so by the definition of β'_x we obtain at most $2^s \beta'_x d^{a'_t}$ total choices for C_t . If $a_t = 0$, then the number of choices for C_t given e is at most $\ell d^{a'_t}$ by (E5). If $a'_t = 0$, then the number of choices for C_t is at most 2^s , and this is compatible with most $\Delta_{a_t,0}(\mathcal{G}^{(a_t,1)}) \leq \beta''_x d_x$ choices for e , by the definition of β''_x . Note that by definition it is always the case that $\sum_{t \in [m]} a'_t = s - j'$, and without loss of generality we may assume that $a_1 \neq 0$, so that the number of choices for (C_1, e) is at most $2^s \beta_x d^{a'_1} d_x$ in all cases. As such we obtain that $q(\bar{F}, \mathcal{P}) \leq \Gamma \beta_x d^{s-j'} d_x$, recalling that $1/\Gamma \ll 1/i^*$. Observe also that $|\mathcal{P}_{m,s,s'}^*| + |\mathcal{F}(F, \mathcal{P})| \leq \Gamma$. We may thus bound

$$\bar{w}_x^{m,s,s'}(\mathcal{Z}_F^{(s)}) = d_x^{-1} \sum_{\mathcal{P} \in \mathcal{P}_{m,s,s'}^*} \sum_{\bar{F} \in \mathcal{F}(F, \mathcal{P})} q(\bar{F}, \mathcal{P}) \leq \Gamma^3 \beta_x d^{s-j'}.$$

Therefore $\beta_x^{-1} w_x^{m,s,s'}(\mathcal{Z}_F^{(s)}) \leq \beta_x^{-1} \bar{w}_x^{m,s,s'}(\mathcal{Z}_F^{(s)}) \leq \varepsilon^{-1} d^{s-j'}$, which implies that $\beta_x^{-1} w_x^{m,s,s'}$ satisfies (W2*). Note that it is always the case that at least one of (W1) and (W1*) holds, so $\beta_x^{-1} w_x^{m,s,s'}$ is either $(d, \varepsilon/10, \mathcal{C})$ -trackable or $(d, \varepsilon/10, \mathcal{C})$ -semi-trackable.

5.5.3. Ignoring untestable sets. In this subsection, we again fix m, s, s' , and claim that

$$(5.8) \quad \text{if } \bar{w}_x^{m,s,s'}(\mathcal{H}_1) \geq d^{s-\varepsilon^3}, \quad \text{then } (1 - d^{-\varepsilon/5}) \bar{w}_x^{m,s,s'}(\mathcal{H}_1) \leq w_x^{m,s,s'}(\mathcal{H}_1) \leq \bar{w}_x^{m,s,s'}(\mathcal{H}_1).$$

Note that in particular, in this case, we have that $\beta_x^{-1} w_x^{m,s,s'}$ satisfies (W1), so is trackable.

Indeed, the upper bound is trivial, so we proceed to prove the lower bound. Start by writing $w_x^{m,s,s'}(E) \geq \bar{w}_x^{m,s,s'}(E) - f(E) - g(E) - h(E)$ where

$$\begin{aligned} f(E) &:= \mathbb{1}(E \text{ not a matching}) \bar{w}_x^{m,s,s'}(E), \\ g(E) &:= \mathbb{1}(E \text{ contains a conflict in } \mathcal{C}) \bar{w}_x^{m,s,s'}(E), \\ h(E) &:= \mathbb{1}(E \text{ contains a conflict-sharing pair}) \bar{w}_x^{m,s,s'}(E). \end{aligned}$$

We first bound $f(\mathcal{H}_1)$ from above by considering the number of choices of $\bar{C} = \{C_t\}_{t \in [m]}$ for which $E = \bigcup_{t \in [m]} C_t$ is not a matching. Given $\mathcal{P} \in \mathcal{P}_{m,s,s'}^*$ and $e \in N_x$, we aim to bound

$q(\mathcal{P}, e)$, which we define as the number of collections $\overline{C} = \{C_t\}_{t \in [m]}$ for which $C_t \cup e \in \mathcal{G}$ for every $t \in [m]$, $\mathcal{P}(\overline{C}) = \mathcal{P}$, and there exist $e_1 \in C_1, e_2 \in C_2$ with some vertex $v \in e_1 \cap e_2$; note that this case suffices without loss of generality, since we regard $\overline{C}$ as unordered.

Let a'_1 and a'_2 be the sizes of C_1 and $C_2 \setminus C_1$, respectively, which are prescribed by $\mathcal{P}$. There are at most $\ell d^{a'_1}$ choices for C_1 by (E5), then at most $a'_1(p+q)$ choices for $v \in \bigcup C_1$, d choices for $e_2 \in \mathcal{H}_1$ with $v \in e_2$, and $2^{a'_1} \beta'_x d^{a'_2-1}$ choices for $C_2 \subseteq \mathcal{H}_1$ with $e_2 \in C_2$ and $C_2 \cup e \in \mathcal{G}$.⁷

As in the previous section, suppose that $3 \leq t \leq m$ and we have already chosen $C_1, \dots, C_{t-1}$ such that their intersections are compatible with $\mathcal{P}$. Similarly, let a_t and a'_t be the desired sizes of $C_t \cap (C_1 \cup \dots \cup C_{t-1})$ and $C_t \setminus (C_1 \cup \dots \cup C_{t-1})$, respectively, as determined by $\mathcal{P}$. As before, we see that the number of choices for C_t is at most $2^s \Delta_{a_t,1}(\mathcal{G}^{(a_t+a'_t,1)}) \leq 2^s \ell d^{a'_t}$, in all cases, by (E5) and (E6). Hence, noting that $\sum_{t \in [m]} a'_t = s$, we obtain that $q(\mathcal{P}, e) \leq \Gamma \beta'_x d^s$. This in particular means that

$$f(\mathcal{H}_1) = d_x^{-1} \sum_{\mathcal{P} \in \mathcal{P}_{m,s,s'}^*} \sum_{e \in N_x} q(\mathcal{P}, e) \leq \Gamma^2 \beta'_x d^s.$$

Therefore, since $\beta_x \leq d^{-\varepsilon/3}$, and by the assumption that $\overline{w}_x^{m,s,s'}(\mathcal{H}_1) \geq d^{s-\varepsilon^3}$, we obtain that

$$f(\mathcal{H}_1) \leq d^{-\varepsilon/4} \overline{w}_x^{m,s,s'}(\mathcal{H}_1).$$

By the very same argument, but using instead the degree conditions on $\mathcal{C}$ and the bound from (5.2) on the number of $\varepsilon/2$ -conflict-sharing pairs $\{e, f\}$ given any fixed edge $e \in \mathcal{H}_1$, respectively, we obtain that $g(\mathcal{H}_1) \leq d^{-\varepsilon/2} \overline{w}_x^{m,s,s'}(\mathcal{H}_1)$ and $h(\mathcal{H}_1) \leq d^{-\varepsilon/3} \overline{w}_x^{m,s,s'}(\mathcal{H}_1)$. Hence the lower bound in (5.8) follows by subtracting these three bounds.

5.5.4. Useful estimates. We finish this section by computing two bounds which we will need in Section 5.6.2. Firstly, we bound the order of magnitude of our test functions, specifically

$$(5.9) \quad \overline{w}_x^{m,s,s'}(\mathcal{H}_1) \leq \Gamma^2 d^s.$$

Indeed, we use a similar counting argument to the previous sections. Fix $\mathcal{P} \in \mathcal{P}_{m,s,s'}^*$ and $e \in N_x$, and attempt to count the number of collections $\overline{C} = \{C_t\}_{t \in [m]}$ with $\mathcal{P}(\overline{C}) = \mathcal{P}$ and $C_t \cup e \in \mathcal{G}$ for every $t \in [m]$. For each $t \in [m]$, let a_t and a'_t be the sizes of $C_t \cap (C_1 \cup \dots \cup C_{t-1})$ and $C_t \setminus (C_1 \cup \dots \cup C_{t-1})$, respectively, as prescribed by $\mathcal{P}$. Suppose now that we have chosen sets $C_1, \dots, C_{t-1}$ compatible with $\mathcal{P}$. If $a_t = 0$ then we have at most $\ell d^{a'_t}$ choices for C_t by (E5). If $a'_t = 0$ then we have at most 2^s choices for C_t . Otherwise we have at most $2^s d^{a'_t-\varepsilon/3}$ choices for C_t by (E6). Since $\sum_{t \in [m]} a'_t = s$ by definition, we obtain in total at most Γd^s choices for the collection $\overline{C}$. Thus we may bound

$$\overline{w}_x^{m,s,s'}(\mathcal{H}_1) \leq d_x^{-1} \sum_{\mathcal{P} \in \mathcal{P}_{m,s,s'}^*} \sum_{e \in N_x} \Gamma d^s \leq \Gamma^2 d^s,$$

as required, recalling that $1/\Gamma \ll 1/i^*$.

Secondly, given $m \in [i^*]$ and $s \in [m\ell - 1]$, we claim further that

$$(5.10) \quad \hat{w}_x^{m,s}(E) := \sum_{s'=s+1}^{m\ell} \overline{w}_x^{m,s,s'}(E) \leq d^{s-\varepsilon/4}.$$

Indeed, consider following the argument above when $s < s'$, in which case there must exist some $t \in [m]$ with $a_t \neq 0 \neq a'_t$. We therefore obtain the improved bound

$$\overline{w}_x^{m,s,s'}(\mathcal{H}_1) \leq \Gamma^2 d^{s-\varepsilon/3},$$

from which the desired result follows.

5.6. Constructing the matching. We are now ready to construct the matching in two stages, as described previously.

⁷Note that $e_2 \notin C_1$, since we assumed that all conflicts in $\mathcal{D}$ are matchings.

5.6.1. *Obtaining $\mathcal{M}_1$.* We start by applying Theorem 4.2 to the hypergraph $\mathcal{H}'_1$ defined in Section 5.2. Since $\mathcal{H}'_1$ may have many more vertices than $\mathcal{H}_1$, we need to ensure that only a small proportion of the vertices in P are left uncovered. To do this, simply define another 1-uniform test function (for $\mathcal{H}'_1$)

$$w_1(e) := \mathbf{1}(e \in \mathcal{H}_1).$$

For any matching $\mathcal{N} \subseteq \mathcal{H}'_1$, the value $pw_1(\mathcal{N})$ is equal to the number of vertices of P covered by $\mathcal{N}$. Note that w_1 is a $(d, \varepsilon/10, \mathcal{C})$ -trackable ℓ' -test function since (H1) and the fact that $|P| \geq d^\varepsilon$ (recalling (S2)) imply that $w_1(\mathcal{H}_1) = |\mathcal{H}_1| > |P|d/2p > d^{1+\varepsilon/2}$, and (W2)–(W4) hold trivially. Observe also that for every other test function w that we have defined previously on subsets of $\mathcal{H}_1$, we may extend w to subsets of $\mathcal{H}'_1$ by setting $w(E) = 0$ for all E with $E \not\subseteq \mathcal{H}_1$, without affecting whether w is trackable or semi-trackable.

We may therefore apply Theorem 4.2 to the hypergraph $\mathcal{H}'_1$, the set of all test functions we have defined, and the conflict hypergraph $\mathcal{C}$, with ℓ' in place of ℓ , to obtain a matching $\mathcal{M}'_1 \subseteq \mathcal{H}'_1$. Since all of our test functions are zero on $\mathcal{H}'_1 \setminus \mathcal{H}_1$, this induces a $\mathcal{C}$ -free matching $\mathcal{M}_1 \subseteq \mathcal{H}_1$ such that, for each of the j -uniform test functions w which we have defined, we have $w(\mathcal{M}_1) = (1 \pm d^{-\varepsilon^3})d^{-j}w(\mathcal{H}_1)$ if w is trackable, and $w(\mathcal{M}_1) \leq d^{\varepsilon/4}$ if w is semi-trackable. In particular, applied to w_1 , this means that $|\mathcal{M}_1| = |\mathcal{M}'_1 \cap \mathcal{H}_1| \geq (1 - d^{-\varepsilon^3})d^{-1}|\mathcal{H}_1| \geq (1 - d^{-\varepsilon^3})(1 - d^{-\varepsilon})|P|/p$ by the degree condition (H1). Hence at most $(1 - (1 - d^{-\varepsilon^3})(1 - d^{-\varepsilon}))|P| \leq d^{-\varepsilon^3/2}|P|$ vertices of P are left uncovered by $\mathcal{M}_1$. Write this set as $P' := \{x_1, \dots, x_M\}$ for $M \leq d^{-\varepsilon^3}|P|$.

5.6.2. *Restricting to safe edges.* As discussed, in order to avoid conflicts in the case $j_2 = 1$, we restrict to a smaller set of safe edges for each vertex in P' when choosing $\mathcal{M}_2$. Recalling that $N_x = \{e \in \mathcal{H}_2 : x \in e\}$ and $d_x = |N_x|$, we show now that for each $x \in P'$, there exists a subset $N_x^s \subseteq N_x$ of safe edges such that if $C \cup e \in \mathcal{D}$ is a conflict with $C \in \binom{\mathcal{M}_1}{j_1}$ for some $j_1 \in [\ell]$ and $e \in N_x$, then $e \notin N_x^s$.

We use the inclusion-exclusion principle to bound the number of safe edges from below, showing that we can choose N_x^s of positive density; that is, we ensure that $|N_x^s| \geq \lambda d_x$, for some constant $\lambda > 0$ to be specified, so that restricting to N_x^s does not significantly limit our choice of edges for $\mathcal{M}_2$. To do this, we need to estimate the number of m -sets $\{C_1, \dots, C_m\}$ such that each $C_t \subseteq \mathcal{M}_1$ forms a conflict with the same edge e . This is possible because $\mathcal{M}_1$ is sufficiently pseudorandom, in an appropriate sense, which we show using the test functions $w_x^{m,s,s'}$, defined in Section 5.5. For the remainder of this section, we again write $\mathcal{G} = \bigcup_{j \in [\ell]} \mathcal{D}^{(j,1)}$ for simplicity, recalling our notation from Section 3.3. Fix $x \in P'$, and recall that $i^* \in \mathbb{N}$ satisfies $1/i^* \ll 1/\ell$.

For each $C \subseteq \mathcal{H}_1$, start by defining $B_C := \{e \in N_x : C \cup e \in \mathcal{G}\}$ to be the set of edges in $\mathcal{H}_2$ containing x which complete a conflict with C , and for each $m \in [i^*]$, recalling the definition of $E_i(\cdot)$ from Section 5.4, set

$$a_m := \sum_{e \in N_x} \binom{|E_1(\mathcal{G}_{[\emptyset,e]}) \cap 2^{\mathcal{M}_1}|}{m}$$

as the sum over all $e \in N_x$ of the numbers of m -sets of partial conflicts (of any sizes) in $\mathcal{M}_1$, all of which form a conflict with e . Define $N_x^s := \{e \in N_x : E_1(\mathcal{G}_{[\emptyset,e]}) \cap 2^{\mathcal{M}_1} = \emptyset\}$ to be the set of safe edges. We may assume that i^* is chosen to be odd, so by the inclusion-exclusion principle, we obtain that

$$(5.11) \quad |N_x^s| = \left| N_x \setminus \bigcup_{C \subseteq \mathcal{M}_1} B_C \right| \geq d_x - a_1 + a_2 - a_3 + \dots - a_{i^*}.$$

We now aim to show that this is suitably close to an exponential series, which guarantees that a constant proportion of the edges of N_x belong to N_x^s .

Fix $m \in [i^*]$ then, considering all of the possible sizes and intersections of a set of m conflicts, we see by the definition of $w_x^{m,s,s'}$ in Section 5.5.1 that

$$(5.12) \quad a_m = d_x \sum_{s' \in [m\ell]} \sum_{s \in [s']} w_x^{m,s,s'}(\mathcal{M}_1).$$

We first claim that

$$(5.13) \quad w_x^{m,s,s'}(\mathcal{M}_1) = (1 \pm d^{-\varepsilon^3/2}) d^{-s} \bar{w}_x^{m,s,s'}(\mathcal{H}_1) \pm d^{-\varepsilon^3/2}$$

for every $s' \in [m\ell]$ and $s \in [s']$. Indeed, we must consider three cases. Firstly, if $\beta_x^{-1} w_x^{m,s,s'}$ is semi-trackable, then Theorem 4.2 tells us that $w_x^{m,s,s'}(\mathcal{M}_1) \leq \beta_x d^{\varepsilon/4} \leq d^{-\varepsilon/12}$, recalling that $\beta_x \leq d^{-\varepsilon/3}$. Secondly, if $\beta_x^{-1} w_x^{m,s,s'}$ is trackable but $\bar{w}_x^{m,s,s'}(\mathcal{H}_1) < d^{s-\varepsilon^3}$, then we obtain that $w_x^{m,s,s'}(\mathcal{M}_1) = (1 \pm d^{-\varepsilon^3}) d^{-s} w_x^{m,s,s'}(\mathcal{H}_1) \leq 2d^{-s} \bar{w}_x^{m,s,s'}(\mathcal{H}_1) \leq 2d^{-\varepsilon^3}$. Since $d^{-s} \bar{w}_x^{m,s,s'}(\mathcal{H}_1) \leq d^{-\varepsilon^3}$ in both of these cases, it is certainly true that $w_x^{m,s,s'}(\mathcal{M}_1) = d^{-s} \bar{w}_x^{m,s,s'}(\mathcal{H}_1) \pm d^{-\varepsilon^3/2}$. Finally, if $\beta_x^{-1} w_x^{m,s,s'}$ is trackable and $\bar{w}_x^{m,s,s'}(\mathcal{H}_1) \geq d^{s-\varepsilon^3}$, then the estimate from Theorem 4.2, combined with (5.8), gives us that $w_x^{m,s,s'}(\mathcal{M}_1) = (1 \pm d^{-\varepsilon^3/2}) d^{-s} \bar{w}_x^{m,s,s'}(\mathcal{H}_1)$, so (5.13) holds in all cases. Plugging (5.13) into (5.12) and recalling (5.10), we obtain that

$$(5.14) \quad \begin{aligned} a_m &= (1 \pm d^{-\varepsilon^3/2}) d_x \sum_{s' \in [m\ell]} \sum_{s \in [s']} d^{-s} \bar{w}_x^{m,s,s'}(\mathcal{H}_1) \pm d^{-\varepsilon^3/2} d_x \\ &= (1 \pm d^{-\varepsilon^3/2}) d_x \sum_{s' \in [m\ell]} d^{-s'} \bar{w}_x^{m,s',s'}(\mathcal{H}_1) \pm d^{-\varepsilon^3/3} d_x. \end{aligned}$$

In order to approximate this value, for each $e \in N_x$ and $j \in [\ell]$, define $\gamma_{e,j} := d^{-j} |\mathcal{G}_{[\emptyset,e]}^{(j,1)}| \leq \ell$ (by (E5)). Then set $\gamma_e := \sum_{j \in [\ell]} \gamma_{e,j} \leq \ell^2$, as an estimate for the total number of partial conflicts completed by e , of any size, which we expect to appear in $\mathcal{M}_1$. For each $s' \in [m\ell]$, write $\mathcal{Q}_{m,s'}$ for the set of sequences $\mathbf{b} = (b_j)_{j \in [\ell]}$ with $\sum_{j \in [\ell]} b_j = m$ and $\sum_{j \in [\ell]} j b_j = s'$. Considering the number of ways to choose m conflicts containing e with the sum of sizes of $\mathcal{H}_1$ -parts equal to s' , we see that we may write

$$(5.15) \quad \sum_{s \in [s']} d_x \bar{w}_x^{m,s,s'}(\mathcal{H}_1) = \sum_{\mathbf{b} \in \mathcal{Q}_{m,s'}} \sum_{e \in N_x} \prod_{j \in [\ell]} \binom{|\mathcal{G}_{[\emptyset,e]}^{(j,1)}|}{b_j} = d^{s'} \sum_{e \in N_x} \sum_{\mathbf{b} \in \mathcal{Q}_{m,s'}} \prod_{j \in [\ell]} \frac{\gamma_{e,j}^{b_j}}{b_j!} \pm d^{s'-1/2} d_x,$$

where the error term is obtained using the fact that $\binom{n}{k} = n^k/k! \pm f(n)$ for a polynomial f of degree at most $k-1$, as well as the fact that $|\mathcal{G}_{[\emptyset,e]}^{(j,1)}| \leq \ell d^j$ by (E5). Recalling (5.9), we see that all summands with $s < s'$ contribute lower order terms to (5.15), so in fact

$$(5.16) \quad d_x \bar{w}_x^{m,s',s'}(\mathcal{H}_1) = d^{s'} \sum_{e \in N_x} \sum_{\mathbf{b} \in \mathcal{Q}_{m,s'}} \prod_{j \in [\ell]} \frac{1}{b_j!} \gamma_{e,j}^{b_j} \pm 2d^{s'-1/2} d_x.$$

Plugging (5.16) into (5.14) gives

$$(5.17) \quad a_m = (1 \pm d^{-\varepsilon^3/2}) \sum_{s' \in [m\ell]} \sum_{e \in N_x} \sum_{\mathbf{b} \in \mathcal{Q}_{m,s'}} \prod_{j \in [\ell]} \frac{\gamma_{e,j}^{b_j}}{b_j!} \pm d^{-\varepsilon^3/4} d_x.$$

Now observe by the multinomial theorem that

$$(5.18) \quad \gamma_e^m = \left(\sum_{j \in [\ell]} \gamma_{e,j} \right)^m = m! \sum_{s' \in [m\ell]} \sum_{\mathbf{b} \in \mathcal{Q}_{m,s'}} \prod_{j \in [\ell]} \frac{\gamma_{e,j}^{b_j}}{b_j!}.$$

Hence, plugging (5.18) into (5.17), we obtain the estimate

$$(5.19) \quad a_m = (1 \pm d^{-\varepsilon^3/2}) \sum_{e \in N_x} \frac{\gamma_e^m}{m!} \pm d^{-\varepsilon^3/4} d_x.$$

For $x \in \mathbb{R}$, define $S(x) = \sum_{m=0}^{i^*} (-x)^m/m!$. Now plugging (5.19) into (5.11) gives

$$(5.20) \quad |N_x^s| \geq \sum_{e \in N_x} 1 - \gamma_e + \frac{1}{2!} \gamma_e^2 - \cdots - \frac{1}{i^*!} \gamma_e^{i^*} \pm i^* d^{-\varepsilon^3/4} d_x \geq \sum_{e \in N_x} S(\gamma_e) - d^{-\varepsilon^3/5} d_x.$$

To bound $\sum_{e \in N_x} S(\gamma_e)$ from below, let $\delta := \exp(-\ell^2)/3 > 0$, then recalling that $1/i^* \ll 1/\ell$, we see that for every $0 \leq x \leq \ell^2$, we have $S(x) = \exp(-x) \pm \delta$. The choice of such an i^* , depending only upon ℓ , is possible by the uniform convergence of the exponential series on the interval $[0, \ell^2]$. Then, since $0 \leq \gamma_e \leq \ell^2$ for every $e \in N_x$, we obtain

$$\sum_{e \in N_x} S(\gamma_e) \geq \sum_{e \in N_x} \exp(-\gamma_e) - d_x \delta \geq d_x \exp\left(-\frac{1}{d_x} \sum_{e \in N_x} \gamma_e\right) - d_x \delta \geq d_x(\exp(-\ell^2) - \delta)$$

by convexity of the exponential function and the bound on γ_e . Substituting this back into (5.20), we see that $|N_x^s| \geq d_x(\exp(-\ell^2) - 2\delta) \geq d_x \exp(-\ell^2)/3$. Hence we may take $\lambda = \lambda_\ell := \exp(-\ell^2)/3 > 0$.

5.6.3. Choosing $\mathcal{M}_2$. We may now proceed to choose the edges of $\mathcal{M}_2$ randomly from the sets we have defined. Recall that $P' = \{x_1, \dots, x_M\}$ is the subset of P not yet covered by $\mathcal{M}_1$, and define $\mathcal{H}'_2 := \mathcal{H}_2[P' \cup R]$. For each $i \in [M]$, choose an edge e_i uniformly at random from the set $N_{x_i}^s$, so each possible edge $e \in N_{x_i}^s$ is taken with probability

$$(5.21) \quad \mathbb{P}[e_i = e] \leq (\lambda d_{\mathcal{H}_2}(x_i))^{-1}.$$

Recall also that (H3') and (H4') together imply that $d_{\mathcal{H}_2}(x_i) \geq d^\varepsilon$, so $|N_{x_i}^s| \geq \lambda d^\varepsilon > 0$. Set $\mathcal{M}_2 := \{e_1, \dots, e_M\}$ and take $\mathcal{M} := \mathcal{M}_1 \cup \mathcal{M}_2$ to be the combination of our matching in $\mathcal{H}_1$ with these new edges chosen from $\mathcal{H}_2$. We now use Lemma 4.3 to ensure that $\mathcal{M}$ is $\mathcal{D}$ -free; recall that by excluding the conflicts added in Section 5.3, this will also ensure that $\mathcal{M}_2$ is a matching.

For each $j_2 \in [2, \ell]$, define bad events $B_D := \{D \subseteq \mathcal{M}_2\}$ for each $D \in \binom{\mathcal{H}'_2}{j_2}$ which could appear as the $\mathcal{H}_2$ -part of a conflict in $\mathcal{D}^{(j_1, j_2)}$ for some $j_1 \in [0, \ell]$, that is $D \in E_2(\mathcal{D}_{[C, \emptyset]}^{(j_1, j_2)})$ for some $C \in \binom{\mathcal{M}_1}{j_1}$.⁸ Let $\mathfrak{A}$ be the set of all such events, for any (j_1, j_2) , which we aim to avoid. Note that, for each event $B_D \in \mathfrak{A}$, we have $\mathbb{P}[B_D] \leq \lambda^{-1} d^{-\varepsilon} < 1/2$, so in order to apply Lemma 4.3, it suffices to show that we also have $\sum_{A \in \mathfrak{B}(D)} \mathbb{P}[A] \leq 1/4$, where $\mathfrak{B}(D) := \{B_{D'} \in \mathfrak{A} : D' \in E_2(\mathcal{D}_x) \text{ for some } x \in V_P(D)\}$, recalling the definition of $V_P(\cdot)$ from Section 4.3; since the event B_D depends only upon the edge choices for the vertices $x \in V_P(D)$, it is mutually independent from the set of all events $B_{D'}$ for which $V_P(D) \cap V_P(D') = \emptyset$.

Now fix some $j_1 \in [0, \ell]$, $j_2 \in [2, \ell]$, and $x \in P'$, and again write $\mathcal{G} := \mathcal{D}^{(j_1, j_2)}$ for ease of notation. Further write $\mathcal{G}^{\mathcal{M}_1} := \{C \cup D \in \mathcal{G} : C \subseteq \mathcal{M}_1 \text{ and } D \subseteq \mathcal{H}'_2\}$. Because $|V_P(D)| = j_2 \leq \ell$, and there are at most ℓ^2 choices for j_1 and j_2 , it is (more than) sufficient to show that

$$(5.22) \quad \sum_{D \in E_2(\mathcal{G}_x^{\mathcal{M}_1})} \mathbb{P}[D \subseteq \mathcal{M}_2] \leq d^{-\varepsilon^3/3}.$$

Given $C \in \binom{\mathcal{H}_1}{j_1}$, say that an $\mathcal{H}_2$ -part $D \in E_2(\mathcal{G}_{[C], x})$ is *blocked* if $V_P(D) \not\subseteq P'$, that is some vertex $y \in V_P(D)$ is already covered by $\mathcal{M}_1$. Note that such conflicts can be ignored as they will never be present in $\mathcal{M}$, because no edge containing y is chosen in $\mathcal{M}_2$; the remainder of the proof is therefore concerned with bounding *unblocked* conflicts. We define $\mathcal{B}(C, x) := \{D \in E_2(\mathcal{G}_{[C], x}) : D \text{ is blocked}\}$ and $\mathcal{U}(C, x) := \{D \in E_2(\mathcal{G}_{[C], x}) : D \text{ is not blocked}\}$. Then by definition

$$(5.23) \quad E_2(\mathcal{G}_x^{\mathcal{M}_1}) = \bigcup_{C \in \binom{\mathcal{M}_1}{j_1}} \mathcal{U}(C, x).$$

Furthermore, observe that by (5.21), and the independence of edge choices for distinct vertices of P' , we have

$$(5.24) \quad \mathbb{P}[D \subseteq \mathcal{M}_2] \leq \prod_{y \in V_P(D)} (\lambda d_{\mathcal{H}_2}(y))^{-1} = \lambda^{-j_2} A(D)$$

⁸Recall here our notation from Sections 3.3, 4.3, and 5.4.

for any (unblocked) $D \in \binom{\mathcal{H}'_2}{j_2}$. Hence to show (5.22), using (5.23) and (5.24), it is enough to prove that

$$(5.25) \quad \sum_{C \in \binom{\mathcal{M}_1}{j_1}} A(\mathcal{U}(C, x)) \leq d^{-\varepsilon^3/3}.$$

We now prove (5.25) using the test functions we have defined, as discussed previously.

5.6.4. Bounding unblocked conflicts. We consider first the case that the function $\alpha_x^{-1}w_x$ is $(d, \varepsilon/10, \mathcal{C})$ -semi-trackable, so Theorem 4.2 tells us that $\alpha_x^{-1}w_x(\mathcal{M}_1) \leq d^{\varepsilon/4}$. By the definition of w_x and the bound $\alpha_x \leq d^{-\varepsilon/3}$, this means that

$$\sum_{C \in \binom{\mathcal{M}_1}{j_1}} A(\mathcal{G}_{[C],x}) = w_x(\mathcal{M}_1) \leq \alpha_x d^{\varepsilon/4} \leq d^{-\varepsilon/12},$$

which in particular is sufficient for (5.25).

Assume instead that both of the functions $\alpha_x^{-1}w_x$ and $\alpha_x^{-1}w'_x$ are $(d, \varepsilon/10, \mathcal{C})$ -trackable. In order to obtain (5.25), we rewrite

$$(5.26) \quad \sum_{C \in \binom{\mathcal{M}_1}{j_1}} A(\mathcal{U}(C, x)) = \sum_{C \in \binom{\mathcal{M}_1}{j_1}} A(\mathcal{G}_{[C],x}) - \sum_{C \in \binom{\mathcal{M}_1}{j_1}} A(\mathcal{B}(C, x)),$$

and use w_x and w'_x to estimate the two sums respectively.

Firstly, by definition and the conclusion of Theorem 4.2, we have that

$$(5.27) \quad \sum_{C \in \binom{\mathcal{M}_1}{j_1}} A(\mathcal{G}_{[C],x}) = w_x(\mathcal{M}_1) = (1 \pm d^{-\varepsilon^3})d^{-j_1}w_x(\mathcal{H}_1) \leq (1 + d^{-\varepsilon^3})d^{-j_1}A(\mathcal{G}_x).$$

To obtain a lower bound for the second term in (5.26), start by observing that, given $D \in E_2(\mathcal{G}_x)$, we have $|V_P(D) \setminus P'| \leq (j_2 - 1)\mathbb{1}(V_P(D) \setminus P' \neq \emptyset) = (j_2 - 1)\mathbb{1}(D \text{ blocked})$. Therefore,

$$\begin{aligned} \sum_{C \in \binom{\mathcal{M}_1}{j_1}} A(\mathcal{B}(C, x)) &= \sum_{C \in \binom{\mathcal{M}_1}{j_1}} \sum_{D \in E_2(\mathcal{G}_{[C],x})} A(D) \mathbb{1}(D \text{ blocked}) \\ &\geq \frac{1}{j_2 - 1} \sum_{C \in \binom{\mathcal{M}_1}{j_1}} \sum_{D \in E_2(\mathcal{G}_{[C],x})} A(D) |V_P(D) \setminus P'| \\ &= \frac{1}{j_2 - 1} \sum_{C \in \binom{\mathcal{M}_1}{j_1}} \sum_{D \in E_2(\mathcal{G}_{[C],x})} A(D) \sum_{y \in V_P(D) \setminus \{x\}} \sum_{e \in \mathcal{M}_1} \mathbb{1}(y \in e) \\ (5.27) \quad &= \frac{1}{j_2 - 1} w'_x(\mathcal{M}_1). \end{aligned}$$

To see the penultimate equality, note that $y \in P \setminus P'$ if and only if there is exactly one edge $e \in \mathcal{M}_1$ containing y . For the final equality, recall the definition of w'_x in Section 5.4.1, and note further that if $C \cup e \subseteq \mathcal{M}_1$, then $C \cup e$ cannot contain any conflict-sharing pair or conflict from $\mathcal{C}$.

Now by Theorem 4.2 and the estimate (5.3), we see that

$$(5.28) \quad w'_x(\mathcal{M}_1) = (1 \pm d^{-\varepsilon^3})d^{-j_1-1}w'_x(\mathcal{H}_1) \geq (j_2 - 1)(1 - d^{-\varepsilon^3/2})d^{-j_1}w_x(\mathcal{H}_1).$$

Hence combining (5.27) and (5.28) we obtain the bound

$$(5.29) \quad \sum_{C \in \binom{\mathcal{M}_1}{j_1}} A(\mathcal{B}(C, x)) \geq (1 - d^{-\varepsilon^3/2})d^{-j_1}A(\mathcal{G}_x).$$

We finish by plugging (5.29) and (5.27) back into (5.26) to obtain

$$\sum_{C \in \binom{\mathcal{M}_1}{j_1}} A(\mathcal{U}(C, x)) \leq 2d^{-\varepsilon^3/2}d^{-j_1}A(\mathcal{G}_x) \leq d^{-\varepsilon^3/3},$$

by (E2), as required for (5.25). Note that all of the arguments made here still work for $j_1 = 0$, when w_x is defined simply on the empty set.

6. ACKNOWLEDGMENTS.

We are grateful to Michelle Delcourt and Luke Postle for helpful comments and for informing us about some references. We would also like to thank the referees for their extensive and helpful comments.

GLOSSARY

- $\mathcal{C}$ -free:** contains no conflict from $\mathcal{C}$. 2
- $d_{\mathcal{G}}(u, v)$: codegree of vertices u and v in hypergraph $\mathcal{G}$. 3
- $\Delta_V(\mathcal{G})$: maximum degree of all vertices in $V \subseteq V(\mathcal{G})$. 3
- $\mathcal{G}^{(j)}$: subhypergraph of $\mathcal{G}$ consisting of edges of size j . 3
- $N_{\mathcal{C}}^{(2)}(e)$: set of edges forming a conflict of size two with e . 5
- (d, ℓ, ε) -**bounded**: conflict hypergraph satisfying (C1)–(C5). 5
- $\mathcal{D}^{(j_1, j_2)}$: conflicts consisting of j_1, j_2 edges from $\mathcal{H}_1, \mathcal{H}_2$, respectively. 5
- $\Delta_{j_1, j_2}'(\mathcal{D})$: maximum codegree among edge sets consisting of j_1, j_2 edges from $\mathcal{H}_1, \mathcal{H}_2$, respectively. 5
- $\mathcal{D}_x$: conflicts containing vertex x in their $\mathcal{H}_2$ -part. 5
- $\mathcal{D}_{x, y}$: conflicts containing both x and y in their $\mathcal{H}_2$ -part. 5
- (d, ℓ, ε) -**simply-bounded**: conflict hypergraph satisfying (D1)–(D4). 5
- j -uniform ℓ -test function**: $[0, \ell]$ -valued function on sets of j edges which is non-zero only for matchings. 6
- $\mathcal{H}_{-v}$: link hypergraph (all sets whose union with v form an edge in $\mathcal{H}$). 6
- $(\varepsilon, \mathcal{C})$ -**conflict-sharing pair**: two edges with large link hypergraph in $\mathcal{C}$. 6
- $(d, \varepsilon, \mathcal{C})$ -**trackable**: test function satisfying (W1)–(W4). 6
- $(d, \varepsilon, \mathcal{C})$ -**semi-trackable**: test function satisfying (W1*), (W2*), (W3), (W4). 7
- $V_P(E)$: all vertices of P covered by $\mathcal{H}_2$ -part of E . 8
- $A(E)$: unavailability of edge set E . 8
- $\mathcal{G}_{[C, D]}$: conflicts in $\mathcal{G}$ containing C, D in their $\mathcal{H}_1, \mathcal{H}_2$ -parts, respectively. 8
- $\Delta_{j_1, j_2}^A(\mathcal{G})$: maximum codegree weighted by unavailability. 8
- $(d, \ell, \varepsilon, \delta)$ -**mixed-bounded**: conflict hypergraph satisfying (E1)–(E6). 8
- $A(v)$: unavailability of vertex v . 9
- ℓ', Γ, i^* : constants used in proof. 9
- $E_i(\mathcal{G})$: $\mathcal{H}_i$ -parts of conflicts in $\mathcal{G}$. 11
- $\mathcal{G}_{[C]}$: conflicts of $\mathcal{G}$ containing edge set C in their $\mathcal{H}_1$ -part. 11
- $\mathcal{G}_{[C], x}$: conflicts containing edge set C and vertex x in their $\mathcal{H}_1, \mathcal{H}_2$ -parts, respectively. 11
- $\mathcal{G}_{[C], x, y}$: conflicts containing edge set C and vertices x, y in their $\mathcal{H}_1, \mathcal{H}_2$ -parts, respectively. 11
- w_x : test function used to count all potential conflicts containing x . 11
- testable**: edge set which is a conflict-free matching with no conflict-sharing pairs. 11
- w'_x : test function used to count blocked potential conflicts containing x . 11
- $b_{C, y}$: edges containing y which could block conflicts containing C . 11
- α_x : scaling constant for test functions w_x, w'_x . 13
- $\mathcal{Z}_E^{(j)}$: all edge sets of size j containing edge set E . 13
- N_x : edges in $\mathcal{H}_2$ containing vertex x . 14
- d_x : size of N_x . 14
- $\mathcal{P}(\overline{C})$: encoding of all intersections of a collection $\overline{C}$ of edge sets. 14
- $\mathcal{P}_{m, s, s'}^*$: all possible $\mathcal{P}(\overline{C})$ for m sets with sum of sizes s' and union size s . 14
- $\overline{w}_x^{m, s, s'}$: function used to count m -sets of potential conflicts with common $\mathcal{H}_2$ -part of size 1. 14
- $w_x^{m, s, s'}$: restriction of $\overline{w}_x^{m, s, s'}$ to testable sets. 14

- β_x : scaling constant for test functions $\overline{w}_x^{m,s,s'}, w_x^{m,s,s'}$. 14
 w_1 : test function used to count vertices in P left uncovered by $\mathcal{M}_1$. 17
 N_x^s : edges in $\mathcal{H}_2$ containing x which do not form a conflict with $j_2 = 1$. 17

REFERENCES

1. Deepak Bal, Patrick Bennett, Emily Heath, and Shira Zerbib, *Generalized ramsey numbers of cycles, paths, and hypergraphs*, arXiv: 2405.15904 (2024).
2. Patrick Bennett, Ryan Cushman, Andrzej Dudek, and Paweł Prałat, *The Erdős-Gyárfás function $f(n, 4, 5) = \frac{5}{6}n + o(n)$ – so Gyárfás was right*, Journal of Combinatorial Theory, Series B **169** (2024), 253–297.
3. Patrick Bennett, Emily Heath, and Shira Zerbib, *Edge-coloring a graph g so that every copy of a graph h has an odd color class*, arXiv:2307.01314 (2023).
4. William Brown, Paul Erdős, and Vera Sós, *Some extremal problems on r -graphs*, New directions in the theory of graphs (1973), 53–63.
5. Michelle Delcourt, Tom Kelly, and Luke Postle, *Clique decompositions in random graphs via refined absorption*, arXiv: 2402.17857 (2024).
6. ———, *Thresholds for $(n, q, 2)$ -steiner systems via refined absorption*, arXiv: 2402.17858 (2024).
7. Michelle Delcourt and Luke Postle, *Finding an almost perfect matching in a hypergraph avoiding forbidden submatchings*, arXiv: 2204.08981 (2022).
8. ———, *Proof of the high girth existence conjecture via refined absorption*, arXiv: 2402.17856 (2024).
9. ———, *Refined absorption: A new proof of the existence conjecture*, arXiv: 2402.17855 (2024).
10. Paul Erdős and Lovász László, *Problems and results on 3-chromatic hypergraphs and some related questions*, Coll Math Soc J Bolyai **10** (1974).
11. Peter Frankl and Vojtěch Rödl, *Near perfect coverings in graphs and hypergraphs*, European Journal of Combinatorics **6** (1985), 317–326.
12. Stefan Glock, Felix Joos, Jaehoon Kim, Marcus Kühn, and Lyuben Lichev, *Conflict-free hypergraph matchings*, Journal of the London Mathematical Society **109** (2024), e12899.
13. Enrique Gomez-Leos, Emily Heath, Alex Parker, Coy Schwieler, and Shira Zerbib, *New bounds on $f(n, 5, 8)$* , arXiv: 2308.16365 (2023).
14. Felix Joos and Dhruv Mubayi, *Ramsey theory constructions from hypergraph matchings*, to appear in Proceedings of the American Mathematical Society.
15. Felix Joos, Dhruv Mubayi, and Zak Smith, *Conflict-free hypergraph matchings and coverings*, arXiv:2407.18144 (2024).
16. Andrew Lane and Natasha Morrison, *Generalized ramsey numbers via conflict-free hypergraph matchings*, arXiv: 2405.16653 (2024).
17. Nicholas Pippenger and Joel Spencer, *Asymptotic behavior of the chromatic index for hypergraphs*, Journal of Combinatorial Theory, Series A **51** (1989), 24–42.
18. Joel Spencer, *Asymptotic lower bounds for ramsey functions*, Discrete Mathematics **20** (1977), 69–76.